1 Euphrasia rostkoviana Hayne

E. campestris auct. angl. non Jord, E. campestris var. neglecta Bucknall., E. fennica sensu Bucknall non Kihlm. ex Mela, E. hirtella auct. angl. non Jord. ex Reut. pro parte, E. officinalis subsp. pratensis Fr., E. officinalis subsp. pratensis Schubl & G. Martens, E. officinalis subsp. rostkoviana (Hayne) F. Towns., E. officinalis subsp. rostkoviana forma tenuis F. Towns., E. rostkoviana var. obscura Pugsley, E. rostkoviana subsp. rostkoviana Hayne

Rostkov's Eyebright


Stems 10-40(50) cm, green or slightly reddish, erect. Cauline internodes shorter to 3 times as long as leaves, lower floral internodes shorter to 4 times as long. Branches 1-12 pairs, slender, ascending or erect, often branched again. Leaves bright green and often conspicuously veined below. Cauline leaves 3-17mm, suborbicular or broadly to narrowly ovate, obtuse at the apex with 1-7 pairs of obtuse to subacute teeth, not much longer than wide, sessile or rarely shortly petiolate, with a truncate or rounded base. Lower floral leaves 6-15mm, generally smaller than the upper cauline, deltoid or broadly ovate to oblong-ovate, obtuse to acute at apex with 4-9 pairs of subacute to acuminate teeth, not much longer than wide, the basal pair patent or retrorse, sessile with a rounded, truncate or cordate base. All (save sometimes the lowest leaves) with more or less dense, long multicellular glandular hairs with the stalk 10-12 times as long as the gland. Lowest flower at node 6-10(14). Corolla (6.5)8-12 mm, white or with a purple or lilac upper lip. Calyx green, teeth narrowly or very narrowly deltoid, acuminate



Euphrasia rostkoviana.



Euphrasia rostkoviana.



Euphrasia rostkoviana.