

Reginald L. Hine F.S.A.

William Bury
Letchworth

N.B. Author's copy with numerous additions
in his hand & that of his wife.



Index Boletorum. Peruvia.

ereus	313	Acetabulum	325.
bovinus	311	Atra	330.
cyanescens	312	bicolor	329.
elegans	314	cerulea	330
flabelliformis	318	calyculus	327.
fomentarius	318	campanulata	331.
frondosus	314	cinernea	331.
gregarius	312	coccinea	326
hepaticus	316	cochleata	327.
hispidus	317	cornucopioides	324
ignarius	318	cyathoides	326
lachrymans	316	epidendra	325
luteus	313	fructigena	327.
medulla panis	315	fulva	328
humulatus	312	hispida	329.
obliquatus	314	Lewis	331.
olivaceus	312	lenticularis	330.
piperatus	313	leporina	329.
rugosus	315	marginata	327.
Salicinus	316	nivea	325.
squamulosus	314	Ochroleuca	324.
suberosus	315	papillaria	328
suaveolens	316	pietri	332.
sulphureus	317	polymorpha	330.
unicolor	317	punctata	325
versicolor.	317	punicea	328
		scutellata	328
		Spadicea	331.
		Stercoraria	328
		striata	331.
		tremelloidea	329.
		tuberosa	326
		viridis.	330.

Clavariae.

Aculeiformis	334
Anthocephala	333
Cinerea	333
Coralloides	332
Conica	334
digitalis	333
fastigiata	332
gyrans	334
Hypoxylon	333
militaris	334
Muscoides	333
pistillaris	332
Ophioglossoides	332.

Lycoperda.

Ardosiaceum	337
Carpobolus	336
Cinereum	337
Epidendrum	337
fragile	338
Gossypium	337
inhatum	338
pedunculatum	336
Protens	336
recollegens	336
stellatum	335
verrucosum	337

Sphaeria.

acuta	341
aggregata	340
atra	342
cirrhata	342
clavata	342
corticalis	340
deprepar	340
echinata	341
fraxinea	339
globularis	341
gregaria	339
lycoperdonides	339
mammosa	340
mori	340
obducta	342
ostreacea	341
sanguinea	340
stigma	341
sulcata	341
tremelloides	339

Trichia

cinerea	343
denudata	343
furfuracea	344
fuscoater	345
globulifera	344
lichenoides	345
minuta	344
nuda	343
olivacea	344
pumiliformis	345
recentita	345
terribilis	345
typiformis	343

Mucoræ.

aurantius	348
botrytis	347
cespitosus	347
casens	348
crustaceus	347
chrysospermus	347
embolus	346
erisyphe	348
glaucus	346
granulosus	348
ligni fractus	347
Mucedo	346
roridus	346
urceolatus	346

Tremella.

albida	270.
arborea	271.
carbonaria	273.
cinnabarinæ	272.
deliquescent	272.
glandulosa	272.
mesenterica	271.
posticæ	271.
nigricans	272.
violacea	271.

Reticularia.

ovata	338
segetum	339
Septica	338.

Byssæ.

auræa	276
candolanis	277
candida	277
flos aquæ	276
fulva	277
incana	278
phosphorea	276
Septica	277
velutina	276.

Uloæ.

crispa	273.
intestinalis	273.
pruniformis	274.

Conferæ.

amphibia	275
bullosa	274
canalicularis	274
gelatinosa	275.
glomerata	275
reticulata	275
riularis	274

Merulizæ.

caryophyllous	278
chantarellus	279
membraneous	279
squamula	278.

Hydnæ.

- auriscalpium* 319
barba Jovis 319
vibracatum 319
sublamellosum 319.

Helvella.

- agariciformis* 320
esculenta 321
gelatinosa 321
nitra 320
Velutina 320
spatulata 321.

Auricularia

- corticalis* 323
ferruginea 322
phylacteris 323
papyrina 322
reflexa 322
tabacina 323
tremelloides 322

Phallus.

- caninus* 324
impudicus 328.

Tubera.

- cibarium* 335
radicatum 335

Squisetæ.

- arvense* 222
fluviatile 222
hyemale 222
limosum 222
palustre 222
sylvaticum 221.

Osmunda.

- lunaria* 223
regalis 224
Spicant. 224

Ophioglossum.

- vulgatum* 223.

Pteris.

- agilina* 224.

Asplenia.

- adianthum nigrum* 225
ruta muraria 225
scolopendrium 225
trichomanes 225.

Polypodia.

- aculeatum* 227
cristatum 226
filix fœminæ 226
filix mas 226
thelypteris 227
vulgare 226

Lycopodiæ.

- clavatum* 228
mundatum 228.

Sphagna.

- arboreum* 229
palustre 228.

Phasca.

- alternifolium* 229
cuspidatum 229
curvicolle 230
piliferum 230
sphaero carpon 230
subulatum 229.

Fontinalides

- antipyretica* 231
minor. 231.

Splachnum

- ampullaceum* 231.

Polytricha.

- commune* 232
juniperinum 232
subrotundum 232.

Marchantia

- polymorphia* 255.

Supplementary.continued.

- ~~X Clavaria~~ ~~epiphylla~~ D. 293
~~X Clavaria~~ ~~coccinea~~ D. 294
~~X Sphagnum~~ ~~communis~~ D. 295.
~~X~~ ~~subcorticalis~~ D. 296.
~~X~~ ~~spinosa~~ D. 299.

- ~~X Bryum~~ ~~Dicksonii~~
~~X~~ ~~Lichen~~ ~~trebella~~
~~X~~ ~~Phascum~~ ~~bryoides~~
~~X~~ ~~Lichen~~ ~~arenarius~~
~~X~~ ~~Lichen~~ ~~punctiformis~~
~~X~~ ~~Bryum~~ ~~Fennæ~~

- Hypnum* *curtipendulum*

- Jungernaria* *lanceolata*,
~~X~~ *Bryum* *infuscans*,
~~X~~ *annulatum*.

- Lichen* *inclusive*,
~~X~~ *corticatus*.

- ~~X~~ *luteus*.

- ~~X~~ *tremelloides* +
~~X~~ *fallax* +

- juniperinum* 232
~~X~~ *fragile* +

- sericeum* +

- Hypnum* *denticulatum* +

- ~~Erodium~~ ~~moschatum~~.

- ~~Moss~~ ~~muraria~~.

- ~~Celid~~ ~~Rufobrunneum~~.

Bedstraw multiflorum
Glechoma glauca
Geum robustum
Caleptra verrucosa
Heather wood Between Haws & Hitchin.
Cypripedium
Lathyrus
Linton &
Barton.

Pennisetum muralis
 near Hitchin on the road
 to London.

Plants already collected
for a supplement. Nov? 1899.

- Eriophorum angustifolium* Lichen
x — coarctatus
- Habenaria nodosa* x — hematoma.
- Festuca tenuifolia* x — lychnis
- ~~Primula primuloides~~ x — cerinus
- Euphorbia Cyparissias* x — sinuosus — 36
- Rosa villosa* x — crenulatus.
- ~~Heuchera villosa~~ x — palmatus
- x — sativa x — hispidus. 2
- x — subspicata
- x — gentilis
- ~~Funaria heterophylla~~ x — Agaricus lobatus.
- Turritis glabra* x — opacus.
- Lactaria scrobicularia* x — Boletus biennis.
- Comfreylla decolorans* x — Peziza junci.
- Salix amygdalina* x — miclinans
- x — cuneata
- x — acuminata
- x — Sphaeria miniata. 40 —
- Ruscus aculeatus* x — Lichen dendriticus
- x — *Taxus baccata* x — pulverulentus.
- Crypt. x — rupestris.
- Phasium hispida* x — Herniaria muscigenus.
- x — muticum.
- x — Bryum rubens.
- Marchantia conica* x — Agaricus miniatus
- Lichen hebraicus* x — polygrammus.
- x — graniformis 20
- x — confertus
- x — repellatus
- x — chartaceus
- x — emarginatus
- x — turfosus
- x — sultans.
- x — Herniaria
- x — Peziza aurea
- x — Clavaria gracilis. 50
- x — Auricularia pulverulenta
- x — Trichia sphaeroccephala.
- x — Boletus pectinatus
- x — Jungermannia circinata

<i>Lichen linitatus.</i>	56. <i>Agaricus mutabilis</i> ^{Coprophilus.}
<i>Sphaera minuta</i> corticata.	* <i>Agaricus pectinatus.</i>
<i>Boletus impuber</i>	<i>Agaricus congregatus.</i> 8.F. 261.
<i>Peruvia hydnoides.</i>	* <i>Ag. Stercorarius</i> 8.F. 262.
<i>Lichen glabratus</i> X <i>Dickson</i> unpublished-60.	* <i>Ag. nigripes.</i> 8.F. 263.
<i>Boletus albidus</i>	<i>Lycoperdon aurantiacum</i> 8.F. 268.
* <i>ferrugineus</i>	* <i>Sphoria spiculifera</i> 8.F. 274
* <i>pelloporus-</i>	* <i>Agaricus ciliatus</i>
* <i>rubeotarius.</i> X	* <i>deliciosus?</i>
	* <i>Sphoria papillosa</i> 8.F.
<i>Sphoria decorticans</i>	8.F. <i>Agaricus hybridus</i>
* <i>tentaculata</i>	8.F. <i>Ag. scaber</i> 8.F.
<i>Agaricus vaginalis</i> 69. 80. <i>Lichen Sphaeroides</i> ^{Dickson.}	
<i>Chama hispida</i>	<i>Agaricus acicula</i>
<i>Bromus diandrus</i>	E.F. 282.
<i>Rosa canina</i>	<i>Agaricus monstrosus</i> 8.F. 283.
<i>Persicaria amplexicaulis.</i>	<i>Ag. radicans.</i> 8.F. 287.
<i>Scrophularia agricola.</i>	
<i>Agaricus vaginalis.</i> 80.	<i>Boletus hybridus.</i>
<i>Scrophularia viridis</i>	8.F. 289.
<i>Robus corymbiferus</i>	* <i>Agaricus armillatus</i> 8.F. 291.
<i>Leontopodium canescens</i>	<i>Lycoperdon equinum</i> 8.F. 292.
<i>Valeriana tuberosa</i>	* <i>Sphoria crenata</i>
<i>Osmunda cinnamomea</i>	* <i>tuberosa.</i>
<i>Vitis vulpina.</i>	<i>Ricium Paceratum.</i>

F L O R A
BEDFORDIENSIS,
COMPREHENDING SUCH PLANTS
AS GROW WILD IN THE COUNTY OF
BEDFORD,
ARRANGED
ACCORDING TO THE SYSTEM OF LINNÆUS,
WITH OCCASIONAL REMARKS.

B Y
CHARLES ABBOT, M. A. F. L. S.
CHAPLAIN TO THE
RIGHT HON. THE MARQUIS OF TWEEDALE,
VICAR OF OAKLEY RAYNES IN BEDFORDSHIRE,
AND LATE FELLOW OF NEW COLLEGE, OXFORD.

"Full many a gem of purest ray serene
 The dark unfathom'd caves of Ocean bear:
"Full many a flower is born to blush unseen,
 And waste it's sweetness on the desert Air."
Gray's Elegy.

Bedford:
PRINTED AND SOLD BY W. SMITH;
MAY BE HAD OF THE BOOKSELLERS IN OXFORD
& CAMBRIDGE, AND OF G. G. & J. ROBINSONS, LONDON.

1798.

Epistle Dedicatory.

To the Queen.

M A D A M,

THE condescension manifested by YOUR MAJESTY in permitting your name to adorn and sanction the pages of the following Work, cannot but inspire some degree of confidence in the heart of the Author, conscious as he is of the inadequate value of the offering. — To the august Patroness of the "H O R T U S K E W E N S I S,"

iv EPISTLE DEDICATORY.

whose scientific researches have justly obtained for her the character of the first female Botanist in the wide circle of the British Dominions, a character still farther embellished by the display of every domestic and relative Virtue, this Volume, chiefly designed for the amusement and instruction of that sex, which boasts YOUR MAJESTY as it's most distinguished Ornament, is humbly dedicated by

MADAM, Your MAJESTY's

Most devoted,

Most dutiful

And obedient Servant,

Charles Abbot.

BEDFORD, Nov. 20th. 1798.

PREFACE

WHEN, in pursuit of his favorite Amusement, the Botanist ranges with eager step amid the wild scenery of nature to pluck some simple flowret that has hitherto escaped observation, the World, disdainful of his humble Toil, marks him as a sort of phænomenon whose "extravagant and erring Spirit" wanders from the common track of human Life without any rational end or object. — But whatever branch of Science is the source of entertainment or instruction has it's use and importance: and the study of Botany exhibits to our view a succession of wonders, which tend to enlarge our knowledge and to heighten our admiration of the works of nature, and more especially of the Power Wisdom and Goodness of their omnipotent Author:

"These are thy glorious Works, Parent of Good"!

Such reflections ought surely to awe the cavalier into silence, whilst they induce the candid to spread the broad shield of public protection over the labours of the Botanist. — Impressed with

with these sentiments the fair daughters of Albion have evinced a zeal and ardour in Botanical researches which have not only done the highest honor to themselves, but have eminently contributed to rescue these pursuits from unmerited reproach, to elevate them into reputation, and to impart to them, if not a superior value, at least a superior currency and fashion. — That such excellence should have been attained in this branch of science by so many of the female sex, notwithstanding the disadvantages they labour under from the want of scholastic and technical instruction, is a convincing proof of the liberality with which Nature has endowed the female mind: and how little reason there is to suppose that their intellectual are from any other cause than want of cultivation, in any degree inferior to their personal accomplishments. In the work now before the public, although the technical terms are in a great measure softened and Anglicized, the Author has by no means the presumption (like some of his more adventurous contemporaries) to attempt to new-model the Linnaean System, than which for the extension of Botanical knowledge, no other or better foundation can possibly be laid. — The several Universities of Oxford and Cambridge are indebted to the indefatigable researches of men of eminence and learning for the exhibition of their respective Floras. — The present publication forms

a connecting link between the two, comprehending an extensive district in the very center of England, where the GODDESS has scattered her beauties with the gayest and most partial profusion. — Nor is it from a transient or careless view of this delightful tract that the following work is composed.—Many of the specimens have been brought into his own garden, where under the constant superintending care of a fair Associate, (whose assistance it would be injustice not gratefully to acknowledge) their various properties and manner of growth have been daily and hourly presented to his observation. — To the same amiable and interesting partner of his pursuits and labours he is also indebted for the preparation of an HERBARIUM, the admiration of all who have knowledge and judgment to discern it's superior beauty and excellence. — But this is only one of the innumerable obligations, for which he is proud to acknowledge himself indebted to her assiduity and attachment. — To the Rev. T. MARTYN, Professor of Botany in the University of Cambridge, his thanks are due for an able and candid revision of his work, the XXIVth. Class excepted. — To the Right Honorable the COUNTESS of UPPER-OSSORY, for the very generous occasional loan of books from her Ladyship's collection: and lastly, since a late mournful event forbids the public exhibition of his thanks, he deems it

it both his duty and his pride thus to acknowledge the constant and free access he has had to the valuable and well-furnished Library of his late regretted friend THOMAS VAUX, Esq. of Bedford. — Upon the whole the author submits his work to the public with unfeigned respect and diffidence; and he hopes the Linnæan Society, of which he has the honor to be a member, will excuse the mode in which it now appears, in consideration of the motives, which induce him to consult as a leading object the convenience and accommodation of that Class of his Readers, who though comparatively and in the common acceptation of the term unlearned, may not and ought not on that account to be regarded as un-initiated, or as unenlightened.

C. A.

A U T H O R S

A U T H O R S Q U O T E D. ix

- Aiton.* Hortus Kewensis. Lond. 8vo. 1789.
Batsch. Elenchus Fungorum. 4to. 1783, &c.
Blackwell. Mrs.—Herbal of 500 Plants. Lond. 2 vols. fol. 1759.
Bolton. Filices Britannicæ. 4to. Huddersfield. 1790.
 — History of Fungus's. Halifax. fol. 4 vols. 1788. &c.
Bulliard. Histoire des Champignons. Paris. fol.
Curtis. Flora Londinensis. 6 vols. fol. Lond.
Dickson. Three Fasciculi of Cryptogamic Plants. 4to. 1785. 1790. 1793.
Dillenius. Historia Muscorum. Oxoniæ. 1741. 4to.
Dubamel. Traite des Arbres. 4to. Paris. 1755.
Flora Danica. By Oeder, Muller and Vahl. Hafniæ. 1761. &c. fol.
Johnson's Gerard. Lond. fol. 1636.
Haller. Historia Stirpium, &c. 3 vols. Berne. fol. 1768.
L'Heritier. Geraniologia. fol. Paris. 1788.
Hoffman. Enumeratio Lichenum. 3 Fasciculi. 4to. Erlang. 1784, &c.
 — Historia Salicum. vol. 1. Lipsiæ. fol. 1787.
 — Vegetabilia Cryptogamia. 2 Parts. Erlang. 4to. 1787. 1790.
 — Germany's Flora. 12mo. Erlang. 1791.
Hudson. Flora Anglica. 8vo. Lond. 1768.
 Ditto. 2 vols. 8vo. Lond. 1778.
Jacquin. Collectanea. 3 vols. 4to. 1786, &c.

x AUTHORS QUOTED.

- Jacquin.* *Flora Austriaca.* 5 vols. 1773 to 1778.
fol.
— *Miscellanea Austriaca.* 2 vols. 4to.
1773. 1781.
- Leers.* *Flora Herbornensis.* 8vo. 1789.
- Lightfoot.* *Flora Scotica.* 2 vols. 8vo. Lond. 1779.
- Gen. Pl.* *Caroli Linnæi Genera Plantarum.*
- Loesel.* *Flora Prussica.* 4to. Regiomonti. 1703.
- Martyn.* *Flora Rustica.* 4 vols. 8vo. Lond.
1792, 1793, 1794.
- Petiver.* *Herbarium Britannicum.* fol. Lond.
1702.
— *Gazophylaceum.* fol. Lond. 1711.
- Plukenet.* *Phytographia.* 4to. Lond. 1691. 1692.
- Ray's Synopsis.* Third Edition. 8vo. 2 vols.
Lond. 1724.
- Relhan.* *Flora Cantab.* 8vo. Cambridge. 1785, &c.
- Retz.* *Obs: Retzius Botanical Observariions,*
1779, &c.
- Sibthorp.* *Flora Oxoniensis.* 8vo. Oxon. 1794.
- Schæffer.* *Icones Fungorum.* 4 vols. 4to. 1762.
- Schreber.* *de Phasco Observationes.* 4to. Lip-
siæ. 1770.
- Sowerby.* *English Botany.* 6 vols. and 5 odd
Numbers. 8vo. Lond. 1790, &c.
— *Plates of English Fungi.* fol. 1 vol.
& 3 Numbers. Lond. 1796, 1797, 1798.
- Stillingfleet.* *Miscellaneous Tracts.* Lond. 8vo.
1775.
- Vaillant.* *Botanicon Parisiense.* fol. 1727.
- Weigel.* *Observationes Botanicæ.* 4to. 1772.
- Withering.* *Botanical Arrangements.* 1787, &c.
- Woodville.* *Medical Botany.* 4 vols. 4to. Lond.
1793.

CLASSICAL ABBREVIATIONS. xi

M o.	Monandria.
Di.	Diandria. <i>X</i> <i>Dioscuria</i>
Tri.	Triandria. <i>X</i> <i>Crocus</i>
Tetr.	Tetrandria. <i>X</i> <i>Aletris</i>
Pen.	Pentandria. <i>X</i> <i>giola</i>
Hex.	Hexandria. <i>X</i> <i>Narcissus</i>
Oct.	Octandria. <i>X</i> <i>Daphne</i> <i>S. Daphne</i>
Enn.	Enneandria.
Dec.	Decandria.
Dodec.	Dodecandria.
Ico.	Icosandria.
Poly.	Polyandria.
Did.	Gymn. Didynamia. Gymnospermæ. <i>X</i> <i>Lamnum</i>
—	Ang. — Angiospermæ.
Monad.	Pentand. Monadelphia. Pentandria.
—	Decand. — Decandria.
—	Polyand. — Polyandria.
Diad.	Hexand. Diadelphia. Hexandria.
—	Octand. — Octandria.
—	Decand. — Decandria.
Polyad.	Polyand. Polyadelphia. Polyandria.
Polyg.	Æq. Polygamia. Æqualis.
—	Super. — Superflua.
—	Necess. — Necessaria.
—	Frustra. — Frustranea.
Gynand.	Gynandria.
Monoec.	Monoecia.
Dioec.	Dioecia.
Polyg.	Polygamia.
And	Crypt. Cryptogamia.
	Mo.

Mo. Di. Tri. Tetr. Pen. Hex. and Poly.
serve also in the orders to express Monogynia,
Digynia, Trigynia, Tetragynia, Pentagynia,
Hexagynia and Polygynia.

However contrary to the author's wishes, it became necessary, in order to give the volume a more portable appearance, thus to abbreviate, and he is convinced that the scientific Botanist, wanting no such key to the explanation of the Botanical terms, will rather applaud than condemn it.

FLORA BEDFORDIENSIS.

Mr. M. J. L. M. I. M. P. E. C. A. S. E. D. M. R.

zilosisus, C. — 1926. *Quadrivittata*

2011-12 budget - proposed grants - 2011-12 budget - 2011-12 budget

C L A S S. I.

as found at the top, usually within small cells.

I Male, I Female, M_o, M_g male

Mare's-tail. HIPPURIS. Gen. Pl. 15.

Cup none, Blossom none, Stigma simple,
1 Seed.

i. Common Mare's-tail. — *H. vulgaris*.

Leaves in eights, awl-shaped. — Curtis.

Fasc. IV. fig. 1. Flora Danica, tab. 87.

Ditches, May, P. Common.

I Male. II Females. Mo. Di.

Starwort. CALLITRICE. Gen. Pl. 17.

Cup none, Petals 2, Capsule 2-celled, 4 Seeds.

2. Vernal Starwort. — *C. verna*

Upper leaves oval, male and female organs in separate flowers. — Flora Danica.

129. admirable. Sowerby. 722.

Ditches, April, A. Common.

I Male. II Females. Mo. Di.

3. Autumnal Starwort. — C. autumnalis.

All the leaves strap-shaped, nicked at the end, male and female organs in the same flower. — Loesel. Fl. Prus. 38.

Ditches, September, A. Common.

Note.

The three foregoing plants are all to be found in the ditches above Bedford, between the Town and Ford-End Farm.

C L A S S. II.

II Males. I Female. Di. Mo.

Privet. LIGUSTRUM. Gen. Pl. 23.

Blossom 4-cleft, Berry containing 4 Seeds.

4. Common Privet. — L. vulgare.

Curtis. V. 1. Sowerby. 764.

Woods, Hedges, Common, S. June.

Obs:

Leaves ovate, blunt, panicle single or three forked.

Speedwell. VERONICA. Gen. Pl. 32.

Blossom with a border of 4 divisions, lower segment narrowest, Capsule of 2 divisions.

5. Common Speedwell. — V. officinalis.

Spikes lateral on fruit-stalks, leaves opposite, stem trailing. — Curtis. III. 1. admirable. Woodville. 219. Sowerby. 765.

Mountainous

one of Sowerby's desiderata.

II Males. I Female. 1 Di. Md.

Mountainous and Sandy Pastures, P. June,
Rare.

Aspley, — Ampthill, — Clophill.

6. *Smooth Speedwell.* — *V. serpyllifolia.*

Bunch of flowers terminal, resembling a
spike, leaves ovate, smooth, scolloped.

Curtis. I. 3. admirable. **Sowerby. 1075.**

Moist Meadows and Pastures, May, P.
Common,

7. *Brooklime Speedwell.* — *V. Beccabunga.*

Bunches lateral, leaves flat, ovate, stem
creeping. — Curtis. II. 3. **Sowerby. 655.**

Ditches and Rivulets, June, A. Common.

8. *Water Speedwell.* — *V. Anagallis.*

Bunches lateral, leaves spear-shaped, serrated,
stem upright. — Curtis. V. 2. **Sowerby. 701.**

Ditches, July, P. Common.

9. *Narrow-leaved Speedwell.* — *V. scutellata.*

Bunches lateral, little fruit-stalks hanging
down, leaves strap-shaped, seldom toothed. —
Curtis. V. 3. **Sowerby. 702.**

Turfy Bogs and Marshy Places, July, P. Rare.

Fenlake, — Stevington, — Ampthill.

II Males. I Female. I Di. Mo.

Obs:

Stem nearly upright, weak: bunches bearing many flowers, upper segment of the Blossom striped with blue Veins — Runners of the Root mostly barren, so as not to justify it's being called a creeping Plant.

10. *Germander Speedwell.* — *V. Chamædrys.*

Bunches lateral, leaves ovate, sitting, wrinkled, toothed, stem hairy on two sides. —

Sowerby. 623

Martyn. 66. admirable. Curtis. I. 2.

Pastures, May, P. Common.

11. *Mountain Speedwell.* — *V. montana.*

Bunches lateral, with few flowers, cups hairy, leaves ovate, wrinkled, scolloped, on leaf-stalks, stem limber. — Curtis. IV. 2.

Hoffman. Ger. Flo. 1. Sowerby. 766.

Woods and Hedges, May, P. Rare.

Eversbott.

Obs:

Root greatly creeping, leaves deeply notched, Fruitstalks distinct and very long, flowers paler and fewer than in the Chamædrys.

12. *Garden Speedwell.* — *V. agrestis.*

Sowerby. 783. Flowers single, leaves heart-shaped, jagged, shorter than the fruit-stalk. — Curtis. I. 1.

Gardens and Fields, May, A. Common.

Wall

II Males. I Female. Di. Mo.

13. Wall Speedwell. — *V. arvensis*. Sowerby. 734.

Flowers single, leaves heart-shaped, jagged,
longer than the fruit-stalk. — Curtis. II. 2.

Walls and Fields, May, A. Common.

14. Ivy-leaved Speedwell. — *V. hederæfolia*.

Flowers single, leaves heart-shaped, flat,
5-lobed. — Curtis. II. 1. Sowerby. 784.

Fields, April, A. Common.

Butterwort. *PINGUICULA*. Gen. Pl. 40.

Blossom gaping, with a Spur. Cup 2-lipped
5-cleft, capsule 1-celled.

15. Common Butterwort. — *P. vulgaris*.

Honey-cup cylindrical, the length of the
Petal. — Sowerby. 70. admirable.

Bogs, May, P. Rare.

Ampthill Bogs.

Bladderwort. *UTRICULARIA*. Gen. Pl. 41.

Blossom gaping, with a Spur. Cup of 2 leaves,
equal. Capsule 1-cell.

16. Common Bladderwort. — *U. vulgaris*.

Nectary conical, stalk with few flowers. —
Sowerby. 253.

Ditches

II Males. I Female. I Di. Mo.

Ditches and Stagnant Waters, July, P. Rare.

A Ditch in Bromham Parish.

First found by T. VAUX, Esq.

17. *Small Bladderwort.* — *U. minor.*

Nectary keeled. — *Flora Danica.* 128.

Sowerby. 254.

Bogs, June, July, P. Rare.

Amphibill, — *Potton Marshes.*

Water-Horehound. *Lycopus.* Gen. Pl. 44.

Blossom of four divisions, one segment notched at the end, stamens distant; Seeds four, rounded.

18. *Common Water-Horehound.* — *L. europaeus.*

Leaves indented and serrated. — Curtis.

III. 2. *Sowerby.* 1105 — .

Banks and Ditches, August, P. Common.

Sage. *SALVIA.* Gen. Pl. 50.

Blossom unequal. Threads fixed transversely to a little fruit-stalk.

19. *Meadow Sage.* — *S. pratensis.*

Leaves oblong, heart-shaped, scolloped, upper leaves embracing the stem: Flowers in whorls, almost naked: Helmet gummy in the Blossoms. — Sowerby. 153.

Dry

II Males. I Female. II Di. Mo.

Dry Pastures, June, P. very Rare.

A Pasture near Ford-End Farm.

Found by the Rev. T. O. MARSH, F. L. S.

Obs:

Blossom labiate, upper Lip superbly arched, style bent inwards, stigma forked.

20. *Vervain Sage*. — *S. verbenaca*.

Leaves serrated, waved, rather smooth, Blossoms narrower than the Cup. — Sowerby.

154.

Path-ways and Road-sides, June. P. Common.

Enchanter's Nightshade. CIRCAEA. Gen. Pl. 3¹.

Blossom 2 Petals. Cup 2 leaves, superior: Seed-vessel, 1 or 2 divisions.

21. *Common Enchanter's Nightshade*. — *C. lutetiana*.

Stem upright, bunches of flowers many, leaves ovate. — Curtis. III. 3. Sowerby. 1056.

Woods, June, P. Common.

II Males. II Females. Di. Di.

Vernal-Grass. — ANTHOXANTHUM. Gen. Pl. 58.

Cup, Chaff of 2 valves, containing 1 flower; Blossom, a Chaff of 2 valves, pointed. Seed 1.

Sweet

II Males. II Females. Di. Di.

22. Sweet Vernal-Grass. — *A. odoratum.*

Spike ovate-oblong, florets on short fruit-stalks, longer than the Beard. — Curtis.

I. 4. Martyn. 23. Sowerby. 647.

Meadows and Pastures, May, P. Common.

C L A S S. III.

III Males. I Female. Tri. Mo.

* Flowers with Petals.

Valerian. — *VALERIANA.* Gen. Pl. 60.

Cup none. Blossom hunched at the base on one side. superior: Seed 1.

23. Marsh Valerian. — *V. dioica.*

Sowerby. 628. Flowers diæcious, three-stamen'd, leaves winged, very entire. — Curtis. IV. 3.

Moist Meadows and Marshes, May, P. Rare.

Caldwell, — Bromham, near the Mill.

24. Common Valerian. — *V. officinalis.*

Flowers three-stamen'd, all the leaves winged. — Curtis. VI. Sowerby. 698.

Woods, June, P. Common.

25. Corn-Sallad Valerian. — *V. Locusta.*

Flowers three-stamen'd, stem forked, leaves strap-shaped. — Curtis. V. 4. Martyn. 24.

Corn-fields, April, A. Common.

Flag.

23 X Red Valerian. *V. rubra*

Garden walls. July. Rare.

Potton. 1798.

III Males. I Female. I Tri. Mo.

Flag. IRIS. Gen. Pl. 79.

Blossoms with 6 divisions, petals alternate,
bent back, the Stigmas resembling petals.

26. *Yellow Iris.* — *I. pseudacorus.*

Blossoms not fringed, inner petals smaller
than the Stigma, leaves sword-shaped. —

Curtis. III. 4. Flora Danica. 494. admirable. Sowerby. 578.

Banks of Rivers and wet Ditches, June, P.
Common.

27. *Stinking Gladwin Iris.* — *I. fætidissima.*

Blossoms not fringed, inner petals expanding
very much, stem with 1 angle, leaves
sword-shaped. — Blackwell. 158. Sowerby. 596.

Hedges, July, P. Rare.

Brombam, — Stevington, — Pavingham.

Obs:

After flowering the triangular Germ opens and the Seeds,
which are red adhering to the pericarp, are visible in that
situation great part of Autumn and Winter.

** Flowers without Petals.

Bog-Rush. SCHENKUS. Gen. Pl. 92.

Husks chaffy, of 1 valve, rounded, blossom
none, Seed 1. roundish within the Husks.

28. *Black Bog-Rush.* — *S. nigricans.*

Straw round, naked, head of flowers ovate,
1 valve of the 2-leaved fence long and awl-
shaped. Sowerby. 1121.

C Bogs

III Males. I Female. Tri. Mo.

Bogs, July, P. Rare.

Amptbill Moor, — Potton Marshes.

29. *White Bog-Rush. — S. albus.*

Straw nearly three-cornered, leafy, flowers in bundles, leaves bristle-shaped. — Flora Danica. 320. Sowerby. 905.

Bogs, July, P. Rare.

Amptbill Moor, — Potton, — Aspley.

Club-Rush. SCIRPUS. Gen. Pl. 94.

Husks chaffy, tiled all round. Blossom none: Seed 1. beardless.

30. *Marsh Club-Rush. — S. palustris.*

Straw round, naked; spike terminal and nearly ovate. — Flora Danica. 273. Sowerby. 131.

Banks of Rivers and wet Meadows, June, P. Common.

31. *Tall Club-Rush. — S. lacustris.*

Straw round, naked, spikes ovate, many in number, on fruit-stalks terminating. Sowerby 666.

Rivers, Ditches and standing Waters, July, P. Common.

Obs:

A good figure of this Rush would be very desirable.

32. *Dwarf Club-Rush. — S. cespitosus.*

Straw

one of Sowerby's desiderata.

* 29 Compressed Bog-Rush. — S. compressus.

Stem roundish, naked, spike two-ranked shorter than the bracts, which is simple, spikelets many-flowered. Leaves flat. English Botany —

Sowerby. 791.

Bogs and meadows. June. P. rare.

Potton Marshes & Fenlake Bog —

one of Sowerby's desiderata.

30 * Least Club-rush. — S. aciculans.

Straw with 4 angles, spike ovate solitary terminal, sheath at the base of the spike blunt.

Sowerby. 749.

Marsky places Aug. P. rare
Ford End Meadow.

III Males. I Female. Tri. Mo.

Straw scored, naked, spike terminating with 2 valves the length of the Cup. Roots divided by scales. — Flora Danica. 167. Sowerby. 1029. Bogs and Marshes, July, P. Rare.

Amphill and Flitton Moors.

33. Least Club-Rush. — *S. setaceus*. Sowerby. 1693.

Straw bristle-shaped, naked, spikes lateral, sitting, almost double. — Flora Danica. 311.

Wet Places, July, A. Rare.

Amphill Moor, — Near Bromham Bridge.

Cotton-Grass. *ERIOPHORUM*. Gen. Pl. 95.

Husks chaffy, tiled all round; Blossom none, Seed 1. surrounded with very long Wool.

34. Many-headed Cotton-Grass. — *E. polystachion*.

Straws round, leaves flat, spikes on fruit-stalks. — Curtis. IV. 9. Sowerby. 563.

Bogs, April, P. not uncommon. Rare

Amphill Moor, Stockington Bass, near Dunstable, h? Dickson.

*** Flowers Grass-like.

Mat-Grass. *NARDUS*. Gen. Pl. 97.

Cup none: Blossom of 2 valves.

35. Mat Weed. — *N. stricta*.

C 2

Spike

for places read places.

* 34. Narrow-leaved Cotton-grass.

S. angustifolium.

Stems round, leaves channelled, triangular at the tips: spikes several on flower-stalks.

Sowerby. 564.

English Botany.

Bogs - April P. Common.

Obs:

The Botanist is desired to seek for this plant near Dunstable and to erase the habitats given in the former edition of this work, it being now admitted that the former is the rare and this the common species.

III Males. I Female. Tri. Mo.

Spike bristle-shaped, upright, flowering on one side. — Martyn. 27. excellent. Sowerby. 290. well designed.

Heaths, June, P. Common.

III Males. II Females. Tri. Di.

Fox-tail. ALOPECURUS. Gen. Pl. 102.

Cup 2 valves. Blossom 1 valve.

36. Meadow *Fox-tail*. — A. pratensis.

Straw upright, spiked; chaff woolly, blossoms without awns. — Martyn. 6. good.

Curtis. V. 5. *Sowerby*. 759.

Meadows and Pastures, June, P. Common.

37. *Jointed Fox-tail*. — A. geniculatus.

Spiked, straw uninterrupted, blossoms awn-

Sowerby. 1250. less. — Martyn. 97. Curtis. V. 6.

Moist Meadows, sides of Ponds, June, P. Common.

38. *Field Fox-tail*. — A. agrestis.

Spiked, straw upright, husks smooth. —

Sowerby. 848. Martyn. 22. Curtis. II. 7. A. myosuroides.

Cornfields, June, A. too common.

Cat's-Tail. PHLEUM. Gen. Pl. 109.

Cup 2 valves, sitting, strap-shaped: lopped, point bifid; blossom 2 valves inclosed.

Meadow

* 39. Knobby rooted Cat's tail - *P. nodosum*.

Spike cylindrical, stem ascending, leaves oblique, root bulbous.

Road Sides July. P. Common.

* 40 Birds Canary Grass. *P. canariensis*.

Panicle nearly ovate, in the form of a spike chaff boat-shaped.

Sowerby 1310.
May. July. R. Rare.

Mill and Offa Lanes. Bedford.
1799.

III Males. II Females. Tri. Di.

39. Meadow Cat's-Tail. — *P. pratense*.

Spike cylindrical, very long, fringed, straw upright. — Martyn. 5. *Sowerby 1076*.

Meadows and Pastures, July, P. Common.

Canary-Grass. *PHALARIS*. Gen. Pl. 106.

Cup of 2 valves, with 1 flower, the small valves nearly equal, blossom very short, stigmas feather-shaped.

40. Reed Canary-Grass. — *P. arundinacea*.

Panicle oblong, bellied. — Flora Danica.

259.

Rivers and Ditches, July, P. Common.

Millet-Grass. *MILUIM*. Gen. Pl. 110.

Cup of 2 valves, with 1 flower, the small valves nearly equal, blossom very short. Stigmas feather-shaped.

41. Wood Millet-Grass. — *M. effusum*.

Flowers panicled, scattered, awnless. Curtis. IV. 12. *Sowerby 1106* ~~Z~~.

Woods, May, P. Common.

Bent-Grass. *AGROSTIS*. Gen. Pl. 111.

Cup 2 valves, 1 floret, somewhat smaller than the Blossom, Stigmas longitudinally rough with Hair.

Broom

for miluum read milium

III Males. II Females. Tri. Di.

42. *Brown Bent-Grass.* — *A. canina.*
Panicle expanding, Blossom bare, 1 of the valves twice as long as the other, awned: Awn from the back straightish. — Leers. tab. IV. fig. 2.

Heaths and Meadows, July, P. Rare.

Amphibill Warren,—Sandy Heath.

43. *Silky Bent-Grass.* — *A. spica-venti.*

Sowerby 951. Awn straight, stiff, very long, fixed to the outer petal, panicle expanding.

Cornfields, June, A. Rare.

Ford-End Farm,—Luton Downs.

44. *Creeping Bent-Grass.* — *A. stolonifera.*

Branches of the panicle spreading, naked,

Sowerby 1532. culm creeping, cups equal. — Martyn. 120.
Cornfields, July, P. Rare.

St. Leonard's Farm, near Bedford.

45. *Fine Bent-Grass.* — *A. capillaris.*

Panicle hair-like, expanding; cups awl-shaped, equal, rather rough, colored, florets awnless. — Flora Danica. 163.

Sowerby 1671. Pastures and Woods, June, P. Common.

46. *White Bent-Grass.* — *A. alba.*

Panicle loose, cups awnless, equal, straw creeping.

Bogs

III Males. II Females. Tri. Di.

Bogs and wet Meadows, June, P. Common.

Hair-Grass. AIRA. Gen. Pl. 112.

Cup 2 valves, containing 2 florets without
the Rudiment of a third between them.

47. *Water Hair-Grass.* — A. aquatica.

Panicle expanding, florets awnless, smooth,
longer than the cups, leaves flat. — Curtis.

I. 5. Sowerby. 1557.

Ditches and Road-sides, June, P. Common.

48. *Turfy Hair-Grass.* — A. cespitosa.

Leaves flat, panicle expanding, petals woolly
and awned at the base; awn straight, short.
Sowerby. 1453.

Moist Woods and Pastures, June, P. Common.

49. *Heath Hair-Grass.* — A. flexuosa.

Leaves bristle-shaped, straws nearly naked,
panicle diverging, fruit-stalk zigzag. —

Flora Danica. 147. Leers. V. 1. Sowerby. 1519.

Heaths, July, P. Common.

50. *Early Hair-Grass.* — A. præcox.

Leaves bristle-shaped, sheaths angular,
flowers between panicle and spiked, florets
awned at the base. — Curtis. III. 7. Sowerby. 1296.

Heaths, April, May, A. Rare.

Amptbill Heath, — Roteney Warren.

Obs:

III Males. II Females. Tri. Di.

Obs:

This dwarf Plant seldom attains to the height of more than 3 Inches, is easily ascertained by the prostrate manner of its growth, and 2-awned spiculae.

51. *Silver Hair-Grass.* — A caryophyllea.

Leaves bristle-shaped, panicle diverging, florets awned, distant. — Flora Danica.

Sowerby. 812. 382. Leers. V. 7. Curtis. VI. Heaths, July, A. Rare.

Amphibill, — Husborne-Crawley.

Melic-Grass. MELICA. Gen. Pl. 113.

Cup 2 valves, containing 2 florets. Rudiment of a third flower between the florets.

52. *Blue Melic-Grass.* — M. cærulea.

Panicle compact, flowers cylindrical. —

Curtis. V. 11. Sowerby. 750.

Moist Heaths, August, P. Rare.

Amphibill.

Obs:

Root fibrous, straw from half a foot to a foot in length, leaves sheathing at short intervals from the bottom of the straw. Spike loose, anthers purple, spikelets ovate.

53. *Single-flowered Melic-Grass.* — M. uniflora.

Panicle not crowded, cups with 2 florets, 1 hermaphrodite, the other neuter. — Martyn.

64. admirable. Curtis. V. 10. Sowerby. 1058.

Woods

III Males. II Females. Tri. Di.

Woods, May, P. Common.

Obs:

Habit of this Grass generally inclining, leaves broad, mostly open, sometimes rolled in: Cup of 2 leaves pointed, concave; Anthers yellow.

Meadow-Grass. Poa. Gen. Pl. 114.

Cup 2 valves, containing many florets. Spikelets ovate, valves skinny-edged, rather pointed.

54. Water Meadow-Grass. — *P. aquatica*.

Panicle spreading, spikelets strap-shaped, containing 6 flowers. — Curtis. V. 12. Leers. V. 5. Sowerby. 315. —

Banks of Rivers, Ditches, August, P. Common.

Obs:

The most gigantic of all the Meadow-Grasses. Panicle a foot in length, at first close, afterwards spreading, Spikelets ovate, Germ ovate, smooth.

55. Rough-stalked Meadow-Grass. — *P. trivialis*.

Panicle spreading, spikelets containing 3 florets, downy at the base, straw straight, round. — Curtis. II. 6. Leers. VI. 2. Sowerby 1072

Meadows and Pastures, June, P. Common.

D

Narrow-leaved

III Males. II Females. Tri. Di.

56. Narrow-leaved Meadow-Grass. — *P. angustifolia*.

Panicle spreading, spikelets containing 4 florets, downy, straw straight, round. — Leers. VI. 3.

Woods and Hedges, June, P. Rare.

Clapham Lane.

57. Smooth Meadow-Grass. — *P. pratensis*.

Panicle spreading, spikelets containing 5 florets, smooth, straw straight, round. — Curtis. II. 5. Sowerby. 1073.

Walls and Meadows, May, June, P. Common.

58. Annual Meadow-Grass. — *P. annua*.

Panicle spreading horizontally, spikelets blunt, straw obliquely compressed. —

Sowerby. 1141. Curtis. I. 6. admirable. Martyn. 98.

Meadows and Road-sides, April, A. Common.

59. Hard Meadow-Grass. — *P. rigida*.

Sowerby. 1371. Panicle spear-shaped, somewhat branched, flowering on one side, branches alternate, flowering on one side. — Curtis. II. 4.

Walls and dry Places, May, A. Common.

60. Flat-stalked Meadow-Grass. — *P. compressa*.

Panicle flowering on one side, compact, straw oblique, compressed. — Flora Danica.

742. Sowerby. 365. Leers. V. 4.

Walls

Veronica montana.
Centunculus minimus.
Sagina erecta.
Pulmonaria officinalis.
Lysimachia nemorum.
Anagallis corulea.
Campanula latifolia.
Caucalis danoides.
Caucalis latifolia.
Lisianthus sagetum.
Drosera anglica.

Ornithogalum pyrenaicum.
Convallaria multiflora.

Rumex aureus

Polygonum Bistorta.

Dianthus deltoides — *Pandurina*.

Zygophyllum hyssopifolium

Rosa villosa.

Geum rivale

Ranunculus flavis.

Helleborus foetidus

Mentha sylvestris.

Lathra squamaria.

Limosella aquatica.

Geranium phaeum

Zizyphus spinosa.

Vicia sylvatica

Trifolium subterraneum.

Trifolium ochroleucon

Trifolium filiforme.

Lactuca scariola.

Aicracium umbellatum.

Argemone glabra.

Hyoscyamus niger.

Inula pulicaria.
Vista pratensis.
Ophrys spiralis
Ophrys muscifera.
Malaxis paludosa.
Hydrocharis m. ranae.
Ruscus aculeatus.

19

III Males. II Females. Tri. Di.
Walls, June, P. Common.

61. *Wood Meadow-Grass.* — *P. nemoralis.*
Panicle tapering, spikelets nearly three-flowered, pointed, rough, straw bent inwards. — *Flora Danica.* 749. *Leers.* V. 3. *Sowerby.* 1265.
Woods, June, P. Rare. — *Clapham-Park Wood.*

62. *Loose Meadow-Grass.* — *P. distans.*
Panicle with subdivided branches, spikelets containing 5 florets, florets distant, blunt. — *P. retroflexa.* *Curtis.* VI. *Sowerby.* 986. —
Watery Places, June, P. Rare. — *Clapham Lane.*

63. *Crested Meadow-Grass.* — *P. cristata.*
Panicle spiked, cups somewhat hairy, about 4 florets in each, longer than the fruit-stalk, petals awned. *Leers.* V. 6. *Sowerby.* 640. 640.
Dry Pastures, June, P. Common.
Quaking-Grass. *BRIZA.* *Gen. Pl.* 115.
Cup 2 valves, containing many florets, little spikes in 2 rows, valves heart-shaped, blunt, inner valve smallest.

64. *Middle Quaking-Grass.* — *B. media.*
Spikelets ovate, cup shorter than the florets. — *Flora Danica.* 253. *Sowerby.* 340. *Martyn.*
39. — *D 2* — *Dry*

III Males. II Females. Tri. Di.

Dry Meadows and Pastures, July, P. Common.

Cock's-foot-Grass. DACTYLIS. Gen. Pl. 117.

Cup 2 valves flat, 1 valve larger, keeled.

65. Rough Cock's-foot-Grass. — *D. glomerata*.

Panicle crowded, flowering on one side. —

Sowerby. 335. Martyn. 14.

Meadows, Pastures, July, P. Common.

Fescue-Grass. FESTUCA. Gen. Pl. 119.

Cup 2 valves, spikelet oblong, almost cylindrical, chaff pointed.

66. Sheep's Fescue-Grass. — *F. ovina*.

Panicle flowering on one side, compact, awned, straw four-square, almost naked, leaves bristle-shaped. — Leers. VIII. 3. Martyn. 102. admirable. Sowerby. 595.

Dry Pastures, June, July, P. Common.

67. Hard Fescue-Grass. — *F. duriuscula*.

Panicle flowering on one side, oblong: spikelets oblong, smooth, leaves bristle-shaped. — Sowerby. 470. Flora Danica. 848.

Dry Pastures, June, July, P. Common.

68. Purple Fescue-Grass. — *F. rubra*.

Panicle

* 66. Fine Sheep's Fescue. *F. tenuifolia*.
Leaves bristle-shaped, rather long, erect:
straw naked, spikelets roughish.

Leers. VIII 4.

Dry spots, June P. Rare.

Milton Grays.

Obs:
The want of awns, and the roughness
of the spikelet, seem to form a
sufficient specific character, and indeed
to justify the late professor Sibthorpe
in keeping it separate from *F. ovina*.

one of Sowerby's desiderata. tent.

* 72 Darnel Fescue. — F. loliacea.

Spiked, spikelets in alternate order,
sitting, compressed, awnless. Hudson.

Curtis. VI. f.

Moist Meadowos — June. Rare.

Friar's pasture — Oakley —
Meadows near Oakley House.

Obs.

Nature between that of the common
perenne, and F. elatior: The
absence of the inner calyx-valve
and the hollow in that case
visible in the rachis for the re-
ception of the spikelet mark.

The species, and at once put an
end to that confusion which is at
first caused in the systematic
Botanist, when he finds in a
hybrid production the two distinct
characters of its unnatural parents.

III Males. II Females. Tri. Di.
Panicle flowering on one side, rough:

spikelets containing 6 florets, awned; last
floret awnless, culm half cylindrical. —

Stillingfleet. tab. IX. Leers. VIII. I.

Hills, July, P. Rare.

Luton Downs, — Barton Hill.

69. Wall Fescue-Grass. — F. myurus.

Panicle spiked; cups very small, awnless;
florets rough, awns long. — Leers. III. 5.

Walls, May, A. Rare. Sowerby. 1412.

Oakley Walls.

70. Meadow Fescue-Grass. — F. pratensis.

Panicle flowering on one side, oblique: Sowerby. 1592.
spikelets nearly strap-shaped, awnless,
leaves flat. — Martyn. 84. Curtis. VI.

Meadows, June, July, P. Common.

71. Tall Fescue-Grass. — F. elatior.

Panicle spreading, spikelets ovate, pointed, Sowerby. 1593.
leaves broad. — Leers. VIII. 6. Curtis. VI.

Meadows and Pastures, June, July, P.
Common.

72. Flote Fescue-Grass. — F. fluitans.

Panicle branched, straight, spikelets nearly
sitting, cylindrical, awnless. — Curtis. I.

7. admirable. Martyn. 113. Sowerby. 1520.

Wet

III Males. II Females. Tri. Di.

Wet Ditches and Ponds, June, P. Common.

Obs:

The most succulent of the Grass tribe, affording an excellent Fodder for Cattle.

73. *Decumbent Fescue-Grass.* — *F. decumbens.*

Panicle straight, spikelets nearly ovate, awnless, cup larger than the florets, straw

Sowerby 792. leaning. — Flora Danica. 162.

Pastures and dry Heaths, July, P. Rare.

Mouse's Pasture, near Bromham.

74. *Wood Fescue-Grass.* — *F. sylvatica.*

Spiked, spikelets alternate in 2 rows, sitting, awned. — Flora Danica. 164.

Woods and Hedges, July, P. Common.

75. *Spiked Fescue-Grass.* — *F. pinnata.*

Spiked, nearly 2-rowed, spikelets alternate, round, nearly awned. — Leers. X. 3.

Hedges, July, P. Rare.

Clapham between Woods.

Obs:

Joints smooth, silvery, spikelets not (as in the preceding) pressed close to the spike-stalk, but very distant.

Brome Grass. BROMUS. Gen. Pl. 120.

Cup of 2 valves, spikelet oblong, cylindrical, in 2 rows. Awns arising from beneath the End.


76. *Field Brome-Grass.* — *B. secalinus.*

Broom

III Males. II Females. Tri. Di.

Panicle spreading, spikelets awnless, straight, awns wanting. Leers.

one of Sowerby's desiderata. sent.



* 76

Dowry Rye Brome Grass. *B. multiflorus.*

Panicle expanding, fruit stalks nearly simple, spikelets ovate: lance-shaped compressed containing fifteen florets, florescules somewhat tilted, nearly cylindrical — Leers. XI. 2.

Fields & Roadsides. July. A. Rare.

Maulden, Chicksands.
1702

III Males. II Females. Tri. Di.

Panicle expanding, spikelets ovate, awns
straight, seeds distinct. Leers. XI. 2. Sowerby 1171.
Cornfields, May, A. Rare.

Bury Field, near Bedford.

77. Soft Brome-Grass. — B. mollis.

Panicle rather straight, spikelets ovate,
downy. Awns straight, leaves woolly,
very soft. — Curtis. I. 8. Martyn. 99. Sowerby 1078.
Walls, Pastures, May, A. Common.

78. Corn Brome-Grass. — B. arvensis.

Panicle spreading, its stalks branched,
spikelets lanceolate, about 8-flowered.
Florets elliptical, imbricated, depreped,
smoothish. — P. Botany. Sowerby. 920.
Clapham Hill — Church Yard Walls.
Harrowton Church —
Rising Ground behind Clapham
June. 1799. Church

79. Barren Brome-Grass. — B. sterilis.

Panicle spreading, spikelets oblong, in two
rows, chaff with awl-shaped awns. — Curtis.
I. 9. Martyn. 125. Sowerby 1030 —
Walls and Hedges, June, A. Common.

80. Rough Brome-Grass. — B. hirsutus.

Panicle nodding, rough, spikelets strap-
shaped, containing 10 florets, awns straight,
sheaths of the leaves hairy, — Curtis. II. 8.
Martyn. 126. Sowerby 1172.

Woods, Hedges, July, P. Common.

Tall

III Males. II Females. Tri. Di.

81. *Tall Brome-Grass.* — *B. giganteus.*

Panicle nodding, spikelets with 4 florets,
shorter than the awns. — Curtis. V. 7.
Leers. X. 1.

Woods and Hedges, July, P. Rare.

Bromham, — Hazel-wood Lane.

Oat-Grass. AVENA. Gen. Pl. 122.

Cup of 2 valves, many-flowered, with a
twisted awn on the back.

82. *Bearded Oat-Grass.* — *A. fatua.*

Panicled, cup with 3 florets, all awned and
hairy at the base. — Martyn. 81. admirable.

Leers. IX. 4.

Cornfields, July, A. Common.

83. *Rough Oat-Grass.* — *A. pubescens.*

Almost spiked, cups with about 3 florets,
hairy at the base, leaves flat, downy. —
Sowerby. 1640. Ray's Synopsis. tab. XXI. fig. 2.

Pastures, June, P. Common.

84. *Meadow Oat-Grass.* — *A. pratensis.*

Panicle undivided, chaff with 5 florets,
leaves rolled in, naked. — Ray's Synopsis.
XXI. 1.

Dry Pastures, July, P. Common.

III Males. II Females. Tri. Di.

85. *Yellow Oat-Grass.* — *A. flavescens.*

Panicle loose, cups short, 3 florets in each,
all awned. — Curtis III. 5. Martyn 112. Sowerby. 952.
Meadows, Pastures, June, P. Common.

86. *Tall Oat-Grass.* — *A. elatior.*

Panicled, cups with 2 florets, the one with
both male and female organs nearly awn-
less; the other with only male, awned. —
Curtis. III. 6. Martyn. 7. Sowerby. 913.

Obs:

I have observed with regret this Plant arranged with the
Holci, it were better, if Linnæus must be departed from,
to return the Holci to their natural station amongst the
triandrous Grasses.

Reed-Grass. ARUNDO. Gen. Pl. 1201.

Cup, 2 valves. Florets crowded, surround-
ed with wool.

87. *Common Reed-Grass.* — *A Phragmites.*

Panicle loose, cups with 5 florets. — Sow-
erby. 401.

Wet Ditches and Banks of Rivers, July, P.
Common.

88. *Wood Reed-Grass.* — *A Calamagrostis.*

Cups with 1 floret, smooth. Blossoms downy,
straw branched. — Flora Danica. 280.
Sowerby. 403.

III Males. II Females. Tri. Di.

Woods, June, P. Rare.

Sheer-batch Wood.

89. *Small Reed-Grass.* — *A. epigejos.*

Cups with 1 floret, panicle straight, leaves smooth underneath. — Sowerby. 402.

Woods, July, P. Rare.

Maulden Wood.

Darnel-Grass. *LOLIUM.* Gen. Pl. 126.

Cup 1 leaf fixed, containing 1 floret.

90. *Perennial Darnel-Grass.* — *L. perenne.*

Spike awnless, spikelets flat, with many flowers. — Sowerby. 315. Martyn. 4.

Pastures, June, P. Common.

91. *White Darnel-Grass.* *L. temulentum.*

Spike awned, spikelets flat, with many flow-

Sowerby. 1124. ers. — Leers. XII. 2. Martyn. 33.

Cornfields, July, A. Rare.

Near Fenlake.

Barley. *HORDEUM.* Gen. Pl. 129.

Cup lateral, 2 valves, 1 floret: Cup trifid.

92. *Wall Barley.* — *H. murinum.*

Florets on the side, male ones awned, intermediate fences fringed. — Curtis, V. 9.

Martyn. 43.

Walls

Oat-fields near Crow Hills to the East

III Males. II Females. Tri. Di.

Walls and Road-sides, June, P. Common.

93. *Wood-Barley*. — *H. sylvaticum*.

All the florets with male and female organs,
awned: fences growing together at the base,
shorter than the awn. — Martyn. 45. *Sowerby* 1317

Woods, June, P. Rare.

Thurlegh, — *Putnoe Woods*.

94. *Meadow-Barley*. — *H. pratense*.

Florets on the side, male ones awnless, fences
bristle-shaped, rough. — Martyn. 108.

Sowerby. 409.

Meadows, Pastures, June, P. Common.

Wheat-Grass. *TRITICUM* Gen. Pl. 130.

Cup of 2 valves, solitary, containing about
3 florets, flowers bluntish, tapering.

95. *Creeping Wheat-Grass*. — *T. repens*.

Cups with 4 florets, awl-shaped, pointed,
leaves flat. — Martyn. 124. *Sowerby* 909.

Fields, July, P. too common.

96. *Bearded Wheat-Grass*. — *T. caninum*.

Cups awl-shaped with nearly 4 florets, florets
awned. *Sowerby*. 1372

Woods, Hedges, July, P. Common.

Dogs-tail-Grass. *CYNOSURUS*. Gen. Pl. 118.

III Males. II Females. Tri. Di.

Cup of 2 valves, many florets, proper receptacle leafy, fixed to the side.

97 Crested Dog's-tail-Grass. — *C. cristatus*.

Floral leaves with winged clefts. — Martyn.

106. Sowerby. 316.

Meadows and Pastures, June, P. Common.

III Males. III Females. Tri. Tri.

Blinks. MONTIA. Gen. Pl. 133.

Cup 2 leaves : Blossom 1 petal, irregular : Capsule 1-celled, 3-valved.

98. Water Chickweed or Blinks. — *M. fontana*.

Curtis. III. 8.

Wet Places, May, A. Rare.

Amphill Moor, — Potton Marshes.

Obs:

Leaves oblong, growing together, sharpish at the points. Root creeping, very fibrous. Flowers minute, seldom entirely open, whence it's English name (Blinks) and the buds nod before blowing.

CLAS. IV.

IV Males. I Female. Tetr. Mo.

Teasel. DIPSACUS. Gen. Pl. 148.

Common cup of many leaves, proper cup superior : Receptacle chaffy.

Wild

IV Males. II Females. Tetr. Mo.

Wild Teasel. — *D. officinalis*.

Leaves entire, chaff straight — Curtis. III. 9. Sowerby. 1032.

Hedges and Road-sides July, P. Common

and Small Woods or Fences. — Curtis. D. plur.

Leaves on leaf-stalks with appendages.

Curtis. I. 16. Chodat. 1.

Wet Meadows, June, P. Common.

Keepers' Field, — Curtis. 1.

Height of the Plant from 1 to 6 feet or more, about

less than in the previous but larger than in the last species.

This plant is the best of the genus.

Scavions. Scavions. — Curtis. 1.

Common cup of many leaves, proper cup double, superior receptacle chaffy or naked.

101. Devil's-bit Scabious. — *S. succisa*.

Flowers with equal segments, each one-sided, being the upper part white, the lower between lance-shaped and white. — Curtis. VII. 10. Sowerby. 1033.

Meat Woods and Fields, July, P. Common.

102. Field Scabious. — *S. pratense*.

Flowers with 5 pointed lobules, leaves

winged, suggesting a white wavy line.

Curtis. V. 10. Sowerby. 1034.

Common

IV Males. I Female. Tetr. Mo.

99. *Wild Teasel*. — *D. sylvestris*.
Leaves entire, chaff straight. — Curtis. III. 9. Sowerby. 1032
Hedges and Road-sides, July, P. Common.

100. *Small Teasel or Shepherd's Rod*. — *D. pilosus*.
Leaves on leaf-stalks with appendages. —
Curtis. I. 10. Sowerby. 877.
Wet Hedges, July, B. Rare.

Kempston, — Medbury, — Goldington.

Obs:

Height of the Plant from 3 to 6 feet or more. Heads less than in the preceding, not thorny, whitish, with dull-purplish Anthers, stem prickly all the way up: terminal Head solitary in the Fork of the main Branch.

Scabious. SCABIOSA. Gen. Pl. 149.

Common cup of many leaves, proper cup double, superior: receptacle chaffy or naked.

101. *Devil's-bit Scabious*. — *S. succisa*.

Blossoms with 4 equal segments, stem undivided, branches approaching, leaves between lance-shaped and ovate. — Curtis. III. 10. Sowerby. 878.

Moist Woods and Fields, July, P. Common.

102. *Field Scabious*. — *S. arvensis*.

Blossoms with 4 segments, radiate, leaves winged, jagged, stem rough with hair. — Curtis. IV. 13. Sowerby. 659.

Cornfields

IV Males. I Female. Tetr. Mo.

Cornfields and Pastures, June, P. Common.

103. *Small Scabious*. — *S. Columbaria*.

Blossoms with 5 segments, radiate, leaves at the root ovate, notched, stem-leaves winged, Sowerby 1311 bristle-shaped, — *Flora Danica*. 314.

Mountainous Pastures, June, P. Common.

Chaff-weed. *CENTUNCULUS*.

Cup with 4 segments, blossom uniform, open: Threads short, seed-vessel cut round as in the Genus *Anagallis* with this difference, it has a pointed beak.

104. *Pimpernel Chaff-weed*. — *C. minimus*.

Vaillant. tab. IV. 2. Curtis. III. 11. admirable. Sowerby 531.

Boggy Ground, July, A. Rare.

Ampthill Moor.

Obs:

Flowers sitting in the wings of the leaves, flesh-color'd, very seldom expanded, leaves ovate, sharp. The whole Plant varies in stature from half an Inch to 3 Inches.

Plantain. *PLANTAGO*, Gen. Pl. 186.

Cup with 4 segments, blossom uniform, with the border reflected. Threads very long, seed-vessel 2-celled, cut round.

105. *Great-Plantain*. — *P. major*.

Leaves

IV Males. I Female. Tetr. Mo.

Leaves ovate, smooth, stem-leaves

with blossoms — *Flora Danica*.

Pastures and Roadsides, June, P. Common.

106. *Middle-Plantain*. — *P. media*.

Leaves broad, smooth,

seed-vessel 2-celled.

Medical and Garden.

Pastures, June, P. Common.

107. *Round-Plantain*. — *P. rotundifolia*.

Leaves round, smooth, stem-leaves

with blossoms — *Flora Danica*.

Leaves round, smooth, stem-leaves

with blossoms — *Flora Danica*.

Leaves round, smooth, stem-leaves

with blossoms — *Flora Danica*.

Leaves round, smooth, stem-leaves

with blossoms — *Flora Danica*.

Sandy meads, June, P. Common.

108. *Smooth-Plantain*. — *P. glabra*.

Smooth, narrow, lanceolate,

leaves smooth, stem-leaves

IV Males, I Female, I Tetr. Mo.

Leaves ovate, smooth, stalk round, spike
tiled with blossoms. — Curtis. II. 11. **Sowerby. 1558.**

Pastures and Road-sides, June, P. Common.

106. *Middle-Plantain.* — *P. media.*

Leaves between ovate and spear-shaped, downy,
spike and stalk cylindrical. — Flora
Danica. 581. Curtis. IV. 14. **Sowerby. 1559.**

Pastures, June, P. Common.

107. *Ribwort-Plantain.* — *P. lanceolata.*

Leaves spear-shaped, spike naked, nearly
ovate, stem angular. — Martyn. 67.
admirable, Curtis. II. 10. **Sowerby - 507.**

Woods and Pastures, June, P. Common.

108. *Bucks-born-Plantain.* — *P. Coronopus.*

Leaves strap-shaped, toothed, stem cylindrical.
— Flora. Danica. 272. **Sowerby. 892.**

Sandy Fields, June, A. Rare.

Amphill, — Aspley.

Bloodwort. **SANGUISORBA.** Gen. Pl. 190.

Cup 2-leaved. Germ between the cup and
blossom.

109. *Burnet-Bloodwort.* — *S. officinalis.*

Spikes ovate. — Flora Danica. 97. Martyn.
142. Sowerby. 1312.

Moist Meadows, July, P. Rare.

Bromham, — Fenlake, — Cow Meadows.

Field

III Males. I Female. Tetr. Mo. VI

Field Madder. SHERARDIA. Gen. Pl. 156.

Blossom 1 petal, funnel-shaped, seeds 2, with
3 teeth.

101 Little Field-Madder. — *S. arvensis*.

All the leaves whorled, flowers terminal. —

Curtis. V. 13. Flora Danica, 439. admirable. Sowerby. 891.

Cornfields, June, A. Rare.

Clapham Hill, — Aspley, — Ampthill Warren.

Woodruff. ASPERULA. Gen. Pl. 157.

Blossom 1 petal, funnel-shaped. Seeds 2,
round.

III. Sweet Woodruff. — *A. odorata*.

Leaves eight at a joint, lance-shaped, flowers
in bundles on fruit-stalks. — Curtis. IV.

15. Blackwell. 60. Sowerby. 755.

Woods, May, P. Rare.

Aspley, — King's Wood, near Ampthill.

Obs:

Root truly creeping. Leaves in whorls 6, 7, or even 8 at
a joint, between ovate and spear-shaped, flowers white, crowded,
smelling like new-mown hay.

112 Squinancy Woodruff. — *A. cynanchica*.

Leaves in fours, strap-shaped, upper ones
opposite, stem upright, flowers with 4 seg-
ments.

Sowerby.

Line 5
for 101 read for III males
110. read IV males.

IV Males. I Female. Tetr. Mo.

Sowerby. 33. admirable.

Chalky Hills, July, P. Rare.

Dunstable, — Barton.

Obs:

The stems trail among grass, leaves differing in length. Flowers blush-colored, mouth striped with red veins, upper leaves 3 at a joint, or even 2.

Bedstraw. *GALIUM.* Gen. Pl. 162.

Blossom 1 petal, flat. Seeds 2, roundish.

113. *White Bedstraw.* — *G. palustre.*

Leaves in fours, roundish-ovate, unequal, stem spreading. — Flora Danica. 423. admirable.

Wet Ditches, July, P. Common.

114. *Marsh Bedstraw.* — *G. uliginosum.*

Leaves in sixes, spear-shaped, with prickly serratures bowed back, pointed, stiff, blossoms larger than the seed. — Petiver. tab.

XXX. fig. 6.

Bogs, July, P. Rare.

Amptbill, — Potton, — Stevington.

115. *Smooth Bedstraw.* — *G. spurium.* Sowerby. 1641.

Leaves in sixes, spear-shaped, keeled, rough with prickles pointing backwards, joints simple, fruit smooth. — Vaillant. IV. 3.

Cornfields, June, A. Common.

IV Males. I Female. Tetr. Mo.

116. Trailing Bedstraw. — *G. procumbens.*

Leaves in sixes, round, ovate, pointed,
smooth, stem prostrate, fruit-stalks trebly
forked. — Petiver. XXX. 6. Sowerby. 815.

Heaths, June, P. Rare.

Ampibill, — Sandy Warren.

117. Yellow Bedstraw. — *G. verum.*

Leaves in eights, strap-shaped, furrowed,
flowering branches short. — Martyn. 54.
Curtis. VI. 13. Sowerby. 660.

Pastures, July, P. Common.

118. Great Bedstraw. — *G. Mollugo.*

Leaves in eights, ovate-strap-shaped, almost
serrated, very open, pointed. Stem weak,
branches spreading. — Flora Danica. 455.
admirable. Sowerby. 1673.

Hedges, July, P. Common.

119. Cleavers Bedstraw. — *G. Aparine.*

Leaves in eights, spear-shaped, keels rough,
prickles pointed backward, joints woolly,
fruit rough. — Curtis. II. 9. Martyn. 104.
Flora Danica. 495. admirable. Sowerby. 816.

Hedges and Cornfields, July, A. Common.

120. Upright Bedstraw. — *G. erectum.*

Leaves

IV Males. I Female. Tetr. Mo.

Leaves in eights, smooth, spear-shaped, pointed, stem nearly erect, terminal panicle with forked fruit-stalks. — Jacquin Flora Austr. tab. 80.

Woods and Pastures, June, P. Rare.

Ford End,—Woods near Dunstable.

121. Least Bedstraw. — *G. pusillum*.

Leaves about eight in a whorl, rough, strap-shaped, pointed, fruit-stalks forked, fruit smooth. — Sowerby, 74. admirable.

Fields, July, P. Rare.

Shefford.

Obs :

The whorls of the leaves at the Root are very compact, habit of the plant rather rigid. Flowers more elegant than in the rest of the Family.

Cornel. CORNUS. Gen. Pl. 194.

Fence often four-leaved. Petals 4, superior, nut of 2 cells, covered with a pulp.

122. Female Cornel. — *C. sanguinea*.

Branches upright, leaves ovate, of an uniform colour, Cymes flat. — Flora Danica. 481. Sowerby. 249. admirable.

Woods and Hedges, June, P. Common.

Ladies-Mantle. ALCHEMILLA, Gen. Pl. 222.

Cup eight divisions. Blossom none; 1 seed.

IV Males. I Female. Tetr. Mo.

123. Common Ladies-Mantle. — *A. vulgaris.*

Leaves lobed. — Flora Danica. 693. Blackwell. 72. (See our fig. No. I.) Sowerby. 597.

Dry Pastures, June, P. Rare.

Whipsnade, — Eversholt.

Obs:

Of all our Natives this is the most elegant Plant. The leaves most simply, yet beautifully plaited, flowers palish-green, of a grassy colour.

124. Pursley-Piert Ladies-Mantle. — *A. Aphanes.*

Hairy, leaves having three lobes, which are twice or thrice subdivided. — Petiver. IX.

12.

Fields, May, A. Common.

IV Males. II Females. Tetr. Di.

Dodder. *Cuscuta.* Gen. Pl. 227.

Cup 4 segments. Blossom 1 petal. Capsule 2-celled.

125. Great Dodder. — *C. europaea.*

Flowers nearly sitting, blossom without scales, stigma simple. — Sowerby. 378. admirable.

Nettles, Beans, &c. July, A Common.

126. Least Dodder. — *C. epithymum.*

Flowers sitting, a fringed scale close at the base of each stamen, Stigmas simple. — Sowerby. 55. admirable.

On



IV Males. II Females. Tetr. Di.

On Mother of Thyme, Heath, &c. June,
A. Rare.

Ampthill Warren, — Barton Hill.

IV Males. IV Females. Tetr. Tetr.

Holly. ILEX. Gen. Pl. 232.

Cup with 4 teeth, blossom wheel-shaped,
style none. Berry containing 4 seeds.

127. *Common Holly.* — I. aquifolium. Sowerby. 496.

Leaves ovate, sharp, thorny. — Flora Danica.

508.

Hills and Woods, S. Rare.

Aspley.

Pondweed. POTAMOGETON. Gen. Pl. 234.

Cup none, petals 4, style none, seeds 4.

128. *Floating Pondweed.* — P. natans,

Leaves oblong-ovate, on leaf-stalks, floating. — Flora Danica. 165.

Ditches and Rivers, July, P. Common.

129. *Perfoliate Pondweed.* — P. perfoliatum.

Leaves heart-shaped, embracing the stem. —
Sowerby. 168.

Rivers, July, P. Common.

130. *Shining Pondweed.* — P. lucens.

Leaves

IV Males. IV Females. Tetr. Tetr.

Leaves spear-shaped, flat, ending in leaf-stalks. — Flora Danica. 195. Sowerby. 376.
Rivers, July, P. Common.

131. Forked Pondweed. — *P. densum*.

Leaves ovate, pointed, opposite, crowded, stems forked, spike containing 4 flowers. — Sowerby. 397.

Ditches, June, P. Rare.

Ford-End.

Obs:

One flower arises from the fork of the stem, sitting on a short fruit-stalk, and resembles the flower of the *Adoxa*.

132. Flat-stalked Pondweed. — *P. compressum*.

Leaves strap-shaped, blunt, stem flattened. — Flora Danica. 203. Sowerby. 418.

Ditches, June P. Rare.

Ditch near Fenlake Osier-holts.

133. Curled Pondweed. — *P. crispum*.

Leaves spear-shaped, alternate or opposite, waved, serrated. — Curtis. V. 15. admirable.

Rivers and Ponds, June. P. Common.

134. Fennel-leaved Pondweed. — *P. pectinatum*.

Leaves bristle-shaped, parallel, approaching, in 2 rows. — Sowerby. 323.

Rivers

IV Males. IV Females. Tetr. Tetr.

Rivers and Ditches, June, P. Rare.

Ditch near Fenlake Osier-holts.

135. Least Pondweed. — *P. pusillum*.

Leaves strap-shaped, opposite and alternate, separate, even at the base, stem round. — Sowerby. 215.

Ditches and Ponds, near the mouth of the main River, June, P. Common.

Brackish. Salix. Gen. Pl. art.

Cup of 4 leaves. Petals & Seed-vessel 4-lobed, valves membranous.

136. Perennial Pondweed. — *P. perenne*.

Branches reading. — Curtis. Ed. 3. Sowerby.

More situations on Rivers, June, P. Common.

137. Spear-leaved Pondweed. — *P. hastatum*.

Stem rather upright, downy, flowers white, without petals.

Walls, June, P. Rare.

Mucky-bottoms near Ditches.

IV Males. IV Females. Tetr. Tetr.

Rivers and Ditches, June, P. Rare.

Ditch near Fenlake Osier-holts.

*35. Least Pondweed. — *P. pusillum*.

Leaves strap-shaped, opposite and alternate, separate, open at the base, stem round. — Sowerby. 215.

Ditches and Ponds, that communicate with the main River, June, P. Common.

Pearlwort. *SAGINA*. Gen. Pl. 236.

Cup of 4 leaves. Petals 4. Seed-vessel 1-celled, 4-valved : many seeded.

*36. Procumbent Pearlwort. — *S. procumbens*.

Branches trailing. — Curtis. III. 12. Sowerby. 880.

Moist situations on Heaths, July, P. Common.

*37. Apetalous Pearlwort. — *S. apetala*.

Stem rather upright, downy, flowers alternate, without petals. — Curtis. V. 14. Sowerby. 881.

Walls, June, A. Rare.

Market-Street, near Dunstable.

Obs:

Stem erect from 1 and a half to 3 Inches high. Petals very minute, four, nicked, almost heart-shaped; Segments of the cup ovate, concave.

Upright

IV Males. IV Females. Tetr. Tetr.

138. *Upright Pearlwort.* — *S. erecta.*

Sowerby. 609. Stem erect, with seldom more than one flower. — *Curtis.* II. 12. admirable.

Heaths, April, A. Rare.

Clophill and Ampthill Warrens.

Obs:

Stem erect 3 or 4 inches high, root fibrous, leaves springing together, sea-green, strap-shaped, sharp : Segments of the cup sharpish : Petals 4, larger than in the other Species, yet shorter than the cup.

C L A S S. V. *brevifolia*.

V Males. I Female. Pen. Mo.

Scorpion-Grass. *MYOSOTIS.* Gen. Pl. 240.

Blossom salver-shaped, 5 divisions, nicked ; mouth closed with vaulted Roofs.

139. *Marsh Scorpion-Grass.* — *M. palustris.*

Cups roundish-oval, blunt, leaves spear-shaped. — *Curtis.* III. 13.

Ditches, June, P. Common.

140. *Field Scorpion-Grass.* — *M. arvensis.*

Seeds smooth, points of the leaves callous.—

Flora Danica. 583. *Sowerby*. 480.

Cornfields, May, A. Common.

Gromill.

V Males. I Female.

Gromill. Linnaeus's Species Plantarum.

Blossom roundish-oval, 5 divisions.

The mouth of Cup in the division.

44. *Common Gromill.* Linnaeus's Species Plantarum.

Stems smooth, base smoothish.

Cup in division.

VIII. 243.

Hedges and Roadsides. — Flora Danica.

Common Name. — *Gromill.*

Vol Males. I. Female. Pen. Mo.

Gromill. *LITHOSPERMUM.* Gen. Pl. 241.

Blossom funnel-shaped: open and naked at the mouth. Cup in five divisions.

141. *Common Gromill.* — *L. officinale.*

Seeds smooth, blossoms hardly exceeding the cup in length. — Sowerby. 134. Woodville. 213.

Hedges and Road-sides, May, P. Rare.

Newenham Wall.

142. *Corn Gromill.* — *L. arvense.*

Seeds rough, blossoms hardly exceeding the cup in length. — Flora Danica. 456. Sowerby. 123.

Cornfields, May, A. Common.

Hound's-tongue. *CYNOGLOSSUM.* Gen. Pl. 243.

Blossom funnel-shaped. Mouth closed by vaulted roofs.

143. *Common Hound's-tongue.* — *C. officinale.*

Threads shorter than the blossom, leaves broad, spear-shaped, downy, sitting. — Curtis. IV. 16. Woodville. 216. Sowerby. 921.

Road-sides, June, P. Common.

Lungwort. *PULMONARIA.* Gen. Pl.

Blossom funnel-shaped, mouth with a passage. Cup a prism of 5 angles.

V Males. I Female. P. Pen. Mo.

144. *Broad-leaved Lungwort.* — *P. officinalis.*

Root leaves between ovate and heart-shaped,
rough. — Sowerby. 118. Woodville. 212.

Woods and Hedges, April, P. Rare.

Between Thurleugh and Milton-Ernys.

Obs :

Leaves spotted with white.

Comfrey. SYMPHYTUM. Gen. Pl. 245.

Border of the blossom between tubular and
bellying, the mouth closed with awl-shaped
rays.

145. *Common Comfrey.* — *S. officinale.*

Leaves between ovate and lance-shaped, run-
ning down the stem, cup closing the whole
length of the blossom's tube. — Curtis. IV.
18. Woodville. 215. Sowerby. 817.

Banks of Rivers, May, P. Rare.

Banks of the Ouse.

146. *Red Comfrey.* — *S. patens.*

Leaves between ovate and lance-shaped, run-
ning down the stem, cup open, shorter than
the tube of the blossom.

Ditch Banks, June, P. Rare.

Banks of the Ouse.

Ditches on the Ival *Borage.*
near Biggleswade

V Males. I Female. Pen. Mo.

Borage. BORAGO. Gen. Pl. 248.

Blossom wheel-shaped: Mouth closed with rays.

147. *Common Borage.* — *B. officinalis.*

All the leaves alternate, cups expanding. — Sowerby. 247. 36.

Road-sides, May, A. Rare.

Duck Mill, Bedford, — Ampthill Warren.

Bugloss. LYCOPSIS. Gen. Pl. 250.

Blossom with a bent tube.

148. *Wild Bugloss.* — *L. arvensis.*

Leaves spear-shaped, rough with hair, cups when in flower erect. — Curtis. V. 17. Sowerby. 938.

Flora Danica. 435. admirable.

Cornfields, June, A. Common.

Viper's Bugloss. ECHIUM. Gen. Pl. 251.

Blossom irregular. Mouth naked.

149. *Common Viper's Bugloss.* — *E. vulgare.*

Stem rough with hairy tubercles, stem leaves spear-shaped, rough with hair, flowers in spikes lateral. — Martyn. 136. Flora Danica. 445.

Cornfields, June, B. Common.

V Males. I Female. Pen. Mo.

Primrose. PRIMULA. Gen. Pl. 258.

A small umbel with a fence. Tube of the blossom cylindrical, mouth expanding.

150. *Common Primrose.* — P. acaulis.

Stemless, leaves wrinkled, toothed, hairy underneath, stalk with one flower. — Curtis. VI. 16. Sowerby. 4. admirable.

Woods and Hedges, March, P. Common.

151. *Cowslip.* — P. officinalis.

Leaves wrinkled, toothed, hairy underneath, stalks with many flowers, nodding, border of the blossom short. — Curtis. VI. 15. Sowerby. V. admirable.

Pastures, April, P. Common.

152. *Oxlip.* — P. elatior. Sowerby. 513.

Leaves wrinkled, toothed, hairy, stalks with many flowers, outermost flower nodding, central one upright. — Flora Danica. 434.

Woods and Thickets, April, P. Common.

Buck-bean. MENYANTHES. Gen. Pl. 253.

Blossom hairy, stigma bifid, capsule 1-cell-ed.

153. *Common Buck-bean.* — M. trifoliata.

Leaves in threes. Curtis. IV. 17. Sowerby. 495.

Boggy Places, May P. Rare.

V. Males. I. Female. Pen. Mo.

*Paradise, near Bedford,—Biddenham South Field
opposite Kempston Church,—Ampthill Moor.*

Obs:

Blossom elegantly fringed, the red and white forming a most exquisite contrast.

Hottonia. **HOTTONIA.** Gen. Pl. 265.

Blossom salver-shaped, threads fastened to the tube of the blossom. Capsule 1-celled.

*54. *Water Violet.* — **H. palustris.**

Fruit-stalk whorled, with many flowers.—

Curtis. I. 11. admirable. Sowerby. 364.

Ditches, May, P. Rare.

Potton Marshes,—Eaton-Soccon Mills.

Loosestrife. **LYSIMACHIA.** Gen. Pl. 269.

Blossom wheel-shaped, capsule globose, pointed, with 10 valves.

*55. *Common Loosestrife.* — **L. vulgaris.**

Panicled, bunches terminal. — Curtis. V. 19.

Watery Places, July, P. Common. **Sowerby. 761.**

*56. *Wood Loosestrife.* — **L. nemorum.**

Leaves ovate, sharp, flowers solitary, stem trailing. — Curtis. V. 18. **Sowerby. 527.**

Woods, June. P. Rare.

Aspley Wood.

Obs:

V Males. I Female. Pen. Mo.

Obs:

Plant trailing, root creeping: Flowers large, deep yellow.
Capsule globular, without the spherical scissure, therefore
a true *Lysimachia*.

157. *Moneywort Loosestrife*. — *L. nummularia*.

Leaves nearly heart-shaped, flowers solitary,

Sowerby. 528. stem creeping. — Curtis. III. 14.

Wet Meadows and Ditches, June, P. Common.

Pimpernel. *ANAGALLIS*. Gen. Pl. 270.

Blossom wheel-shaped. Capsule cut round.

158. *Common Pimpernel*. — *A. arvensis*.

Leaves entire, stem trailing. — Curtis. I. 12.

Sowerby. 529. Cornfields, May, A. Common.

159. *Blue Pimpernel*. — *A. coerulea*.

Leaves between ovate and spear-shaped, seg-
ments of the cup more sharply pointed. —

Hoffman. Blackwell. 274.

Cornfields, May, A. Rare.

Oakley West Field, — *Barton Hill*.

Obs:

Petals nicked, sometimes undulated, and deeply serrated. It
were vain to attempt a description of the elegance of colour and
habit in this little rare Plant.

160. *Bog Pimpernel*. — *A. tenella*.

Leaves ovate, sharpish, stem creeping. —

Curtis. III. 15. Sowerby. 530.

Bogs

V Males. I Female. Pen. Mo.

Bog Pimpernel. — *A. tenella*.

Decumbent Bog-pimpernel. — *A. tenella*.

Root fibrous, leaves broad lanceolate, pointed at the center, and rather narrow at the base.

Common Pimpernel. — *A. arvensis*.

Blossom wheel-shaped. Capsule round, cut round.

Common Pimpernel. — *A. arvensis*.

Blossom bell-shaped, pointed. Capsule round, cut round.

Common Pimpernel. — *A. arvensis*.

Blossom bell-shaped, pointed. Capsule round, cut round.

Bog Pimpernel. — *A. tenella*.

Leaves lanceolate, pointed at the center, and rather narrow at the base.

Bog Pimpernel. — *A. tenella*.

Leaves lanceolate, pointed at the center, and rather narrow at the base.

Faint-scented Pimpernel. — *A. tenella*.

Leaves lanceolate, pointed at the center, and rather narrow at the base.

V. Males. I Female. I Pen. Mo.

Bogs, June, P. Rare.

Stevington Bogs, — Amptbill Moor.

Obs :

Root fibrous, branches produce small fibres towards the center, and take hold of the Ground. Leaves small, opposite, sharp, round. Flowers bell-shaped, small, flesh-colored.

Chironia. CHIRONIA. Gen. Pl. 349.

Blossom wheel-shaped. Female organ bowed down. Threads sitting on the tube of the blossom, anthers corkscrew-shaped, capsule 2-celled.

161. *Common Centaury.* — *C. Centaurium.*

Blossoms with 5 segments, funnel-shaped, stem forked, female organ single. — Curtis.

IV. 22. Sowerby. 417.

Pastures, July, A. Common.

Bindweed. CONVOLVULUS. Gen. Pl. 287.

Blossom bell-shaped, plaited. Stigmas 2. Capsule 2-celled, each 2-seeded.

162. *Field Bindweed.* — *C. arvensis.*

Leaves arrow-shaped, pointed on each side, fruitstalks with about 1 flower. — Curtis.

II. 13. Sowerby. 343. 312

Fields, June, P. Common.

Great

V Males. I Female. I Pen. Mo.

163. Great Bindweed. — *C. sepium*.

Leaves arrow-shaped, lopped behind, fruit-stalks square with 1 flower. — Curtis. I. 13. Sowerby. 313.

Hedges, July, P. Common.

Bell-flower. *CAMPANULA*. Gen. Pl. 290. Blossom bell-shaped, closed at the bottom by thread-bearing valves. Stigma trifid. Capsule inferior, opening by lateral orifices.

164. Round-leaved Bell-flower. — *C. rotundifolia*.

Root leaves kidney-shaped, stem leaves strap-shaped. — Curtis. IV. 21. Sowerby. 866.

Heaths and dry Pastures, June, P. Common.

165. Giant-Throatwort Bell-flower. — *C. latifolia*.

Leaves ovate-spear-shaped, stem very simple, round, flowers single, on fruit-stalks, capsules nodding. — Sowerby. 302.

Woods and Hedges, July, P. Rare.

Whipsnade near Dunstable, — Studham Common.

Obs:

Root milky, stem 5 feet high, slightly angular, crowded with leaves. Blossom splendid purple, leaves of the cup slightly serrated.

166. Throatwort Bell-flower. - *C. Trachelium*.

Stem angular, leaves on leaf-stalks, cups fringed, fruit-stalks in threes. — Sowerby. 12.

Woods

Woods and Hedges, July, P. Common.

167. Clustered Bell-flower. — *C. ciliolata*.

Stem angular, simple, upper part divided into several branches terminating in clusters of flowers. — Sowerby. 312. Common.

Heaths and dry Places, July, P. Common.

168. Grecian Bell-flower. — *C. grecia*.

Stem angular, branched, leaves opposite, pointed, lanceolate, longer than the flowers, petioles long.

Leaves often slightly curved, serrated.

Leaves often slightly curved, serrated.

Common in the neighbourhood.

165
in a lane between Hawnes
and Maudlin on the bank
of a copice on the left.

1807

V Males. I Female. Pen. Mo.

Woods and Hedges, July, P. Common.

167. *Clustered Bell-flower.* — *C. glomerata.*

Stem angular, simple, flowers sitting, head terminating. — Sowerby. 190. admirable.

Heaths and dry Places, July, P. Common.

168. *Corn Bell-flower.* — *C. hybrida.*

Stem somewhat branched at the base, straight, leaves oblong, crenated, cups incorporated, longer than the Blossom, capsules shaped like a prism. — Sowerby. 375. admirable.

Cornfields, June, A. Common.

Brookweed. *SAMOLUS.* Gen. Pl. 294.

Blossom salver-shaped. Threads defended by the scales of the blossom. Capsule 1-celled, inferior.

169. *Pimpernel Brookweed.* — *S. valerandi.*

Curtis. IV. 20. *Sowerby.* 703.

Banks of Rivers and Bogs, July, P. Rare.

Banks of the Ouse above Bedford.

Honeysuckle. *LONICERA.* Gen. Pl. 519.

Blossom of 1 petal, irregular : Berry with many seeds, 2-celled.

170. *Common Honeysuckle.* — *L. Periclymenum.*

Heads ovate, tiled, terminal, all the leaves distinct. — Curtis. I. 15. *Sowerby.* 800.

V Males. I Female. Pen. Mo.

Woods and Hedges, July, S. Common.

Mullein. VERBASCUM. Gen. Pl. 331.

Blossom wheel-shaped, nearly unequal, with
2 cells and 2 valves.

171. Great Mullein. — V. Thapsus.

Leaves running down the stem, woolly on
both sides: stem undivided. — Flora Da-

Sowerby. 549. nica. 631. Woodville. 125.

Road-sides, July, P. Common.

172. Black Mullein. — V. nigrum.

Leaves oblong, heart-shaped, on leafstalks.—
Sowerby. 59. admirable.

Hedges, July, P. Rare.

Sandy,—Warden,—Aspley.

Henbane. HYOSCYAMUS. Gen. Pl. 333.

Blossom funnel-shaped, blunt: threads inclin-
ing; capsule 2-celled, with a lid.

173. Common Henbane. — H. niger.

Leaves embracing the stem, waved, flowers
sitting. — Woodville. 52. admirable.

Road-sides, June, B. Common.

Deadly Nightshade. ATROPA. Gen. Pl. 335.

Blossom bell-shaped, threads distant. Berry
round, 2-celled.

Common

V Males. I Female. Pen. Mo.

174. *Common Deadly Nightshade.* — A. Belladonna.

Stem herbaceous, leaves ovate, very entire. —

Curtis. V. 16. Woodville. 1. Sowerby. 592.

Woods and Road-sides, June, P. Rare.

Near Whipsnade, on the borders of Herts.

Obs. 2.

They who have drawn this Plant have overleaped the modest simplicity of nature, for the flowers are of a dusky colour and not purplish brown; the leaves pale not a deep green. The style is but little longer than the tube of the blossom.

Nightshade. — SOLANUM. Gen. Pl. 537.

Blossom wheel-shaped. Anthers almost united, opening at the point with a double pore. Berry 2-celled.

175. *Woody Nightshade.* — S. Dulcamara.

Stem thornless, shrubby, zigzag, upper leaves spear-shaped. Bunches in half-umbels. —

Curtis. I. 14. Sowerby. 565.

Moist Hedges, July. S. Common.

176. *Garden Nightshade.* — S. nigrum.

Stem thornless, herbaceous, leaves ovate, tooth-angular, bunches in two rows, nodding. — Curtis. II. 14. Woodville. 226. Sowerby. 566.

Dunghills, July, A. Common.

V Male. I Female. Pen. Mo.

Buckthorn. RHAMNUS. Gen. Pl. 358.

Cup tubular, with scales defending the threads; Blossom none: Capsule a berry.

177. *Purging Buckthorn.* — *R. catharticus.*

Thorns terminating, flowers with 4 segments, dioecious, leaves ovate, stem upright. — Blackwell. 135. Sowerby. 1b29.

Woods, Hedges, May, S. Rare.

Clapham,—Ravensden.

178. *Berry-bearing Alder.* — *R. Frangula.*

Flowers with one female, florets hermaphrodite, leaves very entire. — Sowerby. 250. admirable.

Woods, Hedges, May, S. Rare.

Eversholt,—Aspley Wood.

Spindle-tree. EVONYMUS. Gen. Pl. 373.

Blossom 5 petals. Capsule 5 angles, 5 cells, 5 valves, colored: seeds with a veil.

179. *Common Spindle-tree.* — *E. europaeus.*

Flowers with mostly 4 segments, leaves sitting. — Sowerby. 362. admirable.

Woods, Hedges, May, S. Rare.

Renhold,—Cople,—Aspley.

Currant. RIBES. Gen. Pl. 390.

one of Sowerby's desiderata.

one of Sowerby's desiderata.

V Males. I Female. Pen. Mo.

Petals 5, and the threads inserted in the cup:
Style bifid, berry many-seeded, inferior.

180. *Black Currant.* — R. nigrum.

Thornless, bunches hairy, blossoms oblong. —
Flora Danica. 556. admirable. Sowerby. 1295.

Banks of Rivers, April, May, S. Rare.

Blunham.

Ivy. HEDERA. Gen. Pl. 395.

Petals 5 oblong. Berry with 5 seeds sur-
rounded by the cup.

181. *Common Ivy.* — H. Helix.

Leaves both ovate and lobed. — Curtis. I. 16. Sowerby. 1267.
Woods, Hedges, September, S. Common.

Periwinkle. VINCA. Gen. Pl. 419.

Contorted. Bags 2, upright. Seeds naked.

182. *Least Periwinkle.* — V. minor.

Stems trailing, leaves spear-ovate, flowers
on fruitstalks. — Curtis. III. 16. Sowerby. 917.

Woods, Hedges, April, P. Rare.

Potton, — Aspley.

183. *Great Periwinkle.* — V. major.

Stems erect, leaves ovate, slightly fringed. —
Curtis. IV. 19. Sowerby. 514.

Hedges, April, P. Rare.

Ravensden, — Clapham.

Goosefoot.

V Males. II Females. Pen. Di.

Goosefoot. CHENOPODIUM. Gen. Pl. 435.

Cup 5 leaves. Blossom none. Seed 1, lentil-shaped, superior.

184. *Mercury Goosefoot.* — C. bonus Henricus.

Leaves triangular, arrow-shaped, very entire, spikes compound, leafless, axillary. —

Curtis. V. 7. Sowerby. 1033. —

Dunghills and Church-Yards, June, P. Common.

185. *Upright Goosefoot.* — C. urbicum.

Leaves triangular, somewhat toothed, bunches crowded, very upright, lying close to the stem, very long. — Petiver. VIII. 8.

Road-sides, August, A. Rare.

St. Cuthbert's Bedford, behind the Church.

186. *Red Goosefoot.* — C. rubrum.

Leaves heart-three-cornered, bluish, toothed, bunches erect, compound, leafy, shorter than the stem. — Curtis. VI. 21.

Dunghills, August, A. Rare.

Goldington, — Aspley.

187. *Wall Goosefoot.* — C. murale.

Leaves ovate, shining, toothed, sharp, bunches branched, naked. — Curtis. VI. 20.

Dunghills, August, A. Common.

Fig-leaved

one of Sowerby's desiderata

*185. Bak-leaved Goosefoot. Ch. glaucum.

V Males. II Females. Pen. Di.

188. *Fig-leaved Goosefoot.* — *C. serotinum.*

Leaves trowel-shaped, indented and toothed, wrinkled, smooth, uniform, bunches terminating. — Curtis. II. 16. *C. viride.*

Dunghills, August, A. Common.

189. *White Goosefoot.* — *C. album.*

Leaves diamond-three-cornered, gnawed, entire behind, uppermost oblong, bunches upright. — Curtis. II. 15. *C. supina.*
Cornfields, July, A. Common.

190. *Thorn-apple-leaved Goosefoot.* — *C. hybridum.*

Leaves heart-shaped, pointed-angular, bunches branched, naked. — Curtis. IV. 23.

Dunghills, August, A. Rare.

Mill Lane, Bedford.

Elm. *ULMUS.* Gen. Pl. 443.

Cup of 5 segments. Blossom none. Berry dry, flattened, skinny.

191. *Common Elm.* — *U. campestris.*

Leaves doubly-serrated, unequal at the base, flowers nearly sitting, in clusters.

Hedges, April, T. Common.

192. *Wycke-hazel Elm.* — *U. effusa.*

Leaves doubly-serrated, unequal at the base, flowers scattered, on long fruitstalks.

Hedges,

V Males. II Females. Pen. Di.

Hedges, April, T. Common.

Gentian. *GENTIANA*. Gen. Pl. 450.

Blossom 1 petal. Cup 2 valves, 1 cell, with 2 longitudinal receptacles.

193. *Fellwort Gentian*. — *G. Amarella*.

Blossoms with 5 segments, salver-shaped, bearded at the mouth, segments of the cup equal. — Flora Danica. 328. Sowerby. 236. admirable.

Pastures, July, A. Rare.

Turvey, opposite Snelsor, — Bromham.

194. *Field Gentian*. — *G. campestris*.

Blossoms with 4 segments, salver-shaped, bearded at the mouth, segments of the cup unequal. — Flora Danica. 367. Sowerby. 237. admirable.

Hilly Pastures, September, A. Very rare.

Barton Hill.

Obs:

The Synonym expresses that the segments of the cup are unequal, the alternate ones are particular in this respect, the one being strap-shaped, the other ovate-spear-shaped.

Umbellated Plants.

A. with general and partial Fences.

Pennywort. *HYDROCOTYLE*. Gen. Pl. 457.

Umbel

V Males. II Females. II Pen. Di.

Umbel simple, with a fence of 4 leaves,
petals entire, seeds semicircular, flattened.

195. *Marsh Pennywort*. — *H. vulgaris*.
Leaves shield-shaped, umbels with 5 flowers.
— Curtis. VI. 19. Flora Danica. 90.
Bogs, June, P. Rare. *Sowerby*. 751.

Ampthill, — *Aspley*.
Sanicle. *SANICULA*. Gen. Pl. 458.

Umbels crowded, almost in heads. Fruit
rough; fruit of the disc abortive.

196. *Common Sanicle*. — *S. europaea*.
Root leaves simple, all the flowers sitting. —
Sowerby. 98. Flora Danica. 283. admirable.

Woods and Thickets, May, P. Common.

Thorow-wax. *BUPLEURUM*. Gen. Pl. 460.
Fences of the small umbel larger than the
flowers, of 5 leaves; Petals rolled in. Fruit
nearly round, flattened, striped.

197. *Common Thorow-wax*. — *B. rotundifolium*.
No general fences: Stem perforating the
leaves, and growing through them. — Black-
well. 95. Sowerby. 99. admirable.

Cornfields, June, A. Rare.

Barton Hill, — *Woburn*.

V Males. II Females. Pen. Di.

Hensfoot. CAUCALIS. Gen. Pl. 464.

Blossoms with a ray, male florets in the disc.
Petals bent in and nicked. Fruit with
bristles, fences entire.

198. *Fine-leaved Hensfoot.* — *C. daucoides.*

Umbels in threes, leafless, small umbels with
3 seeds and 3 leaves. — Sowerby. 197. ad-
mirable.

Cornfields, June, A. Very rare.

Barton Hill Oakley West Field, discovered by Mr. James
1004 Payne, a discerning Herbarist of that Village.

Obs:

Leaves very finely divided, Germs armed with hooked
bristles, Flowers pinkish-white, Petals uniform.

199. *Broad-leaved Hensfoot.* — *C. latifolia.*

General umbel trifid, partial ones with 5
seeds, leaves winged and serrated. — Sow-
erby. 198. admirable.

Cornfields, June, A. Rare.

Oakley, — Thurleigh. — Obs:

Petals irregular, blush-colored. Seeds hooked as in the
preceding.

200. *Field Hensfoot.* — *C. arvensis.*

V Males. II Females. Pen. Di.

General fence of one leaf, umbels crowded,
seeds round-ovate, leaflets ovate-spear-shaped,
doubly winged, branches straggling. —

Curtis. VI. 23. C. infesta. **Sowerby. 1314.**

Cornfields, July, A. Common.

201. *Hedge Hensfoot.* — C. Anthriscus.

Fences with many divisions, umbel crowded,
seeds oblong, leaflets ovate, doubly-winged,
branches erect. — Curtis. VI. 22. **Sowerby. 987.**

Hedges, July, A. Common.

202. *Knotted Hensfoot.* — C. nodosa.

Umbels lateral, simple, sitting. — Sowerby.
199. admirable.

Road-sides, June, A. Common.

Carrot. DAUCUS. Gen. Pl. 466.

Blossoms nearly radiate, all with both male
and female organs. Fruit rough with hair.
Fences doubly-winged.

203. *Wild Carrot.* — D. Carota.

Seeds rough, leafstalks ribbed underneath. —

Martyn. 82. admirable. **Sowerby. 1174.**

Pastures, July, B. Common.

Earth-nut. BUNIUM. Gen. Pl. 468.

Blossom uniform, umbel crowded, fruit ovate.

V Males. II Females. Pen. Di.

204. *Small Earth-nut.* — *B. flexuosum.*

Stem tapering at bottom, zigzag. — Curtis.

IV. 24. admirable. *Sowerby*. 988.

Pastures and Thickets, May, P. Rare.

King's Wood, Ampthill, — Aspley Wood.

Obs:

The true *Bulbocastanum*, (well figured by Oeder) is not, I believe, a native of this County. The plant before us is truly zigzag towards the root, and the stem tapering at the base.

Hemlock. CONIUM. Gen. Pl. 469.

Partial fences going half way round, generally of 3 leaves. Fruit nearly globular, with 5 scores, notched on each side.

205. *Common Hemlock.* — *C. maculatum.*

Seeds scored. — Curtis. I. 17. admirable.

Road-sides, June, B. Common.

Sulphurwort. PEUCEDANUM. Gen. Pl. 472.

Fruit ovate, scored on both sides, bordered, fences very short.

206. *Common Sulphurwort.* — *P. Silaus.*

Leaflets doubly-winged, segments opposite, general fence of 2 leaves. — Martyn. 128. admirable.

Moist Pastures, July, P. Common.

Hogweed.

V Males. II Females. Pen. Di.

Hogweed. HERACLEUM. Gen. Pl. 477.

Fruit elliptical, nicked, flat, scored; Bloom irregular, bent in and nicked, fence falling off.

207. *Common Hogweed.* — H. Sphondylium.

Leaves doubly-winged, smooth, flowers regular. — Petiver. XXIV. 1. **Sowerby.** 939.

Hedges, July, P. Common.

208. *Narrow-leaved Hogweed.* — H. angustifolium. *one of Dr. Smith's desiderata.*

Leaves with a cross in the wing, leaflets spear-shaped, blossoms with florets. — Jacquin. Flora Aust. 174.

Hedges, July, B. Rare.

Elsow.

Obs:

I am the more convinced of the identity of this species from the circumstance of the Rev. Mr. Relhan's having shewn me, the corresponding Plant page 10 of his second supplement.

Angelica. ANGELICA. Gen. Pl. 479.

Fruit nearly round, angular, firm, with reflected styles. Blossoms equal, petals bent in.

209. *Wild Angelica.* — A. sylvestris.

Leaflets equal, ovate-spear-shaped, serrated. — Woodville. 265. **Sowerby.** 1128.

Woods
for 109 read 209.

V Males. II Females. Peh. Di.

Woods and wet Hedges, July, B. Common.

Water Parsnep. Sium. Gen. Pl. 480.

Fruit rather egg-shaped, scored: fence of many leaves. Petals heart-shaped.

210. *Broad-leaved Water Parsnep.* — S. latifolium.

Leaves winged, umbel terminal. — Sowerby.

204. admirable.

Ditches and Banks of Rivers, July, P. Common.

211. *Narrow-leaved Water Parsnep.* — S. angustifolium.

Leaves winged, umbels axillary on fruitstalks, general fence winged. — Sowerby. 139. admirable.

Rivulets and Ditches, July, P. Rare.

With the above.

Obs:

Leaves deeply serrated, leaves of the fences and fencelets very large and conspicuous.

212. *Trailing Water Parsnep.* — S. nodiflorum.

Leaves winged, umbels lateral, nearly fitting, stem trailing. — Flora Danica. 247. admirable. Sowerby. 639.

Rivulets and Ditches, July, P. Common.

Creeping

V Males. II Females. Peh. Di.

213. *Creeping Water Parsnep.* — S. rotundifolium.

Stem creeping, leaflets rounded, thick, acute, lax. — Jacob's Almanac. 280. 320. 370.

Flowers June, July.

Groundsel-like.

Broad-leaved Parsnep. Gen. Pl.

Fruit oval, scored, fence common, winged.

214. *Creeping Parsnep.*

Leaves winged, umbels terminal, fence common, lax, spreading.

215. *Creeping Water Parsnep.* — S. rotundifolium.

Leaves winged, umbels descending. — Sowerby. 638. admirable.

216. *Creeping Water Parsnep.* — S. rotundifolium.

Leaves winged, umbels descending. — Sowerby. 638. admirable.

217. *Creeping Water Parsnep.* — S. rotundifolium.

Leaves winged, umbels descending. — Sowerby. 638. admirable.

218. *Creeping Water Parsnep.* — S. rotundifolium.

Leaves winged, umbels descending. — Sowerby. 638. admirable.

V Males. II Females. Pen. Di.

Bogs, June, A. Rare.

Amphibolus

Obs :

The leaves out of the water winged, serrated: Those that grow beneath the surface are very fine and hair-like. Blossom irregular.

Dropwort. *OENANTHE*. Gen. Pl. 484.

Florets dissimilar, those in the disc barren, sitting: Fruit crowned with a cup and female organ.

217. Water Dropwort. — *O. fistulosa*.

Root sending forth suckers, stem-leaves winged, thread-shaped, hollow. — Sowerby. 363. admirable.

Boggy Ground, June, P. Common.

218. Sulphurwort-leaved Dropwort. — *O. peucedanifolia*.

All the leaves strap-shaped, root leaves with a double row of wings, stem-leaves winged, no general fence, knobs of the root ovate, sitting. — Sowerby. 348. admirable.

Marshes, June, P. Rare.

Potton Marshes, — Fenlake Meadow.

B. with fencelets, but without fences.

Horse-bane.

219* Parsley leaved Dropwort.
O. primpinelloides.

V Males. II Females. Pen. Di.

Horsebane. PHELLANDRIUM. Gen. Pl. 485.

Florets of the center less, fruit ovate, smooth, crowned with a cup and female organ.

219. *Water Horsebane.* — *P. aquaticum.*

Branches of the foliage straggling. — Woodville. 266. admirable. Sowerby. 684.

Rivers, July, P. Common.

Cowbane. CICUTA. Gen. Pl. 486.

Fruit nearly ovate, furrowed.

220. *Poisonous Cowbane.* — *C. virosa.*

Flora Danica. 208. Woodville. 268. Sowerby. 479.

Ditches, July, P. Rare.

Oakley Springs.

Cicely. ÆTHUSA. Gen. Pl. 487.

Fencelets going half round, with three leaves, hanging down : Fruit scored.

221. *Fool's Parsley.* — *Æ. Cynapium.*

Curtis. I. 18.

Gardens and Fields, July, P. Common.

Shepherd's Needle. — SCANDIX. Gen. Pl. 489.

Blossom radiate, fruit awl-shaped, petals nicked, florets of the disc often males only.

V Males. II Females. Pen. Di.

222. *Common Shepherd's-Needle*. — S. pecten.
Seeds with a very long beak. — Curtis. V.

Sowerby. 1397. 21. Martyn. 38.
Cornfields, May, A. Common.

223. *Rough Shepherd's-Needle*. — S. Anthriscus.
Seeds ovate, rough with hair, blossoms uniform, stem smooth. — Curtis. I. 19. Martyn. 75. *Sowerby. 818.*
Road-sides. May, A. Common.

224. *Chervil Shepherd's-Needle*. — S. Cerefolium.
Seeds glossy, on long fruitstalks. — Blackwell. 236.
Road-sides, May, A. Rare.

Goldington Road-side.

Chervil. CHAEROPHYLLUM — Gen. Pl. 490.
Fence bent back, concave: Petals heart-shaped, bent in. Fruit oblong, smooth.

225. *Common Chervil*. — C. sylvestre.
Stem smooth, scored, knees swollen. — Curtis. IV. 25. Martyn. 96. *Sowerby. 752.*
Pastures under Hedges, May, P. Common.

226. *Small Chervil*. — C. temulentum.
Stem rough, knees swollen. — Curtis. VI.

24. Sowerby. 1521.

Under Hedges, June, B. Common.

C. without

A. with. II. Smooth. Petal.

C. without petiole or fence.

Fence. Pasture. Gen. Pl. 490.

Fruit elliptical, con-pressed flat: Petals
entire, rolled in.

227. *Common Chervil*. — P. corymbosa.

Leaves deeply winged. — Martyn. 96. *Sowerby. 556.*

Stems several, upright.

Leaves few, narrow,

and oblong pointed, smooth.

Fruit oblong, smooth.

Leaves few, narrow,

and oblong pointed, smooth.

V Males. II Females. Pen. Di.

C. without fences or fencelets.

Parsnep. PASTINACA. Gen. Pl. 494.

Fruit elliptical, compressed, flat : Petals entire, rolled in.

227. *Wild Parsnep.* — P. sylvestris.

Leaves simply winged. — Martyn. 83. Sowerby. 556.
Road-sides, July, B. Common.

Alisanders. SMYRNIUM. Gen. Pl. 495.

Fruit oblong, scored ; Petals pointed, keeled.

228. *Common Alisanders.* — S. Olus atrum.

Stem-leaves in threes, on leafstalks, serrated.—
Sowerby. 230. admirable.

Under Hedges, May, B. Rare.

Elstow, — Ravensden, — Oakley.

Dill. ANETHUM. Gen. Pl. 364.

Fruit nearly ovate, flat, scored : Petals rolled in, entire.

229. *Fennel.* — A. Foeniculum.

Woodville. 160. Sowerby. 1208 — .

Springs and chalky Soils, July, B. Rare.

Eversholt, in a deep Ditch near the Church.

Carraways. CARUM. Gen. Pl.

Fruit ovate-oblong, scored ; Fence of 1 leaf.
Petals keeled, bent in, nicked.

V Males. II Females. Pen. Di.

230. *Common Carraways.* — C. Carui.

Sowerby. 1503. Martyn. 55.

Pastures, May, B. Rare.

Thurleigh.

Obs :

This Plant was first shewn to me by John Payne,
brother of James Payne, the Oakley Herbarist.

Burnet Saxifrage. PIMPINELLA. Gen. Pl. 498.

Fruit ovate-oblong. Petals bent in. Stigmas
roundish.

231. *Small Burnet Saxifrage.* — P. Saxifraga.

Leaves winged, root leaves nearly round, up-
per ones strap-shaped. — Martyn. 127.
Sowerby. 407. admirable.

Dry Pastures, July, P. Common.

232. *Jagged Burnet Saxifrage.* — P. dissecta.

All the leaves winged, wings with many di-
visions, segments somewhat leafy, pointed.—
Retz. Fasc. III. fig. 2.

Road-sides, July, P. Common.

233. *Great Burnet Saxifrage.* — P. magna.

All the leaves lobed, the odd lobe divided in-
to three. — Sowerby. 408.

Woods, July, P. Common.

Parsley.

V Males. II Females. Pen. Di.

Parsley. APIUM. Gen. Pl. 499.

Fruit ovate, scored. Fence of 1 leaf, petals equal.

234. *Smallage.* — A. graveolens.

Stem leaves wedge-shaped. — Flora Danica.

790. *Sowerby.* 1210.

Wet Ditches, June, P. Rare.

Medbury, — Wilshamsted, — Goldington.

Goutweed. ÆGOPODIUM. Gen. Pl. 500.

Fruit ovate-oblong, scored.

235. *Herb Gerard.* — Æ. Podagraria.

Upper leaves in threes. — Flora Danica. 670. *Sowerby.* 940.

Woods, Hedges, June, P. Common.

V Males. III Females. Pen. Tri.

Mealy-tree. VIBURNUM. Gen. Pl. 503.

Cup 5 segments. Blossom 5 divisions. Berry 1 seed.

236. *Pliant Mealy-tree.* — V. Lantana.

Leaves heart-shaped, serrated, veined, woolly beneath. — Sowerby. 331. admirable.

Woods, Hedges, May, S. Common.

237. *Guelder Rose.* — V. Opulus.

Leaves lobed, leafstalks with glands. — Sowerby. 332. admirable.

Weeds,

V Males. III Females. Pen. Tri.

Woods, Hedges, May, S. Common.

Elder. SAMBUCUS. Gen. Pl. 505.

Cup 5 segments. Blossom 5 divisions. Berry
3 seeds.

238. *Dwarf Elder.* — S. Ebulus.

Sowerby. 475. Cymes with 3 props, stipulæ leafy, stem her-
baceous. — Curtis. III. 18. Woodville.
260.

Hedges, July, S. Common.

239. *Common Elder.* — S. nigra.

Cymes with 5 props, stem woody. — Flora
Danica. 545. Sowerby 476.

Hedges, June, S. Common.

Chickweed. ALSINE. Gen Pl. 518.

Cup of 5 leaves. Petals 5, equal. Capsule
1 cell, 3 valves.

240. *Common Chickweed.* — A media.

Aellenia media. Petals nicked, leaves ovate-heart-shaped. —
Curtis I. 20. admirable. Sowerby. 537.

Fields, Gardens, April, A. Common.

V Males. IV Females. Pen. Tetr.

Grass of Parnassus. PARNASSIA. Gen. Pl. 523.

Cup 5 segments. Petals 5. Nectaries 5,
heart-shaped, fringed, points globular. Cap-
sule 4 valves.

V Males. IV Females. Pen. Tetr.

241. *Marsh Parnassia*. — *P. palustris*.
Sowerby. 82. Flora Danica. 584. admirable.
Bogs, July, P. Rare.

Stevington. — *Turvey*, — *Amptbill*.

V Males. V Females. Pen. Pen.

- Flax*. *LINUM*. Gen. Pl. 528.
Cup 5-leaved, petals 5. Capsule 5 valves,
10 cells, seeds solitary.

242. *Common Flax*. — *L. usitatissimum*.

Cups and capsules pointed, petals slightly
notched, leaves lance-shaped, alternate, stem
nearly single. — Martyn. 133. Woodville.

111. Sowerby. 1357.
Cornfields, July, A. Rare.

Barton Hill.

243. *Purging Flax*. — *L. catharticum*.

Leaves opposite, ovate-lance-shaped, stem
forked, blossoms pointed. — Curtis. III.
19. Martyn. 135. Sowerby. 382.

Pastures, June, A. Common.

Sundew. *DROSERA*. Gen. Pl. 531.

Cup of 5 divisions, Petals 5. Capsule 1-
celled, opening with 5 valves: Seeds many.

244. *Round-leaved Sundew*. — *D. rotundifolia*.

Stems

V Males. V Females. Pen. Pen.

Stems from the root, leaves round — Flora Danica. 1028. **Sowerby**. 867.

Bogs, July, P. Rare.

Ampthill.

245. *Long-leaved Sundew*, — *D. longifolia*.

Stems from the root, leaves oval-oblong. — Petiver. LXIII. 11. **Sowerby**, 868.

Bogs, July, P. Rare.

Ampthill Bogs.

246. *Great Sundew*. — *D. anglica*.

Leaves oblong, stems from the root, eight female organs in the flower. — Petiver. LXIII. 12. **Sowerby**. 869.

Bogs, July, P. Extremely rare.

Ampthill Bogs.

V Males. Many Females. Pen. Poly.

Mousetail. *MYOSURUS*. Gen. Pl. 538.

Cup 5-leaved, growing together at the base, nectaries 5, awl-shaped, like petals; Seeds numerous.

247. *Least Mousetail*. — *M. minimus*.

Sowerby. 435. Flora Danica, 406, Curtis. IV. 26.
admirable. Cornfields, June, A. Rare.

Biddenham, — *Fenlake*, — *Caldwell*.

Obs:

V Males. Many Females. Pen. Poly.

Obs:

No one has noticed the natural simplicity and elegance of this little plant, and less are even the seeds and parts of fruit & flowers, before the seed-vessel is opened. The reason of the male flower of the flower being open, is that it is the time of flowering.

Obs: A. S. S. W.

V Males. Many Females. Pen. Poly.

Sundew. *GYMNOGYNE*. Gen. Pl.

Petals 5, equal, 1 awl-shaped.

petals awl-shaped, alternate.

248. *Common Sundew*. — *G. Sundew*.

Sowerby. 870.

Bottom of Meadows, April, P. Rare.

Sandy Hills.

Orchard, *Chesham*, *Wotton*, *St. Albans*.

Six petals, equal; flower-cup funnel-shaped,
of 1 leaf. Threads within the flower-cup.

249. *Small Sundew*. — *N. Sundew*.

Sheath with ~~petals~~ — *Grass*, *scrub*, *heath*.

erect, simple, *calyx* *white* *pink*.

Sowerby. 871.

Woods, *Wotton*, *St. Albans*, *W. Herts*.

Cards

V. Males. Many Females. Pen. Poly.

Obs:

No one has observed the natural simplicity and elegance of this little plant, who has not seen the petals and parts of fructification, before the seedspike is produced. The organs of the male division of the flower often far exceed the limits and number of the class.

C L A S S. VI.

VI Males. I Female. Hex. Mo.

Snow-drop. *GALANTHUS.* Gen. Pl.

Petals 3, concave. Honey-cup of 3 small petals, nicked, summit simple.

248. Common Snowdrop. — *G. nivalis.*

Sowerby. 19.

Bottom of Mountains, April, P. Rare.

Sandy Hills.

Daffodil. *NARCISSUS.* Gen. Pl. 550.

Six petals, equal : Honey-cup funnel-shaped, of 1 leaf. Threads within the honey-cup.

249. Pale Daffodil. — *N. pseudo-narcissus.*

Sheath with 1 flower, nectary bell-shaped, erect, crimp, equalling the ovate petals. — Sowerby. 17.

Woods and Hedges, April, P. Rare.

Clapham,—Warden.

L

Garlick.

249.

Melchbourne Wood - 1799.

Rt Hon Mr L. St John

Botanica 300, Class IV, 66.

June, A. Rare.

Malvern—Preston—Cotswold.

VI Males. I Female. Hex. Mo.

Garlick. ALLIUM. Gen. Pl. 557.

Blossom of 6 divisions, expanding. Sheath of many flowers. Umbel crowded. Capsule superior.

250. *Crow Garlick.* — A vineale.

Stem with cylindrical leaves and bulbs, threads with 3 points. — Petiver. LXVI. 1.

Meadows, June, P. Common.

251. *Ramson Garlick.* — A. ursinum.

Stalk naked, three-square, leaves spear-shaped, on leafstalks, umbel flat at the top. — Sowerby. 122. admirable.

Moist Woods, May, P. Rare.

Whipsnade, near Dunstable.

Hyacinth. HYACINTHUS. Gen. Pl. 577.

Blossom bell-shaped. 3 honey-cup pores at the point of the germ.

252. *Hare bells.* — H. non-scriptus

Blossoms with 6 segments, bell-shaped, turned back at the point. — Curtis. II. 18. Sowerby. 377.

Woods, May, P. Common.

Tulip. TULIPA. Gen. Pl.

Blossom of 6 petals, bell-shaped: style none.

Wild

VI Males. I Female. Hex. Mo.

253. *Wild Tulip.* — *T. sylvestris.*

Flower solitary, somewhat drooping, leaves lance-shaped, stigma obtuse triangular, stamens hairy at the base. — (*English Botany.*) Flora Danica. 375. Sowerby. 63. admirable.

Old Chalkpits, April, P. Rare.

Whipsnade. on the borders of Hertfordshire, Rev. D. Jenks.

Obs:

Outer coat of the obovate bulb tawny. Tips of the petals hairy.

Fritillary. *FRITILLARIA.* Gen. Pl. 559. Blossom of 6 petals, bell-shaped, with a cavity containing honey above the claws. Threads as long as the blossom.

254. *Checkered Fritillary.* — *F. Meleagris.*

All the leaves alternate, stem with 1 flower. Curtis. III. 20. Sowerby. 622.

Moist Meadows, April, P. Rare.

Bromham.

Star of Bethlehem. *ORNITHOGALUM.* Gen. Pl. 566.

Blossom of 6 petals, upright, continuing, expanding more than half way. Threads alternate, broad at the base.

L 2

Common

VI Males. I Female. Hex. Mo.

255. Common Star of Bethlehem. — *O. umbellatum.*

Flowers in flat-topped spikes, fruitstalks higher than the stem, threads broadest at the base. — Sowerby. 130.

Pastures and Heaths, May, P. Rare.

Everton Heath.

256. Spiked Star of Bethlehem. — *O. pyrenaicum.*

Bunch of flowers very long, threads spear-shaped, flower-bearing fruitstalks expanding, equal : when in seed approaching the stalk. — Johnson's Gerard. 97. *Sowerby 499.*

Pastures and Hedges, June, P. Rare.

Between Eaton-Socon and Thurleigh.

Convally. CONVALLARIA. Gen. Pl. 575.

Blossom of 6 segments; Berry spotted, 3 cells.

257. Lilly of the Valley. — *C. maialis.*

Stalk bare, half-rounded, leaves elliptical. — (Aiton.) Curtis. V. 24. (See our fig. No. 2.) *Sowerby 1035.*


Woods, May, P. Rare.

Aspley Wood.

Obs:

The leaves are rather egg-spear-shaped than elliptical.

Solomon's



VI Males. I Female. I Hex. Mo.

258. *Solomon's Seal.* — *C. multiflora.*

Leaves alternate, embracing the stem, stem round, fruitstalks from the bosom of the leaves, bearing many flowers. — Sowerby.

279.

Woods, May, P. Rare.

Thurlegh Wood.

Spiderwort. *ANTHERICUM.* Gen. Pl.

Blossom of 6 petals, expanding. Capsule ovate.

259. *Lancashire Asphodel.* — *A. ossifragum.*

Leaves sword-shaped, threads woolly. — Flora Danica. 42. Sowerby. 535.

Bogs, July, P. Rare.

Ampthill Bogs.

Reed. *ACORUS.* Gen. Pl. 586.

Spike cylindrical, covered with florets, Blossoms (having 6 petals) naked, style none. Capsule 3-celled.

260. *Sweet Reed.* — *A. Calamus.*

Sowerby, 356. Woodville, 173.

Stagnant Waters, July, P. Rare.

Moat, at the Hassocks Farm.

Purslane. *PEPLIS.* Gen. Pl. 605.

Cup

VII Males. I. Female. Hex. Mo.

Cup bell-shaped, orifice with 12 clefts. Petals 6, inserted in the cup, capsule 2-celled.

261. *Water Purslane*. — P. Portula.

Stem creeping, flowers in the wings of the leaves, petals very minute and fugacious. — Curtis. IV. 27. admirable. Flora Dani-
ca. 64. Sowerby. 4211.

Bogs, July, A. Rare.

Amphibill Bogs.

Obs:

The plant is absolutely prostrate, bearing reddish flowers sit-
ting at every joint: The allotted number of rose-colored petals
is seldom compleat.

Rush. JUNCUS. Gen. Pl. 590.

Cup of 6 leaves. Blossom none. Capsule
1-celled.

262. *Clustered Rush*. — J. conglomeratus.

Stalk naked, upright, head on one side. —
Leers. XIII. 1. Sowerby. 835.

Meadows and Heaths, July, P. Common.

263. *Soft Rush*. — J. effusus.

Stalk naked, upright, panicle on one side. —

Sowerby. 836. Leers. XIII. 2. ~~Sowerby 836~~

Road-sides, July, P. Common.



264. *Hard Rush*. — J. inflexus. S.D. *dent*

Stalk

VII Males. I. Female. Hex. Mo.

Stalk naked, upright, head on one side. —
Leers. XIII. 2. Sowerby. 836.

Heaths and Bogs, July, P. Common.



VI Males. I Female. I Hex. Mo.

Stalk naked, scored, of a sea-green color, leaf-scales brown at the base, panicle on one side, scattered. — Leers. XIII. 3. Sowerby. 665.

Heaths and Pastures, July, P. Common.

265. Moss Rush. — *J. squarrosum*.

Stalk naked, leaves bristle-shaped, heads leafless, clustered. — Flora Danica. 430. Sowerby. 933.

Heaths, July, P. Rare.

Ampthill Heath.

266. Flat-stalked Rush. — *J. compressus*.

Stalk leafy, trailing, leaves flattened, jointed with knots, panicle compound. — Sowerby.

238.

Wet Places, July, P. Common.

267. Viviparous Rush. — *J. viviparus*.

Stalk branched, leaves bristle-shaped, flowers in bundles, sitting upon the branches.

(Haller.) Plukenet. XXXII. 3. Sowerby. 801.

Bogs, July, P. Rare.

Ampthill Bogs.

268. Bulbous Rush. — *J. bulbosus*. S. D. tent.

Leaves strap-shaped, channelled, capsules blunt. — Flora Danica. 431. Sowerby. 934.

Pastures and Heaths, July, P. Rare.

Ampthill Bogs, — Potton Marshes.

Toad

266.

* Dwarf Meadow Rush. — *J. uliginosus*.

Leaves like bristles, stem upright, transversely jointed, panicle diffuse and compound —

Pastures and Commons. June. P. Rare.

Friars Pastures.

Obs:

This plant does not in its outward appearance bear the smallest resemblance to the *J. compressus* No. 266. of this work, its stem

and leaves are perfectly cylindrical, nor can you perceive the smallest trace of a knot in the whole plant, but when you hold it up to the light, the transverse partitions may be observed —

VI Males. I Female. Hex. Mo.

269. *Toad Rush.* — J. bufonius.

Stem forked, leaves angular, flowers single, sitting. — Leers. XIII. 8. Sowerby. 802.

Wet Places, July, A. Common.

270. *Hairy Wood Rush.* — J. pilosus.

Leaves flat, hairy, flat-topped spike branched, flowers single. — Curtis. V. 25.

Woods, April. P. Common.

271. *Great Hairy Wood Rush.* — J. sylvaticus.

Leaves flat, hairy, flat-topped spike compound, flowers in bundles, sitting. — Curtis.

V. 26.

Woods, May, P. Common.

272. *Hairy Field Rush.* — J. campestris.

Leaves flat, hairy, spikes sitting and on fruit-stalks. — Curtis. II. 19. Sowerby. 672.

Pastures, April, P. Common.

Barberry. BERBERIS. Gen. Pl. 595.

Cup of 5 leaves. Petals 6, with 2 glands at the claws. Style none. Berry 2-seeded.

273. *Common Barberry.* — B. vulgaris.

Fruitstalks in bunches. — Sowerby. 49.

Woodville. 234.

Hedges, May, S. Rare.

Clapham Lane, — Milton Ernys.

Dock.

VI Males. III Females. Hex. Tri.

Dock. RUMEX. Gen. Pl. 613.

Cup 3 leaves. Petals 3, approaching, seed
1 three-square.

274. Bloody Dock. — *R. sanguineus.*

Leaves without red veins. — (Stokes) Curtis.

III. 21. Sowerby. 724.

Woods and Road-sides, July, P. Common.

275. Curled Dock. — *R. crispus.*

Flowers with both male and female organs,
valves entire with grains, leaves spear-
shaped, waved, pointed. — Curtis. II. 20.

Pastures and Road-sides, July, P. Common.

276. Golden Dock. — *R. aureus.*

Whorls crowded, when ripe resembling a
spike, flowers golden yellow. — Petiver. II.

8. Raii. p. 142. *Lapathum aureum.* Sowerby. 725.

Marshes, July, P. Rare.

Goldington Green.

277. Blunt-leaved Dock. — *R. obtusifolius.*

Flowers with both male and female organs,
valves toothed, with grains, leaves heart-
shaped, oblong, bluntnish, crenated. — Cur-
tis. III. 22.

Road-sides, July, P. Common.

M

Fiddle

Obs:

The valve with its lateral fringes
forms a not inapt, tho' distant
representation of a Spider.

VI Males. III Females. Hex. Tri.

278. *Fiddle Dock*. — *R. pulcher*.

Sowerby. 1576. Flowers with both male and female organs,
valves toothed, nearly simple, with a grain,
root leaves fiddle-shaped. — Petiver. II. 10.

Church-yards and Road-sides, July, P. Common.

279. *Water Dock*. — *R. Hydrolapathum*.

Flowers with both male and female organs,
valves entire with grains, leaves pointed,
lance-shaped. — Woodville. 178.

Ditches and Rivers, July, P. Common.

280. *Sheep's Sorrel*. — *R. Acetosella*.

Flowers male and female on separate plants,
leaves between spear and halberd-shaped. —

Sowerby. 1674. Curtis. V. 29.

Heaths, July, P. Common.

Obs :

*Plant diaceous. Root brown, leaves mostly halberd-shaped,
often turned inwards. Habit of the plant so various as some-
times to be no more than 3 inches high, in a richer soil it rises
to 8 inches, or even a foot.*

281. *Common Sorrel*. — *R. Acetosa*.

Flowers male and female on separate plants,
leaves oblong, arrow-shaped. — Sowerby.

127.

Pastures, June, P. Common.

Arrow-Grass.

VI Males. III Females. Hex. Tri.

Arrow-Grass. **TRIGLOCHIN.** Gen. Pl. 166.
Cup of 3 leaves. Petals 3, cup-shaped. Style
none. Capsule opening at the base.

282. *Marsh Arrow-Grass.* — *T. palustre.*
Capsules with 3 cells, nearly strap-shaped. —
Sowerby. 366. Flora Danica. 490.

Marsches, July, P. Rare.

Hassock's Meadow, — *Ampthill Bogs.*

Meadow Saffron. **COLCHICUM.** Gen. Pl. 621.
A sheath. Blossom with 6 segments, the
tube coming from the root. Capsules 3,
connected, inflated.

283. *Common Meadow Saffron.* — *C. autumnale.*
Leaves flat, spear-shaped, straight. — Sow-
erby. 133. Woodville. 177. Martyn. 60.
Wet Meadows, September, P. Rare.

Barford, — *Thurleigh.*

VI Males. Many Females. Hex. Poly.

Water Plantain. **ALISMA.** Gen Pl. 625.

Cup of 5 leaves. Petals 3. Seeds many.

284. *Great Water Plantain.* — *A. Plantago.*
Leaves ovate, sharp, seeds 3-square, blunt. —
Curtis. V. 27. Flora Danica. 561. *Sowerby.* 837.
Ditches, July, P. Common.

M 2 Small

VI Males. Many Females. Hex. Poly.

285. *Small Water Plantain.* — *A. ranunculoides.*

Leaves strap-spear-shaped, fruit round, scurfy. — Sowerby. 326. Flora Danica. 122.

Bogs, June, P. Rare.

Amphill Bogs.

C L A S S. VIII.

VIII. Males. I Female. Oct. Mo.

Willow Herb. EPILOBIUM. Gen. Pl. 639.

Cup 4 segments. Petals 4. Capsule oblong, inferior seeds downy.

286. *Rosebay Willow Herb.* — *E. angustifolium.*

Leaves scattered, strap-spear-shaped, flowers unequal. — Curtis. II. 24. Flora Danica.

289.

Woods, June, P. Rare.

Near Dunstable.

287. *Codlings and Cream.* — *E. hirsutum.*

Leaves opposite, spear-shaped, serrated, running down and embracing the stem. —

Sowerby. 330. Curtis. II. 21. Flora Danica. 326.

Wet Ditches, July, P. Common.

288. *Woolly Willow Herb.* — *E. villosum.*

Leaves oblong, spear-shaped, toothed, downy, stem round, woolly. — Curtis. II. 22. Flora Danica. 347. Sowerby. 795. Wet

VIII Males. I Female. Oct. Mo.

Wet Ditches, July, P. Common.

289. *Smooth Willow Herb.* — *E. montanum.*

Leaves opposite, ovate, toothed. — Curtis.

III. 24. *Sowerby* 1177

Woods, July, P. Common.

290. *Square-stalked Willow Herb.* — *E. tetragonum.*

Leaves spear-shaped, with small teeth, stem
4-square, stigma entire. — Curtis. II. 23.

Boggy Places, July, P. Rare.

Amptbill Bogs.

291. *Marsh Willow Herb.* — *E. palustre.*

Leaves opposite, spear-shaped, very entire,
petals nicked, stem upright. — Sowerby.

346.

Bogs, July, P. Rare.

Amptbill Bogs, — Stevington Bogs.

Obs :

Very like the *tetragonum*, but differing in a round stem,
leaves narrower, almost entire.

Yellow Wort. CHLORA. Gen. Pl. 653.

Cup of 8 leaves. Blossom of 1 petal, 8 divisions,
capsule 1-cell, 2 valves, many seeds;
stigma cloven into 4.

Perfoliate

VIII Males. I Female. Oct. Mo.

292. *Perfoliate Yellow Wort.* — *C. perfoliata.*

Leaves perforated by the stem. — Sowerby.

6o.

Pastures in a limestone soil, July, A. Rare.

Milton Hill, — Bromham Pastures.

Whorts. — *VACCINIUM.* Gen. Pl. 658.

Blossom 1 petal. Threads fastened to the receptacle. Berry with 4 cells and many seeds.

293. *Black Whorts.* — *V. Myrtillus.*

Fruitstalks bearing 1 flower, leaves serrated, ovate, falling off, stem angular.—Sowerby.

456. admirable.

Woods, April, S. Rare.

Aspley Wood.

294. *Cranberry.* — *V. Oxycoccus.*

Leaves very entire, turned back, ovate, stems creeping, thread-shaped, naked.—Sowerby.

319. admirable. *Flora Danica.* 8o.

Bogs, June, S. Rare.

Ampthill Bogs, — Potton Marshes.

Obs.: — *Leaves hoary underneath. Berries easily falling off.*

Heath.

VIII Males. I Female. Oct. Mo.

Heath. ERICA. Gen. Pl. 659.

Cup of 4 leaves. Blossom 4 segments.
Threads fixed to the receptacle. Anthers
cloven in two. Capsule 4-celled.

295. *Common Heath.* — E. vulgaris.

Anthers awned, blossoms bell-shaped, nearly
equal, cups double, leaves opposite, arrow-
shaped. — Curtis. V. 30.

Heaths, July, S. Common.

296. *Cross-leaved Heath.* — E. tetralix.

Anthers awned, blossoms ovate, style within
the flower, leaves in fours, fringed, flowers
in heads. — Curtis. I. 21.

Wet Heaths, July, S. Rare.

Amphill Bogs, — Potton Marshes.

Obs :

Leaves fringed, and the fringes surmounted by a globule,
growing in ranks by fours. Flowers round, beautiful pale
pink, flowering in a half circle.

Mezereon. DAPHNE. Gen. Pl. 664.

Cup none. Blossom 4 segments, of the true
figure of a corolla, withering, inclosing the
Threads. Berry with 1 seed.

297. *Spurge Laurel.* — D. Laureola.

Bunches

VIII Males. I Female. Oct. Mo.

Bunches from the bosom of the leaves,
containing 5 flowers, leaves spear-shaped,
smooth. — Sowerby. 119.

Woods, March, S. Common.

VIII Males. III Females. Oct. Tri.

Knot Grass. POLYGONUM. Gen. Pl. 677.

Cup none. Blossom 5 segments of the na-
ture of a cup : Seed 1 angular.

298. *Bistort.* — P. *Bistorta.*

Stem unbranched, with a single spike, leaves
ovate, running down into a leafstalk. —
Curtis. I. 22. Woodville. 34. Sowerby. 509.

Wet Meadows, May, P. Rare.

Hyde Mill, Luton, — Thurlegh.

Obs :

*Root turning inwards at bottom, leaves sheathing, veiny, so
as to have the appearance of network, flowers in a close spike.*

299. *Amphibious Persicaria.* — P. *amphibium.*

Five males in the flower, styles cloven, spike
ovate. — Curtis. IV. 28. Sowerby. 436. *admirable.*

Wet Ditches, July. P. Common.

300. *Common Persicaria.* — P. *Persicaria.*

Six males in the flower, styles cloven, spikes
ovate, oblong, leaves spear-shaped, leaf-
scales, fringed. — Curtis. I. 23.

Sowerby. 750.

Cornfields

VIII Males. III Females. Oct. Tri.

Cornfields and Ditches, August, A. Common.

301. *Pale Persicaria*. — *P. lapathifolium*.

Six males in the flower, styles cloven, leaf-scales without fringes, fruitstalks rough, seeds flattened on both sides. — (*Sibthorpe.*) Curtis. I. 24. **Sowerby** 1382.

Dunghills, August, A. Common.

302. *Biting Persicaria*. — *P. Hydropiper*.

Six males in the flower, style divided half way, leaves spear-shaped, leafscales scarcely fringed. — Curtis. I. 26. Blackwell. 119. **Sowerby**. 989.

Wet Places, September, A. Common.

303. *Small Persicaria*. — *P. minus*.

Six males in the flower, style slightly divided, leaves between strap and spear-shaped, stem creeping at the base. — Curtis. I. 28. **Sowerby**. 1043.

Wet Places, September, A. Rare.

Eustow, — *Goldington*.

Obs:

Our plant is much smaller than Mr. Curtis's figure, and the stems shorter, but that Gentleman's description of the style and leaves is very accurate.

304. *Bird's Knot Grass*. — *P. aviculare*.

N

Eight

VIII Males. III Females. Oct. Tri.

Eight males and 3 females in the flowers,
from the bosom of the leaves. Leaves spear-
shaped, stem herbaceous, trailing.—Mar-
Sowerby. 1252. tyn. 91. admirable. Curtis. I. 27.

Cornfields, June. A. Common.

305. *Buckwheat.* — P. *Fagopyrum.*

Leaves heart-arrow-shaped, stem nearly up-
right, without prickles, angles of the seeds
even. — Martyn. 46. admirable. *Sowerby. 1044.*

Woods, July, A. Rare.

Wootton Wood.

306. *Climbing Buckwheat.* — P. *Convolvulus.*

Sowerby. 941. Leaves heart-shaped, stem twining, angular,
flowers blunted. — Curtis. IV. 29.

Cornfields, July, A. Common.

VIII Males. IV Females. Oct. Tetr.

Herb Paris. PARIS. Gen. Pl. 683.

Cup 4 leaves. Petals 4, narrower than the
cup. Berry with 4 cells.

307. *Common Herb Paris.* — P. *quadrifolia.*

Blackwell. 286. Sowerby. 7.

Woods, May, P. Rare.

Hawnes,—Renhold,—Clapham Park Wood.

Moschatel.

VIII Males. IV Females. Oct. Tetr.

Moschatel. ADOXA. Gen. Pl. 684.

Cup 2 segments, inferior. Blossom 4 or 5 segments, superior. Berry 4 or 5 cells, united with the cup.

308. *Tuberous Moschatel.* — A. *Moschatellina.*

Curtis. II. 26. Sowerby. 453.

Woods, April, P. Rare.

Aspley,—Renhold,—Clapham.

Obs :

Leaves rising from a root that is hollow within, shaped like those of rue and growing in a tuft. One vertical stem, bearing a roundish head of flowers, between 2 opposite leaves. Flowers with 4 divisions, pale brimstone-colored, mushy.

C L A S S. IX.

IX Males. VI Females. Enn. Hex.

Gladiole. BUTOMUS. Gen. Pl. 693.

Cup none. Petals 6. Capsule 6-celled, with many seeds.

309. *Flowering Rush.* — B. *umbellatus.*

Curtis. I. 29. admirable. Sowerby. 653.

Banks of Rivers, June, P. Rare.

Obs :

Stature 3 feet and more, 3 floral leaves like a fence, bent back; Flowers on long fruitstalks, 10 to 24 in number. Leaves rolled in at the end, 3-square.

C L A S S. X.

X Males. I Female. Dec. Mo.

Bird's Nest. MONOTROPA. Gen. Pl. 737.

Cup none. Petals 10, 5 outer ones hollowed at the base, with honey cups. Capsule 5 valves.

310. *Primrose-scented Bird's Nest.* — M. Hy-
popithys.

Sowerby. 69. Flora Danica. 232.

Beech Woods, July. P. Rare.

*Hostler's Wood, near Market-street. (Joseph
Sibley, Esq.)*

Winter Green. PYROLA. Gen. Pl. 752.

Cup 5 segments. Petals 5. Capsule open-
ing with 5 angles.

311. *Small Winter Green.* — P. minor.

Flowers in bunches, scattered, threads and
styles erect. — Sowerby. 158. Flora Danica.

55.

Woods, July, P. Rare.

Whipsnade,—Woods near Luton.

X Males. II Females. Dec. Di.

Golden Saxifrage. CHRYSOSPLENIUM. Gen.
Pl. 763.

Cup 4 or 5 segments, colored. Blossom
none. Capsule with 2 beaks, 1-cell, and
many seeds.

Common

X Males. II Females. Dec. Di.

312. Common Golden Saxifrage. — C. oppositifolium.

Leaves opposite. — Curtis, II. 27. Sowerby.

490. Flora Danica. 365.

Wet Ditches, April, P. Rare.

Eversholt.

Obs:

The terminal flower with us has always 10 males.

Saxifrage. SAXIFRAGA. Gen. Pl. 764.

Cup 5 segments. Blossom 5 petals. Capsule 2 beaks, 1-cell, many seeds.

313. White Saxifrage. — S. granulata.

Stem leaves kidney-shaped, lobed, stem branched. Root with granulations. — Curtis. I. 30. Flora Danica. 514. Woodville.

232.

Dry Pastures, May, P. Common.

314. Rue-leaved Saxifrage. — S. tridactylites.

Stem leaves wedge-shaped, 3-fingered, alternate, stem upright, branched. — Curtis, II.

28.

Sowerby. 501.

Walls, April, A. Common.

Knawel. SCLERANTHUS. Gen. Pl. 767.

Cup of 1 leaf. Blossom none. Seeds 2, enclosed in the cup.

Annual

X Males. II Females. Dec. Di.

315. Annual Knawel. — *S. annuus*.

Cups open when the seed is ripe.—Sowerby.

316.

Cornfields, June, A. Common.

Sopewort. SAPONARIA. Gen. Pl. 769.

Cup of 1 leaf, naked. Petals 5, with small claws. Capsule oblong, 1-celled.

316. Common Sopewort. — *S. officinalis*.

Cups cylindrical, leaves ovate-spear-shaped.—

Curtis. II. 29. Woodville. 251. Sowerby. 1060.

Hedges and Walls, August, P. Rare.

Bromham, — Sharnbrook.

Pink. DIANTHUS. Gen. Pl. 770.

Cup of 1 leaf cylindrical, with 4 scales at the base. Petals 5, with small claws. Capsule cylindrical, 1-celled.


317. Maiden Pink. — *D. deltoides*.

Flowers solitary, cup-scales 2, spear-shaped, blossoms crenated. — Sowerby. 61. admirable. Flora Danica. 577.

Sandy Hills, June, B. Rare,

Everton, — Pegnut Hill near Potton.

Campion.



X Males. III Females. Dec. Tri.

Campion. CUCUBALUS. Gen. Pl. 771.

Cup inflated. Petals 5, with small claws, without a crown at the mouth. Capsule 3-celled.

318. *Spatling Campion.* — C. Behen.

Cups nearly round, smooth, with network veins, capsules 3-celled, blossoms nearly naked. Sowerby. 164.

Cornfields, June, P. Common.

Catch-fly. SILENE. Gen. Pl. 772.

Cups bellying. Petals 5 with small claws, with a crown at the mouth. Capsule 3-celled.

319. *Night Catch-fly.* — S. noctiflora.

Cups with 10 angles, teeth the length of the tube, stem forked, petals nicked. — Sowerby. 291. admirable.

Cornfields, July, A. Rare.

Oakley West Field.

Obs :

The first flower is conspicuous in the fork of the plant when in seed. Cup with the appearance of network. Petals expand at six in the evening and are odoriferous.

320. *English Catch-fly.* — S. anglica.

Hairy,

X Males. III Females. Dec. Tri.

Hairy, petals nicked, flowers upright. Fruit
on reflected fruitstalks, alternate. — Curtis.

IV. 30. *Sowerby* 1170 —

Cornfields, July, A. Rare.

Woburn, — *Barton*.

Stitchwort. STELLARIA. Gen. Pl. 773.

Cup of 5 leaves, open. Petals 5, cloven in
two. Capsule 1-celled, many seeds.

321. *Great Stitchwort.* — S. Holostea.

Leaves spear-shaped, with fine serratures,
petals cloven. — Curtis. II. 30.

Woods, Hedges, May, P. Common.

322. *Small Stitchwort.* — S. graminea.

Leaves strap-shaped, very entire, flowers in
panicles. — Flora Danica. 414. *Sowerby* 803.

Meadows, July, P. Common.

323. *Middle Stitchwort.* — S. media.

Leaves strap-spear-shaped, very entire, sea-
green color, alternately opposite, petals
larger than the cup by the one half. —
(Sibthorpe.) Petiver. LVIII. 2. *Sowerby* 825.

Boggy Ditches, July, P. Rare.

Amphill Bogs, — *Ford-End Ditches*.

324. *Bog Stitchwort.* — S. uliginosa.

Leaves

Sowerby.
511.

X Males. III Females. II Dec. Tri.

Leaves spear-shaped, very entire, stem trailing. — Curtis. VI. 28. admirable. Sowerby. 1074 — Rivulets and Bogs. June, A. Rare.

Amphill Bogs, — Kempston Pits. Clapham —

Sandwort. ARENARIA. Gen. Pl. 774.

Cup of 5 leaves, open. Petals 5, entire. Capsule 1 valve, containing many seeds.

325. Plantain Sandwort. — A. trinervia.

Leaves ovate, pointed, ribbed, on leafstalks. — Curtis. IV. 31. Sowerby. 1403.

Wet Woods and Hedges, June, A. Common.

Obs: V. 11. M. X.
Stem a span high. Twin-sister of *Alsine media* as to habit, but the leaves differ in being 3-ribbed. Capsules as in that plant hanging down. Nevertheless there is no defect in any one of the characteristics of the family.

326. Thyme-leaved Sandwort. — A. serpyllifolia.

Leaves nearly ovate, pointed, sitting, blossoms shorter than the cup. — Curtis. IV. 32. Sowerby. 923.
Walls, June, A. Common.

Obs:
Many stems from 3 to 4 inches long. Branches forked, leaves under a magnifier appearing fringed. Fruitstalks with 1 flower.

O Purple

X Males. III Females. Dec. Tri.

327. *Purple Sandwort*. — *A. rubra*.

Leaves thread-shaped, leafscales skinny at the edge, and sheathing. — Petiver. LIX. 8.
Sowerby. 852.

Sandy Fields, June, A. Rare.

Potton, — *Sandy*, — *Amptbill*.

328. *Fine-leaved Sandwort*. — *A. tenuifolia*.

Leaves awl-shaped, stem panicled, capsules erect, petals spear-shaped, shorter than the cup. — Sowerby. 219.

Walls, June, A. Rare.

Newenham Wall.

X Males. V Females. Dec. Pen.

Stonecrop. *SEDUM*. Gen Pl. 789.

Cup 5 segments. Blossom 5 petals. Honey-cup scales 5 at the base of the germ. Capsules 5.

329. *Orpine*. — *S. Telephium*.

Leaves flattish, serrated, flowers in a leafy flat-topped spike, stem erect. — Curtis. III. 25. Sowerby. 1319.

Woods, July, P. Rare.

Aspley Wood.

330. *Thick-leaved Stonecrop*. — *S. dasypylum*.

Leaves

X Males. V Females. Dec. Pen.

Leaves opposite, ovate, blunt, fleshy; stem
weak, flowers scattered. — Curtis. III. 26.
Walls, June, P. Rare. **Sowerby 656.**

Market-Street.

Obs :

Leaves large in proportion, growing together, often red-
dish. Flowers whitish pink, small. Threads 12.

331. *White Stonecrop.* — *S. album.*

Leaves oblong, blunt, nearly cylindrical, sit-
ting, expanding, tuft of flowers branched.—
Curtis. I. 31. Flora Danica. 66. **Sowerby 1570.**

Walls, July, P. Rare.

Stevington, — Sharnbrook.

Obs :

Flowers white, often flesh-colored.

332. *Reflected Stonecrop.* — *S. reflexum.* **SD.** *sent*

Leaves awl-shaped, scattered, loose at the
base, lower ones bowed back. — Flora
Danica. 113. **Sowerby 695.**

Walls and Roofs, July, P. Common.

333. *Biting Stonecrop.* — *S. acre.*

Leaves nearly ovate, growing together and
sitting, bellying, nearly upright, alternate.
Tuft of flowers parting into three. — Cur-
tis. I. 32. Woodville. 231. **Sowerby 839.**

X Males. V Females. Dec. Pen.

Walls and Banks, June, P. Common.

Wood sorrel. OXALIS. Gen. Pl. 794: *

Cup of 5 leaves. Petals united by the claws.
Capsules opening by angles, 5-cornered.

334. *Common Wood sorrel.* — O. Acetosella.

Stems bearing one flower, leaves in threes,
leaflets inversely heart-shaped, hairy. — Cur-
Sowerby. 762. tis. II. 31. Woodville. 26. admirable.

Woods, April, P. Rare.

Aspley, — Ravensden, — Eversholt.

Cockle. AGROSTEMMA. Gen. Pl. 795.

Cup 1 leaf, leathery. Petals 5, with small
claws. Border blunt, entire. Capsule 1-
celled.

335. *Common Cockle.* — A. Githago.

Hairy, cups longer than the blossom, petals
entire, naked. — Curtis. III. 27. Martyn.
105. Flora Danica. 576. Sowerby. 741.

Cornfields, July, A. Common.

Lychnis. LYCHNIS. Gen. Pl. 796.

Cup of 1 leaf, oblong, smooth. Petals 5,
with small claws. Border generally cloven.
Capsule 5-celled.

336. *Ragged Lychnis.* — L. flos cuculi.

Petals

* 334. O. corniculata.

In the wood behind Oakley
House.

X Males. V Females. Dec. Pen.

Petals divided into four, fruit nearly round.

— Curtis. I. 33. Flora Danica. 590. Sowerby. 573.

Meadows, June. P. Common.

337. *Red & White Lychnis*. — L. dioica.

Male and female organs in separate plants.—

A. Curtis. II. 32. (red) B. Flora
Danica. 792. (white) Sowerby 1579.
1580.

A. Hedges, June, P. Rare.

Aspley,—Hazelwood Lane.

B. Hedges, June P. Common.

Mouse-ear. CERASTIUM. Gen. Pl. 797.

Cup of 5 leaves. Petals cloven. Capsule
1-celled, opening at the point.

338. *Common Mouse-ear*. — C. vulgatum.

Leaves ovate, petals even with the cup, stems
spreading. — Curtis. II. 34. Sowerby. 790.

Walls, Pastures, May, P. Common.

339. *Clammy Mouse-ear*. — C. viscosum.

Upright, woolly and clammy. — Curtis.
II. 35. Sowerby. 789.

Walls, Pastures, April, A. Common.

340. *Least Mouse-ear*. — C. semidecandrum.

Five males in the flower, petals nicked. —
Curtis. II. 33. Sowerby. 1630.

Walls, April, A. Common.

X Males. V Females. Dec. Pen.

341. *Field Mouse-ear.* — *C. arvense.*

Leaves between strap and spear-shaped, blunt,
smooth, blossoms larger than the cup. —
Curtis. VI. 29. Sowerby. 93.

Cornfields, May, P. Rare.

Kempston, — Ford-End, — Barton Hill.

342. *Water Mouse-ear.* — *C. aquaticum.*

Leaves heart-shaped, sitting, flowers solitary,
Sowerby. 538. fruit hanging down. — Curtis. I. 34.

Wet Ditches, July, P. Common.

343. *Dwarf Mouse-ear.* — *C. pumilum.*

Five males in the flowers, petals cloven the
length of the cup. Capsule twice as long
as the cup. — Curtis. VI. 30.

Dry Banks, March, A. Rare.

Amptbill, — Aspley.

Spurrey. SPERGULA. Gen. Pl. 798.

Cup of 5 leaves. Petals 5, entire. Capsule
ovate, 1-celled with 5 valves.

344. *Corn Spurrey.* — *S. arvensis.*

Leaves in whorls, 10 males in the flower. —
Curtis. V. 31. *Sowerby. 1536.*

Cornfields, June, A. Common.

345. *Smooth Spurrey.* — *S. pentandra.* SD.

Leaves

X Males. V Females. Dec. Pen.

Leaves in whorls, 5 males in the flower. —
Ray's Synopsis. 351. Sowerby 1536.

Sandy Fields, August, A. Rare.

Amphill.

346. *Knotted Spurrey.* — *S. nodosa.*

Leaves opposite, awl-shaped, smooth, stem
simple. — Flora Danica. 96. Curtis. IV. 34.

Bogs, July, P. Rare. Sowerby 694.

Amphill, — Stevington.

Obs :

*This plant sometimes lies trailing, and takes up more ground,
more often it is simple and erect. Bearing a white flower, large
out of proportion. The parts of fructification highly embellish
this small but elegant plant.*

C L A S S. XI.

XII. Males. I Female. Dodec. Mo.

Grasspoly. LYTHRUM. Gen. Pl. 825.

Cup with 12 segments. Petals 6, inserted in
the cup. Capsules with 2-cells and many
seeds.

347. *Purple-spiked Grasspoly.* — *L. Salicaria.*

Leaves opposite, heart-spear-shaped, flowers
in spikes, with 12 males in each. — Curtis.

III. 28. Sowerby 1061.

Banks of Rivers, July, P. Common.

Hyssop

XII Males. I Female. Dodec. Mo.

348. *Hyssop Grasspoly.* — L. *hyssopifolium.*

Leaves alternate, strap-shaped, 6 males in each flower. — Sowerby. 292. admirable.

Ditches and wet Situations, July, A. Rare.

Goldington Green, — Oakley West Field. X

Obs:

Stem 6 to 8 inches high, little branched, outermost branches at first spreading, then ascending. Flowers rather delicate, in the bosom of the leaves, pink, fugacious. Segments of the cup 12, as in the whole family, alternate ones bowed back.

XII Males. II Females. Dodec. Di.

Agrimony. AGRIMONIA. Gen. Pl. 830.

Cup with 5 teeth, surrounded by an outer cup. Petals 5. Seeds 2, in the bottom of the cup.

349. *Common Agrimony.* — A. *Eupatoria.*

Sowerby. 1335. Stem leaves winged, the odd one on a leafstalk, seeds rough with hair. — Curtis. V. 32. Woodville. 258. Martyn. 37. admirable.

Pastures and Road-sides, July, P. Common.

XII Males. III. Females. Dodec. Tri.

Dyer's Weed. RESEDA. Gen. Pl. 831.

Cup of 1 leaf, divided. Petals jagged. Capsule opening with a mouth, 1-celled.

Yellow

Do Church Field 1000.

XII Males. III Females. Dodec. Tri.

350. *Yellow Dyer's Weed.* — R. Luteola.

Leaves spear-shaped, entire, with 1 tooth on either side at the base, cups with 4 segments. — Martyn. 40. Sowerby. 320.

Walls and Way-sides, June, A. Common.

351. *Base Rocket.* — R. lutea.

All the leaves in threes, lower ones winged. — Sowerby. 321.

Cornfields, June, A. Rare.

Barton Hill.

Spurge. EUPHORBIA. Gen. Pl. 832.

Blossom 4 or 5 petals, sitting on the cup. Cup 1 leaf, bellying. Capsule 3-berried.

352. *Petty Spurge.* — E. Peplus.

Umbels with 3 divisions, forked, fencelets ovate, leaves very entire, inversely ovate, on leafstalks. — Curtis. I. 35. Sowerby. 959 — .

Gardens and Fields, July, A. Common.

353. *Dwarf Spurge.* — E. exigua.

Umbels with 3 divisions, forked, fencelets spear-shaped, leaves strap-shaped. — Curtis. IV. 36. Sowerby. 1336.

Cornfields, June, A. Common.

354. *Sun Spurge.* — E. helioscopia.

P

Umbels

XII Males. III Females. Dodec. Tri.

Umbels with 5 divisions, smaller ones with 3,
forked, fencelets inversely ovate, leaves
wedge-shaped, serrated. — Curtis. I. 36.

Sowerby. 883.

Gardens and Fields, July, A. Common.

355. *Wood Spurge*. — *E. amygdaloides*.

Umbels with many divisions, forked, fence-
lets perforated by the stem, round, leaves
blunt. — Sowerby. 256.

Woods, April, P. Common.

sent 356. *Broad-leaved Spurge*. — *E. platyphyllus*. S.D.

Umbel with 4 divisions, next 3, then forked,
Fencelets with hairy keels, leaves serrated,
spear-shaped, capsules warty.

Cornfields, July, A. Rare.

In the Corn near Bedford.

XII Males. Many Females. Dodec. Poly.

House-leek. *SEMPERVIVUM*. Gen. Pl. 837.

Cup 12 segments. Petals 12. Capsules 12,
with many seeds.

357. *Common House-leek*. — *S. tectorum*.

Leaves fringed, off-sets expanding. — Cur-
tis. III. 29. Sowerby. 1320

Walls and Houses, June, P. Common.

CLASS.

* one of Dr. Smith's desiderata.

* 356. *Cypress Spurge* *P. Cyperifolia*.

Umbel with many divisions, forked, fence-
lets nearly heart-shaped, branches barren
leaves bristly-shaped, stem-leaves spear-
shaped. Sowerby. 840. excellent.

Blackwell. Tab: 163 f. 3.

Woods & Heaths. July. P. rare

Barton —

C L A S S. XII.

XX Males. I Female. Ico. Mo.

Characters of the XIIth. Class.

- 1st. A cup of one hollow leaf.
- 2d. Petals fixed by the claws to the sides of the cup.
- 3d. Threads more than 19, fixed to the sides of the cup or blossom, and not to the receptacle.

Plumb. PRUNUS. Gen. Pl. 849.

Cup of 5 divisions, inferior. Petals 5. Nut of the fruit a stone, with prominent sutures.

358. Bullace Plumb. — P. insititia. S.D. *sent*

Fruitstalks in pairs, leaves ovate, slightly downy, rolled together, branches thorny. — Duhamel. 41. Sowerby. 841.

Hedges, April, T. Rare.

Oakley, — Renhold.

359. Red Cherry Plumb. — P. Cerasus. in fruit S.D. *sent*

Umbels nearly on fruitstalks, leaves ovate-spear-shaped, smooth, doubled together. —

Blackwell. 449. Sowerby. 706.

Woods, April, T. Common.

360. Mazzard Cherry. — P. avium. S.D. *sent*

XX Males. I Female. Ico. Mo.

Umbels sitting, leaves ovate-spear-shaped, downy underneath, doubled together, leaf-stalks with 2 glands. — Blackwell. 425.

Woods and Hedges, April, T. Rare.

Oakley Hedges, — Aspley Wood.

361. *Sloe-tree.* — *P. spinosa.*

Sowerby. 842. Fruitstalks single, leaves spear-shaped, smooth, branches thorny. — Woodville. 84.

Hedges, March, T. Common.

XX Males. II Females. Ico. Di.

Hawthorn. CRATAEGUS. Gen. Pl. 854.

Cup of 5 segments. Petals 5. Berry inferior, containing 2 seeds.

362. *Whitebeam Tree.* — *C. Aria.* S.D. *cert 1800*

Leaves ovate, cut in, serrated, downy underneath. — Flora Danica. 302.

Woods, Hedges, May, T. Rare.

Between Fenlake and Cardington.

363. *Common Hawthorn.* — *C. monogyna.*

Leaves generally divided into 3, sharp, female organ simple. — Blackwell. 149.

Woods, Hedges, May, T. Common.

Service.

*361. Wild Plum Tree. *P. domestica*

Fruitstalks nearly single, leaves spear-ovate, rolled inwards, branches unarmed.

Woodville. f. 85.

Hedges in Lane. S. April. Rare Oakley. Bromham.

Obs:

The leaves bear a distant resemblance to those of the *Salix cinerea* —

XX Males. III Females. Ico. Tri.

Service. *SORBUS.* Gen. Pl. 623.

Cup of 5 segments. Petals 5. Berry inferior, containing 3 seeds.

364. *Mountain Ash.* — *S. aucuparia.*
Leaves winged, smooth on both sides. —
Sowerby. 337.

Woods, May, T. Rare.

Aspley Wood.

Obs:

Leaflets finely serrated, 5 pair with an odd one. The odd leaf oblong-ovate. Fruit nearly round, red when ripe.

XX Males. V Females. Ico. Pen.

Pear. *PYRUS.* Gen. Pl. 858.

Cup 5 segments. Petals 5. Apple inferior, with 5 cells and many seeds.

365. *Common Pear.* — *P. communis.* S.D. sent.
Leaves serrated, smooth, flowers in flat-topped spikes. — Blackwell. 453. Sowerby. 1784.

Woods, Hedges, May, T. Rare.

Eversholt, — Thurleigh.

366. *Crab Apple.* — *P. Malus.*

Leaves serrated, umbels sitting. — Sowerby. 284. 179.

Woods, Hedges, April, T. Common.

Meadow

XX Males. V Females. Ico. Pen.

Meadow-sweet. SPIRÆA. Gen. Pl. 862.

Cup 5 segments. Petals 5. Capsule with many seeds.

367. Dropwort Meadow-sweet.—S. Filipendula.
Leaves winged, leaflets strap-spear-shaped, irregularly serrated, very smooth, flowers in tufts. — Sowerby. 284.

Dry Pastures, July, P. Common.

Obs:

The greater and smaller leaflets (being arranged along the mid-rib of the leaf in alternate order) at once point out this species.

368. Common Meadow-sweet. — S. Ulmaria.
Leaves winged, leaflets ovate, doubly serrated, hoary underneath, flowers in tufts. — Curtis. V. 33. Sowerby. 960.

Wet Meadows, July, P. Common.

XX Males. Many Females. Ico. Poly.

Rose. ROSA. Gen. Pl. 863.

Petals 5. Cup pitcher-shaped, with 5 segments, fleshy, contracted at the neck. Seeds many, rough with hair, fixed to the inner side of the cup.

369. Sweet Briar. — R. rubiginosa.
Germs round and prickly, prickles bent back
leaves rust-colored underneath. — Jacquin
Flora Aust. 50. Sowerby. 991.
Woods,

*. Cottony Rose. R. tomentosa.
Sowerby. 990 ~.

* 369. Apple Rose. R. villosa.

~~the germs round~~ germs and fruit stalks
~~prickly~~ ^{hirsute} stem with scattered prickles.
leaf stalks prickly, leaves ~~long~~ ^{cottony}.
Sowerby. Withering.
woods. Hedges

Found by R. H. W. Lady John Russell
Aspley Promham Lane, ⁱⁿ Woods

XX Males. Many Females. Ico. Poly.

Woods, Hedges, July, S. Common.

370. *Dog Rose*. — *R. canina*.

Germs ovate, fruitstalks smooth, stem and leafstalks prickly. — Curtis. V. 34. Woodville. 139. Sowerby. 992.

Hedges, July, S. Common.

371. *White Rose*. — *R. arvensis*.

Germs round, fruitstalks smooth, stem and leafstalks prickly, flowers in tufts. — Sowerby. 188.

Hedges, June, S. Common.

Obs:

Flowers in some specimens nearly approaching to umbels, fruitstalks purple.

Bramble. RUBUS. Gen. Pl. 864.

Cup 5 segments. Petals 5. Berry composed of granulations, each containing 1 seed.

372. *Raspberry Bramble*. — *R. idaeus*. SD.

Leaves winged with 5 leaflets, and sometimes with 3, stem prickly, leafstalks channelled. — Woodville. 138.

Woods, May, S. Rare.

Everton, — Aspley.

373. *Dewberry Bramble*. — *R. cæsius*.

Leaves

XX Males. Many Females. Ico. Poly.

Sowerby. 826. Leaves in threes, nearly naked, side-leaves with 2 lobes, stem round, prickly.

Woods, June, S. Common.

Obs:

Leaves mostly in three, not downy underneath.

374. *Blackberry Bramble.* — *R. fruticosus.*

Leaves 5 and 3-fingered, stem and leafstalks prickly. — Blackwell. 45. **Sowerby**. 715.

Woods, Hedges, June, P. Common.

Obs:

Leaves commonly 5, with very thick down underneath.

Strawberry. FRAGARIA. Gen. Pl. 865.

Cup 10 segments. Petals 5. Receptacle of the seeds ovate, like a berry, falling off.

375. *Common Strawberry.* — *F. vesca.*

Runners creeping. — Blackwell. 77. **Sowerby**. 1524.

Woods, May, P. Common.

376. *Barren Strawberry.* — *F. sterilis.*

Stem trailing, flowering branches loose. —

Sowerby. 1785. Curtis. III. 30. Hoffman Ger. Flo. 6.

Woods, Pastures, April, P. Common.

Cinquefoil. POTENTILLA. Gen. Pl. 866.

Cup 8 or 10 segments. Petals 4 or 5. Seeds nearly round, naked, fixed to a small dry receptacle.

374 *

R. coryli folius.
Sowerby. 827.

Aspley & Clapham Park Woods.

XX Males. Many Females. Ico. Poly.

377. *Silver Weed Cinquefoil.* — P. Anserina.

Leaves winged, serrated, stem creeping, fruit-stalks with 1 flower. — Curtis. III. 31. **Sowerby. 861.**
Road-sides, July, P. Common.

378. *Tormentil Cinquefoil.* — P. argentea.

Leaves wedge-shaped, in 5 divisions, cut in, downy underneath, stem upright. — Sowerby. 89.

Hilly Pastures, July, P. Rare.
Ampthill, — Aspley, — Rowntree Warren.

Obs:

Leaves 2 or 3 pair, towards the root more deeply jagged, underneath elegantly silvered, far from being unseemly in decay.

379. *Creeping Cinquefoil.* — P. reptans.

Leaves in fives, stem creeping, fruitstalks with 1 flower. — Curtis. I. 37. **Sowerby. 862.**

Road-sides, June, P. Common.

Obs:

A central nick in each petal. We have seen as many leaflets as 7 or even 9 in some specimens.

380. *Trailing Cinquefoil.* — P. procumbens.

Leaves in fives, stem trailing, lowest flowers with 5, uppermost with 4 petals. — (Sibthorpe.) Petiver. XLI. 10. **Sowerby. 864.**

Q

Woods,

XX Males. Many Females. Ico. Poly.

Woods, Heaths, June, P. Rare.

Ampthill, — *Aspley*.

381. *Septfoil*. — P. *Tomentilla*.

Leaves in threes, cup with 8 segments, blossoms with 4 petals. — Woodville. 27. Curtis. V. 35. *Sowerby*. 863.

Heaths, Woods, June, P. Common.

Bennet. *GEUM*. Gen. Pl. 867.

Cup 10 segments. Petals 5. Seeds with a jointed awn.

382. *Herb Bennet*. — *G. urbanum*.

Flowers upright, fruit globular, woolly, awns hooked, bare, leaves lyre-shaped. — Woodville. 259. Curtis. II. 36. *Sowerby*. 1400.

Woods, Hedges, June, P. Common.

383. *Water Avens*. — *G. rivale*.

Flowers nodding, fruit oblong, awns feathery, twisted. — Sowerby. 106.

Moist Woods, May, P. Rare.

Putnoe Wood.

Obs :

A most elegant plant, leaves winged, greater and smaller leaflets in the root-leaves arranged alternately. Petals flesh-colored, with a notch in the middle, marked with red veins. Cup mahogany-colored.

Marshlocks.

XX Males. Many Females. Ico. Poly.

Marshblocks. COMARUM. Gen. Pl. 383

Cup 10 segments. Petals 5, less in dimensions than the segments of the cup. Receptacle of the seed ovate, spongy, permanent.

384. *Purple Marshblocks.* — C. palustre.

Sowerby. 172. Hoffman. Ger. Flo. 7.

Marshes, June, P. Rare.

Obs:

Leaves winged, leaflets 2 or 3 pair, spear-shaped, serrated, with an odd one, leafscales very large, spear-shaped. Cups red. Petals less, of the same color. Seeds with granulations.

C L A S S. XIII.

Many Males. 1 Female. Poly. Mo.

Character of the XIIIth. Class.

Threads fixed to the receptacle.

Celandine. CHELIDONIUM. Gen. Pl. 880.

Blossom of 4 petals. Cup of 2 leaves. Pod 1-celled, strap-shaped.

385. *Common Celandine.* — C. majus.

Fruitstalks in umbels. — Woodville. 263.

Rubbish, June, P. Common. Sowerby. 1581.

Poppy. PAPAVER. Gen. Pl. 881.

Blossom of 4 petals. Cup of 2 leaves, capsule 1-celled, opening by pores under a permanent stigma.

Bryum
apocaulon

116

Many Males. 1 Female. Poly. Mo.

386. *Round prickly-headed Poppy.* — *P. hybridum.*

Capsules nearly round, with prominences, rough with hair, stem leafy, many-flowered. —

Sowerby. 43.

Cornfields, June, A. Common.

387. *Long prickly-headed Poppy.* — *P. Argemone.*

Capsules club-shaped, rough with hair, stem leafy, many-flowered. — Curtis. V. 38.

Sowerby. 643.

Cornfields, June, A. Common.

388. *Corn Rose.* — *P. Rhæas.*

Capsules smooth, round, stem hairy, many-flowered, leaves winged, jagged. — Curtis.

III. 32. Sowerby. 645.

Cornfields, June, A. Common.

389. *Long smooth-headed Poppy.* — *P. dubium.*

Capsules oblong, smooth, stem many-flowered, with bristles laid to, leaves winged, jagged. — Curtis. V. 37. Sowerby. 644.

Cornfields, June, A. Common.

Lime-tree. TILIA. Gen. Pl. 894.

Cup ^{NOT} segments. Petals 5. Berry dry, globular, 5-celled, 5-valved, opening at the base.

390. *Common Lime.* — *T. europaea.* Sowerby. 610.

Flowers without a honey-cup.

Woods,

Many Males. I. Female. Poly. Mo. M.

Woods, July, T. Rare.

~~Everton, — Aspley, — Potton.~~

~~old bushy shrub with pointed leaves.~~
Sunflower. CISTUS. Gen. Pl. 913.

Blossom of 5 petals. Cup 5 leaves, 2 of the leaflets smaller.

391. Sunflower Cistus. — C. Helianthemum.

Rather shrubby, trailing, leafscales spear-shaped, leaves oblong, turned back, somewhat hairy. — Curtis. V. 36. Flora Danica. 101. Sowerby 1321.

Hilly Pastures, July, S. Common.

Water Lilly. NYMPHÆA. Gen. Pl. 886.

Blossom of many petals. Cup 4 or 5 leaves. Berry many-celled, lopped.

392. Yellow Lilly. — N. lutea.

Leaves heart-shaped, entire; lobes approaching, cup of 5 leaves longer than the petals. — Sowerby. 159. admirable.

Rivers, July, P. Common.

393. White Water Lilly. — N. alba.

Leaves heart-shaped, very entire, lobes round, tiled, cup of 4 leaves. — Sowerby. 160.

Rivers, July, P. Common.

Larkspur.

Many Males. III Females. Poly. Tri.

Larkspur. DELPHINIUM. Gen. Pl. 954.

Cup none. Petals 5. Honey-cup double,
horned behind.

394. *Common Larkspur.* — D. Consolida.

Honey-cup of 1 leaf, stem subdivided. —
Flora Danica. 683.

Cornfields, June, A. Rare.

St. Leonard's Farm, near Bedford.

Many Males. V Females. Poly. Pen.

Columbine. AQUILEGIA. Gen. Pl. 954.

Cup none, petals 5, with small horns among
the petals, capsules 5, distinct.

395. *Common Columbine.* — A. vulgaris.

Honey-cups bent inwards. — Sowerby. 297.
Woods, June, P. Rare.

Barton Leat Wood.

Many Males. Many Females. Poly. Poly.

Anemone. ANEMONE. Gen. Pl. 948.

Cup none. Petals 6. Seeds many.

396. *Pasque Flower.* — A. Pulsatilla.

Fruitstalk with a fence, petals straight, leaves
with two rows of wings. — Flora Danica.

153. Relhan's Flora Cantab. at page 208.

Many Males. Many Females. Poly. Poly.

Hills, April, May, P. Rare.

Barton Hill, — Luton Downs.

Obs:

The fence which approaches the petals before the plant is in flower, is seen at the distance of 1 or 2 inches from the seeds afterwards.

397. *Wood Anemone.* — *A. nemorosa.*

Seeds pointed, leaflets cut in, stem with 1 flower. — Curtis. II. 38. Sowerby. 355.

Woods, April, P. Common.

398. *Mountain Anemone.* — *A. apennina.*

Seeds pointed, leaflets cut in, petals spear-shaped, numerous. — (Curtis.) Sowerby. 1062.

Woods, April, P. Rare.

Luton Hoe.

Obs:

This plant I have never seen in the above habitat, but merely insert it upon the credit of Mr. Knowlton in Ray's Synopsis.

Traveller's Joy. CLEMATIS. Gen. Pl. 950.

Cup none. Petals 4 to 6. Seeds with a tail.

399. *Common Traveller's Joy.* — *C. Vitalba.*

Leaves winged, leaflets heart-shaped, climbing. — Curtis. IV. 37. Sowerby. 612.

Hedges, July, S. Common.

Meadow

Many Males. Many Females. Poly. Poly.

Meadow Rue. THALICTRUM. Gen. Pl. 951.

Cup none. Petals 4 or 5. Seeds without a tail.

400. *Common Meadow Rue.* — T. flavum.

Stem leafy, grooved, panicle of many divisions, upright. — Sowerby. 367. admirable.

Wet Meadows, June, P. Common.

Crowfoot. RANUNCULUS. Gen. Pl. 953.

Cup of 5 leaves. Petals 5, with a honey-cup gland within the claws. Seeds naked.

401. *Small Spearwort.* — R. Flammula.

Leaves ovate-spear-shaped, on leafstalks, stem declining. — Curtis. VI. 37. Sowerby. 387.

Boggy Heaths, July, P. Rare.

Oakley, — Stevington, — Ampthill.

Obs. :

This sportive species varies, with leaves very entire, or deeply serrated.

402. *Great Spearwort.* — R. Lingua.

Leaves spear-shaped, stem erect. — Sowerby. 100. admirable.

Banks of Rivers, Ditches, July, P. Rare.

Geldington, — Oakley.

Pilewort

Adonis. — Adonis. Gen. Pl.

Cup of 5 leaves, Petals 5 or more without a nectary. Seeds naked.

*400. *Corn Adonis.* A. autumnalis.

Petals ~~are~~ about eight, obcordate, concave. Fruit ovate. J. Botany.

Sowerby. 308.

Corn Fields. Sep? A. Rare.

S. Leonards ~~Bakery~~, Pickering's —

Obs. :

Mr. Abbott has often brought this elegant plant in from the corn fields, and it would have been inserted in the body of the work, but from the general suspicion of its being not indigenous.

Many Males. Many Females. Poly. Poly.

403. *Pilewort*. — R. Ficaria.

Leaves heart-shaped, angular, on leafstalks,
stem with 1 flower. — Martyn. 21. **Sowerby**. 584.

Meadows, April, P. Common. observed in flower

Obs: March. 7. 1802.

The honey-cup gland being present, we must, in spite of it's
having only 3 leaves to the cup, place this plant with the rest
of the genus.

C.A.

404. *Goldilocks Crowfoot*. — R. auricomus.

Root-leaves kidney-shaped, crenated, cut in,
stem-leaves fingered, strap-shaped, stem
with many flowers. — Curtis. II. 41. **Sowerby**. 624.

Woods, Thickets, April, P. Common.

Obs:

In the whole system there is no plant more variable in it's
leaves than this. It affords specimens of the palmated, kidney-
shaped, fingered, spear and strap-shaped leaves. Some of the
petals are generally wanting.

405. *Celery-leaved Crowfoot*. — R. sceleratus.

Lower leaves hand-shaped, upper ones fin-
gered, fruit oblong. — Curtis. II. 42. **Sowerby**. 681.

Pools and wet Places, June, A. Common.

Obs:

Fruit ovate.

406. *Bulbous Crowfoot*. — R. bulbosus.

R

Cups

Many Males. Many Females. Poly. Poly.

Cups bent back, fruitstalks grooved, stem upright, with many flowers, leaves compound. — Curtis. I. 38. Martyn. 28. admirable. **Sowerby. 515.**

Pastures, May, P. Common.

Obs:

Root like a turnep in miniature.

407. *Pale-leaved Crowfoot.* — *R. hirsutus.*

Root fibrous, annual, stem hairy, cups rough with projecting points, and turned back. —

Sowerby. 1504. Curtis. II. 40.

Pastures, June, A. Common.

Obs:

Leaves paler than in the preceding species.

408. *Creeping Crowfoot.* — *R. repens.*

Cups open, fruitstalks grooved, suckers creeping, leaves compound. — Curtis. IV. 38. Martyn. 29. **Sowerby. 516.**

Wet Meadows, June, P. Common.

409. *Upright Crowfoot.* — *R. acris.*

Cups open, fruitstalks round, leaves with many divisions parted into three, upper ones strap-shaped. — Curtis. I. 39. Woodville. 246. Martyn. 30. **Sowerby. 652.**

Meadows, June, P. Common.

410. *Corn Crowfoot.* — *R. arvensis.*

Seeds

Many Males. Many Females. Poly. Poly.

Seeds prickly, upper leaves doubly compound, strap-shaped. — Curtis. VI. 36.

Martyn. 56. Flora Danica. 219. Sowerby. 135.

Cornfields, May, A. Common.

411. *Small-flowered Crowfoot.* — *R. parviflorus.*

Seeds prickly, leaves simple, jagged, pointed, hairy, stem spreading. — Sowerby. 120.

Road-sides, May, A. Common.

412. *Ivy-leaved Crowfoot.* — *R. hederaceus.*

Leaves nearly round, of 3 lobes, very entire, stem creeping. — Curtis. IV. 39.

Boggy Places, May, P. Rare.

Obs:

Without an useful remark, the Botanical Pupil may here be led into an error. The leaves correspond not with those of the rest of the family. The number of threads is not perfect. But the honey-cup-gland (the sure mark of the Crowfoot) determines the plant.

413. *Various-leaved Water Crowfoot.* — *R. heterophyllus.*

Leaves under the water hair-like, those above water with central fruitstalks. — Sowerby.

414.

Pools, May, P. Common.

414. *River Crowfoot.* — *R. fluviatilis.*

All the leaves hair-like. — Flora Danica. 376.

Pools and Rivers, June, P. Common.

Many Males. Many Females. Poly. Poly.

Obs :

We are obliged to relinquish the other two varieties, which the late celebrated Dr. Sibthorpe has called *aquatalis* and *circinatus*, for we have not yet determined them by cultivation.

Marsh Marigold. *CALTHA*. Gen. Pl. 957

Cup none. Petals 5. Honey-cups none.
Capsules many, with many seeds.

415. Marsh Marigold. — *C. palustris*.

Curtis. I. 40. *Sowerby*. 506.

Wet Ditches, April P. Common.

Hellebore. *HELLEBORUS*. Gen. Pl. 956.

Cup none. Petals 5 or more. Honey-cups
2-lipped, tubular. Capsules with many
seeds, rather erect.

416. Green Hellebore. — *H. viridis*.

Stem cloven. Branches leafy, with 2 flowers,
leaves fingered. — Curtis. VI. 34. Sowerby.

200. admirable. Woodville. 19.

Pastures, February, P. Rare.

Goldington.

Obs :

Leaves serrated towards the points, petals pointed, permanent.

417. Stinking Bear's-foot. — *H. fætidus*. SD-X

Stem with many flowers, leaves bird-footed. —

Woodville. 57. *Sowerby*. 613.

Thickets, March, P. Rare.

Bromham, — *Stevington*, — *Stagsden*.

C L A S S.

II. Polyg. Polygam.

Character of the New Class.

The characters of the new class are,

II. Polyg. Polygam.

Upon which, the name of the class is founded.

Woodville. P. Collection.

Illustrated Characters. Gen. & Order.

Illustrated Characters. Gen. & Order.

Illustrated Characters. — *Chrysanthemum*.

SD-X sent March 1st 1799.

C L A S S. XIV.

II Powers. Didynamia.

Character of the XIVth. Class.

Four threads in the flower, two higher than the others.

*

II Powers. Seeds naked. Did. Gymn.

Bugle. AJUGA. Gen. Pl. 959.

Upper lip the least. Threads longer than the upper lip.

418. *Common Bugle.* — A. reptans.

Smooth, suckers creeping. — Curtis. II. 43. Sowerby 489.
Woods, May, P. Common.

Germanander. TEUCRIUM. Gen. Pl. 960.

Blossom without an upper lip, divided beyond the base, straddling where the threads are.

419. *Common Germanander.* — T. Chamædrys.

Leaves wedge-ovate, jagged, crenated, on leafstalks, flowers in threes, stems trailing, somewhat hairy. — Woodville. 243. Sowerby. 680.
Road-sides and Walls, June, P. Rare.

Warden.

420. *Wood Sage.* — T. Scorodonia.

Leaves heart-shaped, serrated, on leafstalks, bunches lateral, flowering on one side, stem upright. — Curtis. V. 40. Sowerby. 1543.
Heaths,

II Powers. Seeds naked. Did. Gymn.

Heaths, Woods, July, P. Rare.

Aspley,—Ampthill.

Obs:

Stem 1 or 2 feet high. Suckers creeping, leaves bluntly heart-shaped, lower ones on longish fruitstalks, upper ones almost sitting, veiny, in some specimens like net-work. Floral leaves ovate.

421. *Ground Pine.* — T. Chamæpitys.

Leaves in threes, strap-shaped, very entire, flowers sitting, single stem spreading. —

Sowerby. 77. Ajuga Chamæpitys.

Dry Pastures, June, A. Rare.

Luton Downs. J. Sibley, Esq.

Cat-mint. NEPETA. Gen. Pl. 961.

Blossom, the middle segment of the lower lip scolloped, the lateral ones abridged by the edge of the mouth, bent back. Threads approaching.

422. *Common Cat-mint.* — N. Cataria.

Flowers in spikes, whorls on short fruitstalks, leaves on leafstalks, heart-shaped, tooth-serrated. — Sowerby. 137.

Hedges, July, P. Common.

Vervain. VERBENA. Gen. Pl. 43.

Blossom funnel-shaped, nearly equal, crooked, one tooth of the cup lopped.

Common



II Powers. Seeds naked. Did. Gymn.

Heaths, Woods, July, P. Rare.

Aspley,—Ampthill.

Obs:

Stem 1 or 2 feet high. Suckers creeping, leaves bluntly heart-shaped, lower ones on longish fruitstalks, upper ones almost sitting, veiny, in some specimens like net-work. Floral leaves ovate.

421. *Ground Pine.* — T. Chamæpitys.

Leaves in threes, strap-shaped, very entire, flowers sitting, single stem spreading. —

Sowerby. 77. Ajuga Chamæpitys.

Dry Pastures, June, A. Rare.

Luton Downs. J. Sibley, Esq.

Cat-mint. NEPETA. Gen. Pl. 961.

Blossom, the middle segment of the lower lip scolloped, the lateral ones abridged by the edge of the mouth, bent back. Threads approaching.

422. *Common Cat-mint.* — N. Cataria.

Flowers in spikes, whorls on short fruitstalks, leaves on leafstalks, heart-shaped, tooth-serrated. — Sowerby. 137.

Hedges, July, P. Common.

Vervain. VERBENA. Gen. Pl. 43.

Blossom funnel-shaped, nearly equal, crooked, one tooth of the cup lopped.

Common

II Powers. Seeds naked. Did. Gymn.

423. Common Vervain. — *V. officinalis.*

Spikes thread-shaped, panicled, leaves jagged, with many divisions, stem solitary. — Curtis. I. 41. Woodville, 218. Hoffman. Ger. Flo. 2. Sowerby. 767.

Road-sides, July, P. Common.

Mint. MENTHA. Gen. Pl. 967.

Blossom nearly equal, 4 segments, 1 segment nicked. Threads upright, distant.

424. Rough Water Mint. — *M. hirsuta.*

Flowers in heads, leaves ovate, serrated, nearly sitting, downy, threads longer than the blossom. — Ray. X. 1. Sowerby. 447.

Watery Places, August, P. Common.

M. aquatica minor of Sole tab. 10.

425. Smooth Water Mint. — *M. aquatica.* SD.

Leaves egg-shaped, serrated on leafstalks, threads longer than the blossom. — Petiver.

XXXI. 6. *M. aquatica major.* Sole. tab. 11.

Watery Places, August P. Rare.

Oakley.

426. Field Mint. — *M. arvensis.*

Flowers in whorls, leaves ovate, pointed, serrated, threads even with the blossom. — Flora Danica. 512. Sole. tab. 12.

Wet Places, July, P. Common.

Ground

427. Horse mint. *M. sylvestris.*

Spikes oblong, leaves oblong, woolly, serrated, sitting, threads longer than the blossom. — Blackwell 292.

Hedges July August. P. Rare. Silsoe.

421. * Wild Mint — *M. sativa*

Flowers in whorls, leaves ovate, sharp serrated, woolly. — Sowerby. f. 448. admirable. Watery Places - August. P. rare.

Pophill Hills — [REDACTED]

Oct 7 1899.

427. Red Mint. *M. gentilis.* Elstow.
[REDACTED] below the bridge

* 427. — *M. rotundifolia.*

~~427. Woolly or Horse mint. *M. sylvestris.*~~

Silsoe 1899 P.A.

* 428. Heath Mint. *M. syvestris*

[REDACTED] Oct.

Bedford-Watkin's Haddock

II Powers. Seeds naked. Did. Gymn.

Ground Ivy. GLECHOMA. Gen. Pl. 970.

Each pair of anthers growing together, in the shape of a cross. Cup 5 segments.

427. *Common Ground Ivy.* — G. hederacea.

Leaves kidney-shaped, crenated. — Curtis. II.

44. Martyn. 61. admirable. Sowerby. 853.

Hedges, April, P. Common.

Archangel. LAMIUM. Gen. Pl. 971.

Upper lip of the blossom entire, arched, lower lip with 2 lobes, the mouth with a marginal tooth on each side.

428. *White Archangel.* — L. album.

Leaves heart-shaped, pointed, serrated, on leafstalks, whorls with 20 flowers. — Curtis. II. 45. Martyn. 26. Sowerby. 760.

Rubbish, Gardens, April, P. Common.

429. *Red Archangel.* — L. purpureum.

Leaves heart-shaped, on bluntnish leafstalks. — Curtis. I. 42. Martyn. 25. Sowerby. 769.

Rubbish, Gardens, April, A. Common.

430. *Henbit Archangel.* — L. amplexicaule.

Floral leaves sitting, embracing the stem, blunt. — Curtis. II. 46. Sowerby. 770.

Walls, Cornfields, May, A. Common.

All-beal

Mentha villosa.

found at Silsoe by myself and at Chicksands by H, is represented in Sole at tab. 2^d.
m. villosa secunda.

Mentha sylvestris

Between Silsoe and Barton

m. villosa prima Sole. at tab. 1.

Mentha gentilis. Ilstone.

Sole. at tab. 15.

Mentha rotundifolia.

at Watkins

m. sylvestris Sole tab. 3.

Mentha sativa.

m. sativa — Sole tab. 21.

Mentha viridis

Cophill and Kaus's Close.

Sole. tab. 5.

431. Largeflowered Nettle Hemp. *G. versicolor.*

Stem bristly, swelled below the joints,
blossom three times as long as the
cup, its upper lip inflated.

Sowerby. 667. admirable.

Ditches near Woods. June to Sep. A Rare
^{Aug. 1799.} between Hawes and
White Wood. — between Chicksand and
Priory.

II Powers. Seeds naked. Did. Gymn.

All-heal. *GALEOPSIS.* Gen. Pl. 972.

Upper lip of the blossom somewhat crenated,
arched: lower lip with two teeth above.

431. *Corn All-heal.* — *G. Ladanum.*

Stems of an equal thickness between each
joint, all the whorls remote, cups not pun-
gent. — Petiver. XXXIII. 2. Sowerby. 884.

Cornfields, June, A. Common.

432. *Nettle Hemp.* — *G. Tetrahit.*

Joints of the stem thickened above, upper
whorls somewhat near together, cups some-
what pungent. — Sowerby. 207.

Cornfields, June, A. Common.

Weazole-snout. *GALEOBOLON.* (Hudson.)

Upper lip of the blossom upright, vaulted;
lower lip cloven into 3, segments unequal,
middlemost longer.

433. *Yellow Nettle.* — *G. luteum.*

Whorls with 6 flowers, fence of 4 leaves. —
Curtis. IV. 40. Sowerby. 787.

Woods, Thickets, May, P. Common.

Obs:

Leaves opposite, nettle-shaped. Flowers in each whorl 6 to
12, more crowded in the center, thinner at top and bottom.

S

Betony.

II Powers. Seeds naked. Did. Gymn.

Betony. BETONICA. Gen. Pl. 973.

Cup awned. Upper lip of the blossom ascending, flattish, tube cylindrical.

434. *Wood Betony.* — *B. officinalis.*

Spike interrupted, the intermediate segment of the lip nicked. — Curtis. III. 33. Woodville. 241. Sowerby. 1142.

Woods, July, P. Common.

Woundwort. STACHYS. Gen. Pl. 974.

Upper lip of the blossom arched, lower one bent back at the sides, intermediate segment greater and nicked. Threads after flowering bent towards the sides.

435. *Hedge Nettle.* — *S. sylvatica.*

Whorls with 6 flowers, leaves heart-shaped, on leafstalks. — Curtis. III. 34. Sowerby.

416.

Hedges, July, P. Common.

436. *Clown's All-heal.* — *S. palustris.*

Whorls about 6 flowers in each, leaves strap-spear-shaped, half embracing the stem and sitting. — Curtis. III. 35.

Ditches, July, P. Common.

Henbit. BALLOTA. Gen. Pl. 975.

Cup

Sowerby. 1675.

✗ *S. germanica.*

about Laton. Clapham Hill.
Sowerby. 829.

✗ *S. arvensis.*

Near Dunstable.

II Powers. Seeds naked. Did. Gymn.

Cup salver-shaped, with 5 teeth and 10 scores.

Upper lip of the blossom scolloped, concave.

437. *Stinking Henbit.* — *B. nigra.*

Leaves heart-shaped, entire, serrated, cups pointed. — Sowerby. 46. Blackwell. 136.

Hedges, July, P. Common.

Horehound. *MARRUBIUM.* Gen. Pl. 976.

Cup salver-shaped, stiff, with 10 scores.

Upper lip of the blossom with 2 segments, strap-shaped, straight.

438. *Common Horehound.* — *M. vulgare.*

Teeth of the cup bristle-shaped, hooked. — Sowerby. 410. Woodville. 97.

Road-sides, July, P. Rare.

Elstow, — *Cople*, — *Everton*.

Lion's-tail. *LEONURUS.* Gen. Pl.

Anthers sprinkled with shining dots.

439. *Mother-wort.* *L. Cardiaca.*

Stem leaves with 3 lobes, spear-shaped. — Sowerby. 286.

Road-sides, June, P. Rare.

Ford-End.

II Powers. Seeds naked. Did. Gymn.

Basil. CLINOPodium. Gen. Pl. 980.

Fence of many bristles, placed under the whorl.

440. *Wild Basil.* — *C. vulgare.*

Sowerby. 1401. Heads roundish, rough with hair, floral leaves bristle-shaped. — Flora Danica. 930.

Hedges, July, P. Common.

Marjoram. ORIGANUM. Gen. Pl. 981.

Cup a spiked 4-cornered cone.

441. *Common Marjoram.* — *O. vulgare.*

Spikes roundish, panicled, clustered, floral leaves ovate, longer than the cup. — Curtis. V. 39. Woodville. 164. Sowerby. 143.

Chalky Hills, July, P. Rare.

Streatley, — *Luton Downs.*

Thyme. THYMUS. Gen. Pl. 982.

Cups with 2 lips, mouth closed with soft hairs.

442. *Mother of Thyme.* — *T. Serpyllum.*

Flowers in heads, stems creeping, leaves blunt, flat, fringed at the base. — Curtis.

II. 47. Woodville. 110. Sowerby. 1514.

Banks, July, B. Common.

443. *Basil Thyme.* — *T. Acinos.*

H Powers. Seeds naked. Did. Gymn.

Flowers in whorls, fruitstalks wih 1 flower,
stems upright, somewhat branched, leaves
pointed, serrated. — Curtis. I. 43. Sow-
erby. 411.

Cornfields, July, A. Rare.

Barton Hill,—Aspley.

Obs:

A white semilunar spot in the center of the lower lip at once determines this species.

Calamint. MELISSA. Gen. Pl. 983.

Cup dry, somewhat flat above, upper lip nearly flat-topped; upper lip of the blossom arched, middle lobe of the lower heart-shaped.

444. *Field Calamint.* — M. Calamintha.

Fruitstalks from the bosom of the leaves, the length of the leaves. — Blackwell. 166. Sowerby. 167b.
Road-sides, August, A. Common.

Self-heal. PRUNELLA. Gen. Pl. 990.

Threads with 2 forks, one point supporting the anthers, stigma cloven in two.

445. *Common Self-heal.* — P. vulgaris.

All the leaves ovate-oblong, serrated, on leaf-
stalks. — Curtis. IV. 42. Sowerby. 961.
Pastures, June, P. Common.

Skull-cap.

II Powers. Seeds naked. Did. Gymn.

Skull-cap. SCUTELLARIA. Gen. Pl 989.

Cup, mouth entire, after flowering closed,
with a lid.

446. Common *Skull-cap.* — *S. galericulata.*

Sowerby 523. Leaves heart-spear-shaped, scolloped, flow-
ers from the bosoms of the leaves. — Cur-
tis. III. 36.

Ditches, July, P. Common.

* *

II Powers. Seeds covered. Did. Ang.

Rattle. RHINANTHUS. Gen. Pl. 997.

Cup of 4 segments, bellying. Capsule 2
cells, blunt, compressed.

447. *Yellow Rattle.* — *R. Crista galli.*

Upper lip of the blossoms flat and shorter.—
Curtis. V. 43. Martyn. 138. *Sowerby* 657.

Meadows and Pastures, June, A. Common.

Eyebright. EUPHRASIA. Gen. Pl. 998.

Cup of 4 segments, cylindrical, capsule 2
cells, ovate-oblong. Anthers below, with
1 point, thorny at the base.

448. Common *Eyebright.* — *E. officinalis.*

Leaves ovate, streaked and sharply toothed.—

Sowerby 1416. Curtis. V. 42. Woodville. 220.

Pastures, July, A. Common.

II Powers. Seeds covered. Did. Ang.

449. *Corn Eyebright.* — E. Odontites.

Leaves strap-shaped, all serrated. — Curtis.

I. 44. Martyn. 42. Sowerby. 1415.

Pastures and Cornfields, July, A. Common.

Cow-wheat. MELAMPYRUM. Gen. Pl. 999.

Cup of 4 segments, upper lip of the blossom flat, with a margin folded back. Capsule 2 cells, oblique, opening on one side. Seeds 2, hunched.

450. *Crested Cow-wheat.* — M. cristatum.

Spikes 4-square, floral leaves heart-shaped, compact, finely toothed, tiled. — Sowerby.

41.

Woods, June, A. Common.

451. *Meadow Cow-wheat.* — M. pratense.

Flowers pointing one way, sidelong, union of the leaves remote, blossoms closed. — Sowerby. 113.

Woods, July, A. Rare.

Aspley Wood.

Toothwort. LATHRÆA. Gen. Pl. 1000.

Cup with 4 segments. A depressed gland at the base of the seam of the germ. Capsule 1-cell.

Great

II Powers. Seeds covered. Did. Ang.

452. *Great Toothwort.* — *L. squamaria.*

Stem very simple, blossoms hanging down,
lower lips cloven into three. — Sowerby. 50.

Hedges and Road-sides, April, P. Rare.

Whipsnade, — *Studham.*

Lousewort. *PEDICULARIS.* Gen. Pl. 1003.

Cup of 5 segments. Capsules 2-celled, pointed, oblique. Seeds coated.

453. *Marsh Lousewort.* — *P. palustris.*

Stem branched, cups crested with callous dots, blossoms oblique in the lip. — Sowerby. 399.

Marshes, June, P. Common.

464. *Common Lousewort.* — *P. sylvatica.*

Stem branched, cups oblong, angular, smooth, blossoms with a heart-shaped lip. — Sowerby. 400.

Heaths, May, A. Rare.

Amptbill Heath.

Snapdragon. *ANTIRRHINUM.* Gen. Pl. 1007.

Cup of 5 leaves, base of the blossom jutting down beneath, with a honey-like liquor. Capsule 2-celled.

Ivy-leaved

136.

for ¹⁰ 464 read 454.

II Powers. Seeds covered. Did. Ang.

455. *Ivy-leaved Snapdragon.* — A. Cymbalaria.

Leaves heart-shaped, with 5 lobes, alternate,
stems trailing. — Curtis. I. 45. *Sowerby.* 502.

Walls, June, P. Rare.

Luton Hoe. Hulcot Church Walls.

Obs:

A very elegant plant. Leaves truly like ivy. Upper lip of
the blossom cloven in two and bent back.

456. *Sharp-pointed Fluellin.* — A. Elatine.

Leaves halberd-shaped, alternate, stems tra-
iling. — Curtis, I. 46. *Sowerby.* 692.

Cornfields, July, A. Rare.

Bromham,—Clapham.

457. *Round-leaved Fluellin.* — A. spurium.

Leaves ovate, alternate, stems trailing. —
Curtis. III. 37. *Sowerby.* 691.

Cornfields, July, A. Rare.

Clapham,—Bromham,—Stevington.

Obs:

Leaves more close in their order, and larger than in the pre-
ceding species.

458. *Least Snapdragon.* — A. minus.

Leaves mostly alternate, spear-shaped, blunt,
stem very much branched, spreading. —
Curtis. V. 41.

T

Cornfields,

H Powers. Seeds covered. Did. Ang.

Cornfields, June, A. Rare.

Oakley West Field.

Obs:

Root fibrous, stem a span high.

459. *Great Snapdragon.* — *A. majus.*

Blossoms without a spur, flowers in spikes,
cups rounded. — Sowerby. 129.

Walls, July, B. Rare.

Elstow, — Bedford.

460. *Common Toadflax.* — *A. Linaria.*

Leaves spear-strap-shaped, crowded, stem
upright, spikes terminal, sitting, flowers
tiled. — Curtis. I. 47. Martyn. 93. Wood-
ville. 221. Sowerby. 659.

Road-sides, August, P. Common.

Figwort. SCROPHULARIA. Gen. Pl. 1014.

Cup 5 segments. Blossom nearly round,
reversed. Capsule 2-celled.

461. *Knobby-rooted Figwort.* — *S. nodosa.*

Sowerby. 1544 Leaves heart-shaped, with 3 nerves in them,
stem with pointed angles. — Blackwell. 87.

Woods, July, P. Common.

462. *Water Figwort.* — *S. aquatica.*

Leaves

II Powers. Seeds covered. Did. Ang.

Leaves heart-shaped, blunt, on leafstalks, running down, stems angular, edged with a membrane, bunches terminating. — Curtis. V. 44. Sowerby. 854.

Ditches, July, P. Common.

Foxglove. DIGITALIS. Gen. Pl. 1017.

Cup 5 segments. Blossom bell-shaped, 5 segments, bellying. Capsule ovate, 2-celled.

463. *Common Foxglove.* — D. purpurea.

Leaves of the cup ovate, pointed, blossoms blunt, upper lip entire. — Curtis. I. 48. Woodville. 24. Sowerby. 1297.

Hedges, Road-sides, July, B. Rare.

Aspley, — Maulden.

Obs :

Before flowering the blossom is closed, during flower tubular and labiate, the tube marked with purple spots.

Mudwort. LIMOSELLA. Gen. Pl. 1039.

Cup 5 segments. Blossom 5 segments, equal. Threads in pairs, approaching. Capsule 1 cell, 2 valves, many seeds.

464. *Plantain Mudweed.* — L. aquatica.

Leaves spear-shaped. — Sowerby. 357. admirable. Flora Danica. 69.

II Powers. Seeds covered. Did. Ang.

Wet Places, July, A. Rare.

Goldington Green.

*Obs:**Leaves like those of plantain, root creeping.*

Broomrape. OROBANCHE. Gen. Pl. 1045.

Cup 2 segments. Blossom gaping. Capsule
1 cell, 2 valves, many seeds; a gland at the
base of the germ.

465. Great Broomrape. — *O. major.*

Stem undivided, downy, threads just emerging. — Curtis. IV. 44. Sowerby. 421.

Pastures, Woods, June, P. Rare.

Oakley, — Ampthill, — Aspley.

C L A S S. XV.

Character of the XVth. Class.

*Six threads in the flower, 4 higher, 2 lower.*Tetradynamia. — *Capsule either a Pouch or a Pod.*


*

Capsule a Pouch. Siliculosa.

Whitlow Grass. DRABA. Gen. Pl. 1076.

Pouch entire, oval-oblong, valves flattish,
parallel to the partition. Style none.

Vernal



467. *

Common Nithidate Mustard.

T. canescens.

Pouches nearly round, dotted, bordered above, leaves blunt arrow-shaped half embracing the stem, hoary and toothed
Curtis. V. 45.

Fields, July. A. Rose

Between Duck Mill and the Sluice
on the sides of the Bank —

Capsule a Pouch. Siliculosa.

466. Vernal Whitlow Grass. — *D. verna.*

Stems naked, somewhat serrated. — Curtis.
I. 49. Sowerby. 586.

Walls and Fields, March, A. Common.

Shepherd's Purse. *THLASPI.* Gen. Pl. 1078.

Pouch nicked, inversely-heart-shaped, with many seeds, valves boat-shaped, keeled and bordered.

467. Common Shepherd's Purse. — *T. Bursa pastoris.*

Pouches inversely heart-shaped, root leaves winged. — Curtis. I. 40. Sowerby. 1405.

Walls and Fields, April, A. Common.

Scurvy Grass. *COCHLEARIA.* Gen. Pl. 1079.

Pouch nicked, swelling, rough, valves hunched, blunt.

468. Swine's Cresses. — *C. Coronopus.*

Leaves winged, stem depressed. — Martyn.
92. Sowerby. 1660.

Road-sides, June, A. Common.

Candy Tuft. *IBERIS.* Gen. Pl. 1080.

Blossom irregular, the two outermost petals largest. pouch with many seeds nicked.

469. Rock Cress. — *I. nudicaulis.*

Herbaceous,

Capsule a pouch. Siliculosa.

Herbaceous, leaves waved, stem naked, undivided. — Curtis. VI. 42. Sowerby. 327.
Flora Danica. 323.

Sandy Banks, May, A. Rare.

Ampthill, — *Clophill*, — *Aspley*.

* * *

Capsule a Pod. Siliquosa.

Ladies' Smock. CARDAMINE. Gen. Pl. 1088.

Pod opening with elasticity, valves rolling back. Stigma entire. Capsule somewhat gaping.

470. Common Ladies' Smock. — C. pratensis.

Leaves winged, root-leaflets nearly round, stem-leaflets spear-shaped. — Curtis. III. 40.

Woodville. 30. Martyn. 95. Sowerby. 776.

Wet Meadows, April, P. Common.

471. Hairy Ladies' Smock. — C. hirsuta.

Leaves winged, 4 males in each flower. — Curtis IV. 48. Sowerby. 492.

Wet Ditches, April, A. Rare.

Eversholt, — Aspley.

Water Cress. SISYMBRIUM. Gen. Pl. 1089.

Pod opening, valves almost straight, cup open, blossom expanding.

Common

Phasium curvirostrum
Dec 23. 1800. Petiver
Saldwell Inclosures.

143

Capsule a Pod. Siliquosa.

472. Common Water Cress. — *S. Nasturtium*.

Pods declining, leaves winged, leaflets nearly heart-shaped. — Curtis. VI. 44. **Sowerby. 855.**

Watery Places, June, B. Common.

473. Creeping Water Cress. — *S. sylvestre*.

Pods declining, oblong-ovate, leaflets spear-shaped, serrated. — Curtis. III. 41.

Meadows, June, P. Common.

474. Jagged-leaved Water Cress. — *S. amphibium*.

Pods declining, oblong-ovate, leaves winged, tooth-serrated. — Petiver. XLIX. 9.

Banks of Rivers, July, P. Common.

475. Annual Water Radish. — *S. terrestre*.

Leaves with winged divisions, pods full of seeds. — Curtis. V. 49.

Wet Places, June, A. Rare.

Elstow, — Goldington.

Obs:

Pods extremely turgid.

476. Flixweed. — *S. Sophia*.

Petals smaller than the cup, leaves winged, and doubly compound. — Martyn. 57. **Sowerby. 963.**

Rubbish, June, A. Common.

Wormseed.

Capsule a Pod. Siliquosa.

Wormseed. ERYSIMUM. Gen. Pl. 1090.

Pod like a pillar, exactly 4-sided, cup closed.

477. *Hedge Mustard.* — E. officinale.

Pods pressed to the spike, leaves notched. —

Curtis. V. 50. Woodville. 244.

Rubbish, July, A. Common.

478. *Garlick Wormseed.* — E. Alliaria.

Leaves heart-shaped. — Curtis. II. 48.

Woodville. 245. Sowerby 796.

Hedges, April, B. Common.

479. *Winter Cresses.* E. Barbarea.

Leaves lyre-shaped, outer segment roundish. —

Sowerby. 443.

Banks, May, P. Common.

Wallflower. CHEIRANTHUS. Gen. Pl. 1091.

Germ with a glandular tooth on each side,
cup closed, 2 of the leaves hunched at the
base. Seeds flat.

480. *Common Wallflower.* — C. Cheiri.

Leaves spear-shaped, pointed, smooth, very
entire, branches angular, stem shrubby. —
Petiver. XLV. 3.

Walls, May, P. Rare.

Walls of Bedford Castle.

Turkey-pod.

*Phasium
multicapsulare.*

Capsule a Pod. Siliquosa.

Turkey-pod. ARABIS. Gen. Pl. 1094.

Honey-cup-glands 4, one within each leaf of the cup, like a reflected scale.

481. Mouse-ear. — A. thaliana.

Leaves on leafstalks, spear-shaped, very entire. — Curtis. II. 49. Sowerby. 901.

Walls and Fields, April, A. Rare.

Ampibill,—Aspley.

Obs:

The bottom leaves form a rose: stem a span high and more, seldom branched, stem leaves ovate, somewhat scalloped, flowers minute, pods long, nail-shaped, and standing on fruitstalks.

Cabbage. BRASSICA. Gen. Pl. 1096.

Cup erect, approaching. Seeds round, a gland between the shorter threads and the female organ, and between the longer and the cup.

482. Turnep. — B. Rapa.

Root continuing from the stem round, flattened, fleshy. — Martyn. 49. 50. admirable.

Fields, April, B. Common.

483. Navew. — B. Napus.

Root continuing from the stem, spindle-shaped. — Martyn. 103.

U

Fields,

Capsule a Pod. Siliquosa.

Fields, April, B. Rare.

Aspley.

Mustard. SINAPIS. Gen. Pl. 1097.

Cup open, claws of the blossom upright, a gland between the shorter threads and the female organ, and between the longer and the cup.

484. *Common Mustard.* — S. arvensis.

Pods with many angles, swollen and branched out, smooth, longer than the two-edged beak. — Curtis. V. 47. Martyn. 101.

Fields, June, A. Common.

485. *White Mustard.* — S. alba.

Pods rough with hair, beak sword-shaped, oblique, and very long. — Curtis. V. 46. Martyn. 70. Sowerby. 1677.

Fields, June, A. Common.

Obs:

This plant is easily distinguished from all the other species, by its scymitar-shaped beak.

486. *Black Mustard.* — S. nigra.

Pods smooth, pressed close to the spikestalk. —

Woodville. 151. Martyn. 51. Sowerby. 969.

Cornfields, July, A. Rare.

Clapham Hill, — Cardington.

Radish.

Capsule a Pod. Siliquosa.

Threads in Cup.

Threads in Cup. The greater number of them are short, crooked, and irregular, and are interwoven with the glandular threads, which are long, straight, and parallel.

Pod round, smooth.

Cross, V. of Mustard.

Comilla.

Cross, V. of Mustard.

Character of the Spikes.

Threads in Cup.

Glands in Cup.

Cross, V. of Mustard.

Pod round, smooth.

Threads in Cup.

Glands in Cup.

Cross, V. of Mustard.

Pod round, smooth.

Capsule a Pod. Siliquosa.

Radish. RAPHANUS. Gen. Pl. 1098.

Cup closed. Pod branching out, mostly jointed, round. Honey-cup-glands 2, between the shorter threads and the female organ, as many between the longer ones and the cup.

487. *Wild Radish.* — R. Raphanistrum.

Pods round, jointed, smooth, 1-celled. —

Curtis. IV. 46. Martyn. 71. Sowerby. 056.

Cornfields, July, A. Common.

C L A S S. XVI.

Character of the XVIth. Class.

The threads are all united at the base, but separate at the top.

Threads in I Set. V Males. Monad. Pentand.

Stork's-bill. ERODIUM. (*L' Heritiere.*)

Cup of 5 leaves. Blossom of 5 petals. Nectary 5 scales, with alternate filaments and honey-cup-glands, sitting on the base of the threads. Fruit 5 seeds, each with a beak. Beaks twisted like a corkscrew, bearded within.

488. *Hemlock-leaved Stork's-bill.* — E. circutarium.

Threads in I Set. V Males. Monad. Pentand.

Fruitstalks with many flowers, leaves winged, leaflets sitting, with many divisions. — Curtis. I. 51.

Walls and Fields, May, A. Common.

489. *Burnet Stork's-bill*. — *E. pimpinellæ-folium*.

Fruitstalks with many flowers, leaves winged,
leaflets doubly winged, 2 upper petals spotted.
— (*Sibthorpe.*)

Sandy Fields, June, A. Rare.

Aspley.

Obs.:

Petals not with 2 spots in each, but the 2 upper petals marked each at the base with 1 greenish spot.

Threads in I Set. X Males. Monad. Decand.

Crane's-bill. GERANIUM. (*L' Heritier.*)

Cup of 5 leaves. Blossom regular, of 5 petals. Nectary 5 honey-bearing glands, growing to the base of the longer threads. Fruit a seed of 5 berries, beaked; beaks simple, naked, neither twisted nor bearded.

490. Meadow Crane's-bill. — *G. pratense*.

Fruitstalks with 2 flowers, leaves with nearly central footstalks, in many divisions, wrinkled, pointed, petals entire. — Curtis. IV. 49. Sowerby. 404.

Meadows,

489 X Musky Stork's Bill. *S. moschatum*.

Fruitstalks with many flowers,
leaves winged, leaflets nearly sitting,
elliptical, irregularly ~~dentate~~.

Sowerby, tab. 902.

A May. Rare
Sandy Heaths on

Amphibians

Threads in 1 Set. X Males. Monad. Decand.

Meadows, June, P. Common.

491. *Perennial Crane's-bill.* — *G. pyrenaicum.*

Leaves rounded, with 3 or nearly 5 lobes, cut in, petals cloven in two. — Curtis. III. 42.

Rubbish, June, P. Rare. Sowerby. 405.

Luton Hooe, — Ford-End.

Obs :

Root rather woody, flowering branches prostrate, adorned with numerous flowers, twice as large as those of the common Dove's-foot. Petals nicked, between rose and flesh-color. Anthers blue.

492. *Dusky Crane's-bill.* — *G. phæum.*

Fruitstalks solitary, opposite the leaves, cups somewhat awned, stem upright, petals waved. — Sowerby. 322.

Pastures and Lanes, June, P. Rare.

Eversholt.

Obs :

The threads almost incorporated at the base afford an excellent instance of the class *Monadophila*. It is distinguished, by its elegantly waved, reflected and dusky brown petals.

493. *Herb Robert.* — *G. robertianum.*

Leaves in fives and threes, cut in, cups with 10 angles. — Curtis. I. 52. Sowerby. 1406.

Walls, Hedges, June, A. Common.

Shining

Threads in I Set. X Males. Monad. Decand.

494. *Shining Crane's-bill.* — *G. lucidum.*

Fruitstalks with 2 flowers, cups pyramidal, angular, angles raised and wrinkled, leaves with 5 lobes rounded. — Flora Danica. 218.

Sowerby. 75.

Walls, Hedges. June, A. Rare.

Elstow, — Caldwell.

495. *Soft Dove's-foot Crane's-bill.* — *G. molle.*

Fruitstalks with 2 flowers, floral leaves alternate, petals cloven, cups awnless, stem nearly upright. — Curtis. II. 50. Sowerby. 778.

Cornfields, May, A. Common.

496. *Longstalked Dove's-foot Crane's-bill.* —
G. columbinum.

Fruitstalks longer than the leaf, leaves with 5 divisions and many segments, cups with 5 angles, coat of the seeds smooth. — Sowerby. 259.

Cornfields, July, A. Rare.

Cox's Pits, Biddenham.

497. *Jagged-leaved Crane's-bill* — *G. dissectum.*

Fruitstalks shorter than the leaf, leaves with 5 divisions, subdivided into three, petals nicked, coat of the seeds woolly. — Curtis. VI. 45. Sowerby. 753.

Hedges

*495. *Dove's-foot Crane's-bill* — *G. rotundifolium.*

Fruitstalks with 2 flowers, petals entire the length of the cup, stem expanding leaves kidney-shaped, notched, seeds ~~reticulated~~, dotted

Sowerby 157.

Dry gravelly Banks, rare.

Bank on the left hand of the ^{Land,} Freeman of Bedfords Common, as you enter it from Gravellays.

July. 3. 1802.

Threads in I Set. X Males. Monad. Decand.

Hedges and Cornfields, June, A. Common.

498. *Small-flowered Dove's-foot Crane's-bill.* —
G. parviflorum.

Stem somewhat downy, 5 males in the flower,
petals nicked, seedcoats smooth, covered
with flattened hairs. — (*Curtis.*)

Curtis. VI. 46. *Sowerby.* 385.

Fallow Fields, June, A. Common.

Threads in I Set. Many Males. Monad. Polyand.

Mallow. *MALVA.* Gen. Pl. 1134.

Cup double, outer one of 3 leaves, seed coats
many, 1 seed in each.

499. *Common Mallow.* — *M. sylvestris.*

Stem herbaceous, upright, leaves with 7 pointed lobes, fruit and leafstalks hairy. — *Curtis.* II. 51. *Sowerby.* 671.

Rubbish, June, P. Common.

500. *Dwarf Mallow.* — *M. rotundifolia.*

Stem prostrate, leaves round, heart-shaped,
imperfectly 5-lobed, fruitstalks when in
seed hanging down. — *Curtis.* III. 43. *Sowerby.* 1092.

Rubbish, June, P. Common.

501. *Jagged Vervain Mallow.* — *M. moschata.*

Stem

Threads in I Set. Many Males. Monad. Polyand.

Stem erect, root leaves kidney-shaped, jagged, stem leaves with 5 divisions, segments between winged and cloven, into many divisions. — Curtis. IV. 50. *Sowerby*. 754.

Woods, June, B. Rare.

Milton, — *Clapham*, — *Bletsoe*.

C L A S S. XVII.

Character of the XVIIth. Class.

Threads in II Sets. Flowers butterfly-shaped.

Threads in II Sets. VI Males. Diad. Hexand.

Fumitory. *FUMARIA*. Gen. Pl. 1154.

Cup of 2 leaves. Blossom gaping. Threads 2, skinny, each with 3 anthers.

502. Common *Fumitory*. — *F. officinalis*.

Capsules in bunches, with 1 seed in each, stem spreading. — Curtis. II. 52. Woodville. 88. Martyn. 68. *Sowerby*. 509.

Fields, May, A. Common.

Threads in II Sets. VIII Males. Diad. Octand.

Milkwort. *POLYGALA*. Gen. Pl. 1155.

Cup of 5 leaves, leaflets wing-shaped, colored, pod inversely-heart-shaped, with 2 cells.

Common

* 502. Yellow Fumitory. *F. lutea*.

Pods nearly cylindrical, shorter than their footstalk. — Stems spreading, angular: floral leaves minute: spur short, rounded. *Sowerby*. 508.

In the vicinity of gardens, not uncommon. May. P.

Threads in II Sets. VIII Males. Diad. Octand.

503. *Common Milkwort*. — *P. vulgaris*.

Flowers awned, in bunches, stems herbaceous, simple, trailing, leaves strap-spear-shaped. — Sowerby. 76.

Barren Spots, June, B. Common.

Threads in II Sets. X Males. Diad. Decand.

Broom. *SPARTIUM*. Gen. Pl. 1166.

Stigma longitudinal, woolly above. Threads adhering to the base. Cup lengthened downwards.

504. *Common Broom*. — *S. scoparium*.

Leaves in threes and solitary, branches without prickles, angular. — Curtis. V. 52. Woodville. 89. Sowerby. 1339.

Sandy Fields, May, S. Common.

Green-wood. *GENISTA*. Gen. Pl. 1167.

Cup with 2 lips, upper consisting of 2, the under of 3 leaflets. Standard oblong, bent downwards from the male and female organs.

505. *Yellow Green-wood*. — *G. tinctoria*.

Leaves spear-shaped, smooth, branches scorched, round, upright. — Sowerby. 94. 44.

Pastures, June, S. Rare.

Threads in II Sets. X Males. Diad. Decand.

Stevington, — Clapham between Woods.

Obs :

*Stems at first trailing, then ascending, leaves ovate, sitting.
Flowers in a loose spike, yellow. Keel and wings upright.*

506. *Needle Furze. — G. anglica.*

Thorns simple, flowering branches without
prickles, leaves spear-shaped. — Sowerby.

132.

Heaths, May, S. Rare.

Ampthill Heath.

Obs :

*Root woody, lying spread on the ground. Flowering branch
erect, without prickles. Pods smooth, turgid.*

Furze. ULEX. Gen. Pl. 1169.

Cup of 2 leaves. Pods scarcely longer than
the cup.

507. *Common Furze. — U. europaeus.*

Leaves woolly, pointed, thorns scattered. —
Flora Danica. 608. *Sowerby. 742.*

Heaths, April, S. Common.

Rest-harrow. ONONIS. Gen. Pl.

Cup of 5 leaves. Segments strap-shaped.
Standard scored. Pod swollen, sitting.
Threads growing together, not separated.

Threads in II Sets. X Males. Diad. Decand.

508. *Field Rest-harrow*. — *O. arvensis*.

Flowers in bunches, leaves in threes, upper ones solitary, branches without prickles, somewhat woolly. — Blackwell. 301.

Cornfields, June, P. Common.

509. *Thorny Rest-harrow*. — *O. spinosa*.

Flowers in bunches, solitary, leaves in threes or single, branches thorny. — Martyn. 129. Sowerby. 682.

Pastures, June, P. Common.

Ladies' Finger. ANTHYLLIS. Gen. Pl. 1174.

Cup bellying. Pod nearly round, covered.

510. *Common Ladies' Finger*. — *A. Vulneraria*.

Herbaceous, leaves winged, unequal, flowers in a double head. — Sowerby. 104.

Dry Pastures, June, P. Rare.

Oakley, — *Cox's Pits*, — *Barton*.

Peaseling. OROBUS. Gen. Pl. 1185.

Style strap-shaped. Cup blunt at the base, upper segments deeper and shorter.

511. *Wood Pease*. — *O. tuberosus*.

Leaves winged, spear-shaped, leafscales half-arrow-shaped, very entire, stem simple. — Curtis. I. 53. Sowerby. 1153.

Woods, April, P. Rare.

Sphagnum arboreum.

Threads in II Sets. X Males. Diad. Decand.

Aspley, — Birchmore.

Obs:

Fruitstalks with 4 flowers, nodding before they are in bloom.

Pods black when ripe.

Vetchling. LATHYRUS. Gen. Pl. 1186.

Style flat, woolly above, broader upwards,
2 upper segments of the cup shortest.

512. *Crimson Grass Vetch.* — L. Nissolia.

Fruitstalks with 1 flower, leaves simple, leaf-scales awl-shaped. — Sowerby. 112. Curtis. VI. 51.

Hedges, June, A. Rare.

Putnoe, — Bromham, — Ampthill.

513. *Narrow-leaved Wood Pea.* — L. sylvestris.

Fruitstalks with many flowers, tendrils with 2 leaves, leaflets sword-shaped, membranes between the joints. — Curtis. VI. 52.

Sowerby. 905.

Woods, June, P. Common.

514. *Broad-leaved Wood Pea.* — L. latifolius.

Sowerby. 1100.

Fruitstalks with many flowers, tendrils with 2 leaves, leaves spear-shaped, membranes between the joints. — Flora Danica. 785.

Woods, July, P. Rare.

Hawnes, — Bromham.

Meadow

Threads in II Sets. X Males. Diad. Decand.

515. *Meadow Vetchling*. — *L. pratensis*.

Fruitstalks with many flowers, tendrils with
2 leaves undivided, leaflets spear-shaped. —

Curtis. III. 44. Martyn. 52. Sowerby. 670.

Meadows, June, P. Common.

Vetch. *VICIA*. Gen. Pl. 1187.

Stigma with a transverse beard on the lower
side.

516. *Wood Vetch*. — *V. sylvatica*.

Fruitstalks with many flowers, leaflets oval,
leafscales finely toothed. — Sowerby. 79.
admirable.

Woods, June, P. Rare.

Sheerbatch Wood, — Eversholt.

Obs:

*The fringes of the leafscales set this plant apart from all
the others.*

517. *Tufted Vetch*. — *V. Cracca*.

Fruitstalks with many flowers, flowers tiled,
leaflets spear-shaped, downy, leafscales en-
tire. — Curtis. V. 54. Sowerby. 1160 —

Woods, Hedges, June, P. Common.

518. *Common Vetch*. — *V. sativa*.

Pods sitting, generally 2 together, upright,
leaves indented, leafscales with a burnt
spot. — Martyn. 116. Sowerby. 334.

Fields,

Threads in II Sets. X Males. Diad. Decand.

Fields, June, A. Common.

519. *Narrow-leaved Vetch.* — *V. angustifolia.*

Pods nearly sitting, generally 2 together, expanding, lower leaflets inversely-heart-shaped, nicked and pointed, upper ones strap-shaped. — Johnson's Gerard. 1227. 4.

Heaths, Woods, June, A. Common.

520. *Bush Vetch.* — *V. sepium.*

Pods on partial fruitstalks, mostly in fours, upright, leaflets ovate, very entire, outer ones decreasing in size to the end. — Flora Danica. 699. Sowerby. 1515.

Woods, May, P. Common.

521. *Strangle Tare.* — *V. lathyroides.*

Pods nearly sitting, solitary, upright, smooth, leaflets 6, lower ones inversely heart-shaped. — Sowerby. 30, admirable.

Grassy spots in sand, May, A. Rare.

Amphill, — *Maulden*, — *Clophill*.

Obs:

Root ornamented with garnet-colored tubercles. Leaf-scales without the burnt spot, leaflets 2 or 3 pair without an odd one or tendrils, ending in a point.

Tine Tare. ERVUM. Gen. Pl. 1188.

Cup of 5 segments, the length of the blossom.

Smooth-podded

Threads in II Sets. X Males. Diad. Decand.

522. *Smooth-podded Tare.* — *E. tetraspermum.*

Fruitstalks with mostly 2 flowers, seeds four, round. — Curtis. I. 55.

Hedges, June, A. Common.

523. *Hairy Tine Tare.* — *E. hirsutum.*

Fruitstalks with many flowers, seeds 2, round. — Curtis. I. 54. Sowerby. 970.

Hedges, June, A. Common.

Bird's-foot. ORNITHOPUS. Gen. Pl. 1199.

Pod jointed, round, arched.

524. *Least Bird's-foot.* *O. perpusillus.*

Leaves winged, pods bowed inward. — Curtis. VI. 53. Sowerby. 369. admirable.

Heaths, June, A. Rare.

Sowthill, — *Amptbill*, — *Aspley*.

Obs:

Root rough with tubercles, fibrous. Capsule jointed. Stems much branched, trailing. Leaves 8 or 10 pair, with an odd one. Flowers in heads, small, standards scored.

Horse-shoe Vetch. HIPPOCREPIS. Gen. Pl. 1200.

Pod flat, crooked, one of the seams with many nicks.

525. *Tufted Horse-shoe Vetch.* — *H. comosa.*

Pods on fruitstalks, crowded, arched, rough, waved on either edge. — Sowerby. 31.

Hilly

Threads in II Sets. X Males. Diad. Decand.

Hilly Pastures, May, P. Rare.

Barton Hill.

Obs :

Pod flattened, zigzag. Standard marked with very delicate veins. Leaves 5 or 6 pair, with an odd one, all alike, and inversely heart-shaped.

Saintfoin. HEDYSARUM. Gen. Pl. 1204.

Keel of the blossom transversely blunt. Pod with 1 seed in each joint.

526. *Common Saintfoin.* — H. Onobrychis.

Leaves winged, shells with 1 seed, prickly, wings of the blossom as long as the cup, stem lengthened out. — Sowerby. 96. Martyn. 47.

Pastures, June, P. Rare.

Barton Hills, — Ford End Pastures.

Cock's-head. ASTRAGALUS. Gen. Pl. 1208.

Pod with 2 cells hunched.

527. *Liquorice Vetch.* — A. glycyphyllos.

Stems leafy, prostrate, pods nearly 3-square, arched, leaves oval, longer than the fruit-stalk. — Sowerby. 203.

Hedges, July, P. Rare.

Bromham, — Oakley.

Obs :

Fruitstalks round, seeds kidney-shaped.

Trefoil.

Threads in II Sets. X Males. Diad. Decand.

Trefoil. *TRIFOLIUM.* Gen. Pl. 1211.

Flowers mostly in heads. Pod scarcely longer than the cup, not opening, falling off.

528. *Melilot.* — *T. Melilotus officinalis.*

Pods in bunches, naked, 2 seeds in each, wrinkled, pointed, stem upright. — Martyn. 72. *Sowerby.* 1340.

Pastures, Woods, June, A. Common.

529. *Dutch Clover.* — *T. repens.*

Heads like umbels, pods with 4 seeds, stem creeping. — Curtis. III. 46. admirable. Martyn. 34.

Meadows, Pastures, May, P. Common.

530. *Subterraneous Trefoil.* — *T. subterraneum.*

Heads woolly, with 5 flowers, central tuft bent back, stiff, enclosing the seed. — Curtis. II. 54. admirable: *Sowerby.* 1040 -- .

Heaths, June, A. Rare.

Ampthill, — *Clophill.*

Obs :

It is an error to suppose that the capsules of this Trefoil take root in their deflected position. The root divides into very small and delicate fibres. Stems prostrate, woolly. Flowers white, tubular, and for the size of the plant beautiful.

Y

Honey-suckle

Threads in II Sets. X Males. Diad. Decand.

531. *Honey-suckle Trefoil.* — *T. pratense.*

Spikes round, somewhat woolly, surrounded by leafscales, which are opposite, with membranaceous borders, blossoms consisting of 1 petal. — Martyn. 3.

Pastures, June, P. Common.

532. *Zigzag Trefoil.* — *T. medium.*

Spikes loose, blossoms nearly equal, leafscales awl-shaped, approaching, stems zigzag, branched. — Sowerby. 190. Martyn. 13. admirable.

Pastures, June, P. Rare.

Bromham, — Milton Hill.

533. *Brimstone Trefoil.* — *T. ochroleucum.* SD.

Spikes woolly, stem upright, downy, lower leaflets inversely heart-shaped. — Curtis. VI. 49. Martyn. 35. admirable.

Pastures, June, P. Rare.

Potton, — Everton, — Clapham.

534. *Hare's-foot Trefoil.* — *T. arvense.*

Spikes woolly, oval, teeth of the cup bristle-shaped, woolly, equal. — Curtis. VI. 50.

Pastures, July, A. Common.

Sowerby. 944. 535. *Rough Trefoil.* — *T. scabrum.*

Heads

sent July 1st 1800 —

Threads in II Sets. X Males. Diad. Decand.

Heads sitting, lateral, ovate, segments of the cup unequal, rigid, bowed back. — Curtis.
VI. 48. *Sowerby. 903*.

Dry Pastures, June, A. Rare.

Potton, — Sandy, — Ampthill.

536. *Knotted Trefoil.* — *T. striatum.*

Heads sitting, nearly lateral, ovate, cups scored, rounded. — Ray's Synopsis. XIII. 3.

Dry spots, June, A. Rare.

Biddenham, — Aspley,

537. *Strawberry Trefoil.* — *T. fragiferum.*

Spikes nearly round, cups inflated with 2 teeth bent back, stems creeping. — Curtis.
II. 55. *Sowerby. 1050.*

Wet Pastures, August, P. Common.

538. *Procumbent Trefoil.* — *T. procumbens.*

Spikes somewhat ovate, tiled, standards permanent, bowed down, stems spreading. — Curtis. III. 45. *Sowerby. 945.*

Cornfields, June, A. Common.

539. *Intermediate Trefoil.* — *T. dubium.*

Spikes nearly round, loosely tiled, standards permanent, bowed down, stems trailing. — Curtis. V. 53. *Sowerby. 1256.*

Pastures, June, A. Common.

Threads in II Sets. X Males. Diad. Decand.

540. *Small Trefoil.* — *T. filiforme.*

Spikes somewhat tiled, standards permanent,
bowed down: cups on partial fruitstalks,
Sowerby 1257. stems trailing. — Ray's Synopsis. XIV. 4.

Pastures, May, A. Rare.

Biddenham, — Bury Field.

Obs :

Every flower stands on a distinct thread-shaped fruitstalk.

Claver. Lotus. Gen. Pl. 1212.

Pod cylindrical, quite straight. Wings ap-
proaching upwards longitudinally. Cup
tubular.

541. *Bird's-foot Claver.* — *L. corniculatus.*

Heads depressed, stems trailing, pods cylind-
rical, expanding. — Curtis. II. 56. Mar-
tin. 53.

Dry Pastures, June, P. Common.

Medick. MEDICAGO. Gen. Pl. 1214.

Pod flat, wreathed. Keel of the blossom bent
down from the standard.

542. *Lucern Medick.* — *M. sativa.*

Flowers in bunches, pods narrow, regular,
twisted, stem upright, smooth. — Johnson's
Gerard. 1189. 2.

Pastures, July, P. Common.

Hop

'Threads in II Sets. X Males. Diad. Decand.

543. *Hop Medick.* — *M. lupulina.*

Spikes oval, pods kidney-shaped, with 1 seed
in each, stem trailing. — Curtis. II. 57. Sowerby. 974.

Dry Pastures, June, A. Common.

C L A S S. XVIII.

Character of the XVIIIth. Class.

Threads in many Sets. Polyadelphia.

Many Sets. Many Males. Polyad. Polyand.

St. John's Wort. HYPERICUM. Gen. Pl. 1224.

Cup of 5 segments. Petals 5. Threads many,
united at the base into 5 sets. A capsule.

544. *Common St. John's Wort.* — *H. perforatum.*

Three females in each flower, stem 2-edged,
leaves blunt with transparent dots.—Curtis.
I. 57. Sowerby. 295.

Hedges, August, P. Common.

545. *St. Peter's Wort.* — *H. quadrangulum.*

Three females in each flower, stem square,
herbaceous.—Curtis. IV. 52. Sowerby. 370.

Bogs and Ditches, July, P. Common.

546. *Trailing St. John's Wort.* — *H. humifusum.*

Three

Many Sets. Many Males. Polyad. Polyand.

Three females in each flower, flowers solitary, from the bosom of the leaves, stems 2-edged, prostrate, thread-shaped, leaves smooth. — Curtis. III. 50.

Heaths, June, P. Rare.

Ampthill, — *Potton*, — *Aspley*.

Obs :

Root fibrous, stems many, branches often forked, leaves red toward the root. Segments of the cup and petals alternate.

547. *Woolly St. Peter's Wort.* — *H. elodes*.

Three females in each flower, stem cylindrical, creeping, leaves roundish, woolly. — Sowerby. 109. admirable.

Bogs, July, P. Rare.

Potton Marshes.

Obs :

Spaces between the upper joints of the stem hairy, smooth below.

548. *Hairy St. John's Wort.* — *H. hirsutum*.

Three females in each flower, cups notched, with glandules, stem round, upright, leaves ^{Sowerby. 1156.} ovate, rather downy. — Curtis. III. 49.

Woods, Hedges, July, P. Common.

549. *Upright St. John's Wort.* — *H. pulchrum*.

Three

Many Sets. Many Males. Polyad. Polyand.

Three females in each flower, cups notched,
with glandules, leaves heart-shaped, smooth,
stem round. — Curtis. I. 56.

Heaths, July, P. Rare.

Ampbill, — *Aspley*, — *Potton*.

Obs:

Stem at first weak and leaning, afterwards upright, leaves undotted, shining : between them rise fruitstalks, each bearing 3 flowers.

550. *Imperforate St. John's Wort*. — *H. dubium*.

Three females in each flower, stem obsoletely
4-edged, leaves obtuse, destitute of pellucid
dots, cup leaves elliptical. — (*English Botany*.)
Sowerby. 296.

Woods, August, P. Rare.

Luton.

Found by the late T. VAUX, Esq.

Obs:

Leaves opposite, hoary underneath.

C L A S S. XIX.

Anthers united. Syngenesia.

Character of the XIXth. Class.

Flowers compound. Anthers united so as to form a cylinder, each individual seed placed on a common bed or receptacle under it's correspondent floret.

In

Equal Polygamy. Polyg. \ae q.

In this first subdivision all the florets both in the ray and disc are furnished alike with male and female organs.

Goat's-beard. TRAGOPOGON. Gen. Pl. 1229.

Receptacle naked, cup simple, feather downy.

551. *Yellow Goat's-beard*. — *T. pratense*.

Cups as long as the rays of the blossom,
leaves entire, straight. — Petiver. XV. 6.

Pastures, June, B. Common.

Ox-tongue. *PICRIS.* Gen. Pl. 1231.

Receptacle naked, cup surrounded by another, feather downy, seeds transversely furrowed.

552. *Common Ox-tongue*. — *P. echiooides*.

Sowerby. 972. Outer cup of 5 leaves, greater than the inner, awned. — Curtis. III. 51.

Woods and Road-sides, July, A. Common.

553. Hawkweed Ox-tongue. — *P. hieracioides*.

Cups loose, leaves entire, fruitstalks covered, with scales up to the cup. — Sowerby. 196.

Borders of Fields, July, B. Common.

Obs:

Stem 2 or 3 feet high, fruitstalks loosely scaled up to the cups. Root-leaves marked with very long teeth. Globe of the feather a perfect ball.

Sowthistle.

Equal Polygamy. Polyg. Eq.

Sowthistle. *SONCHUS.* Gen. Pl. 1233.

Receptacle naked, cup tiled, bellying, feather hairy.

554. *Common Sowthistle.* — *S. oleraceus.*

Fruitstalks woolly, cups smooth. — Curtis.

II. 58. *Sowerby.* 843.

Fields, July, A. Common.

555. *Field Sowthistle.* — *S. arvensis.*

Fruitstalks and cups rough with hair, leaves notched, heart-shaped at the base. — Curtis. IV. 53. *Sowerby.* 674.

Cornfields, July, P. Common.

Obs :

Stem 3 to 4 feet high and more: milky, rising from a milky root, leaves in alternate order, notched: in the upper part of the stem entire, and clothed with harmless prickles. Flowers splendid, golden yellow. Cups with hairs surmounted by glandules, after flowering turgid at the base.

Dandelion. *LEONTODON.* Gen. Pl. 1237.

Receptacle naked, cup tiled with loosish scales, feather downy, sitting.

556. *Common Dandelion.* — *L. Taraxacum.*

Lower scales of the cup bent back, leaves notched, with fine teeth, smooth. — Curtis.

I. 58. *Sowerby.* 510 —

Meadows, Pastures, April, P. Common.

Equal Polygamy. Polyg. AEq.

557. *Autumnal Dandelion.* — L. autumnale.

Stem branched, fruitstalks scaly, leaves spear-shaped, toothed, very entire, smooth. — Johnson's Gerard. 297. 3. Sowerby. 830.

Pastures, August, P. Common.

Obs:

Root bitten off.

558. *Rough Dandelion.* — L. hispidum.

The whole cup upright, leaves toothed, very entire, rough with hair, hairs forked. —

Curtis. V. 56. Sowerby. 554.

Pastures, July, P. Common.

Obs:

The flowers nod before they open.

559. *Deficient Dandelion.* — L. hirtum.

The whole cup upright, leaves toothed, hairy, hairs undivided. — Curtis. VI. 59.

Sowerby. 555. Heaths, Commons, June, P. Common.

Hawkweed. HIERACIUM. Gen. Pl. 1238.

Receptacle naked. Cup tiled, ovate. Feather simple, sitting.

560. *Mouse-ear Hawkweed.* — H. Pilosella.

Leaves very entire, ovate, cottony underneath, suckers creeping, stem with 1 flower. —

Curtis. IV. 54. Sowerby. 1093.

Heaths,

Equal Polygamy. Polyg. Δ Eq.

Heaths, Pastures, June, P. Common.

561. *Wall Hawkweed*. — *H. murorum*.

Stem branched, root leaves ovate, toothed, stem leaf smaller. — Johnson's Gerard.

304.

Walls and Woods, July, P. Rare.

Aspley, — *Everton*, — *Silsoe*.562. *Shrubby Hawkweed*. — *H. sabaudum*.

Stem upright, bearing many flowers, leaves ovate-spear-shaped, toothed, embracing the stem. — Sowerby. 349. excellent.

Woods, July, P. Rare.

Aspley.*Obs*:

Stem 3 to 4 feet high, covered very closely towards the bottom with tooth-spear-shaped leaves.

563. *Umbelliferous Hawkweed*. — *H. umbellatum*.

Leaves strap-shaped, somewhat toothed, scattered, flowers in a kind of umbel. — Curtis. VI. 58.

Heaths and Woods, July, P. Rare.

Aspley.*Hawk's-beard*. *CREPIS*. Gen. Pl. 1239.

Receptacle naked. Cup with a smaller one at the base, scales falling off. Feather downy, on a pillar.

Equal Polygamy. Polyg. AEq.

564. Smooth Hawk's-beard. — *C. tectorum*.

Sowerby. 111. Leaves spearshaped, notched, sitting, smooth, lower ones toothed. — Curtis, V. 55.
Walls and Banks, June, A. Common.

Swine's Succory. HYOSERIS. Gen. Pl. 308

Receptacle naked. Cup nearly equal. Feather hairy, and with a small cup.

565. Least Swine's Succory. — *H. minima*.

Stem branched, naked, fruitstalks thickening upwards. — Sowerby. 95.

Cornfields, May A. Rare.

Ampthill, — Maulden.

Obs :

Leaves in a rose, crowded; stems trumpet-shaped, red towards the root.

Cat's-ear. HYPOCHÆRIS. Gen. Pl. 1246.

Receptacle chaffy. Cup somewhat tiled. Feather downy.

566. Long-rooted Cat's-ear. — *H. radicata*.

Leaves notched, blunt, rough, stem branched, naked, smooth, fruitstalks scaly. — Curtis. III. 52. *Sowerby. 83*.

Pastures, July, P. Common.

Obs : Segments of the cup keeled; keel fringed with stiffish hairs.

Smooth

Smooth Polygamy. Polyg. Sm.

Smooth Cat's-ear. — *H. glabra*.

Cups oblique, tiled, stem branched, naked.

Leaves tooth-waved. — Curtis. II. 53. *Sowerby. 57*.

Heath, June, A. Rare.

Obs :

Nodding Cat's-ear. *H. nodosa*

Receptacle naked. Cup with a narrow neck, each of the inner scales channelled.

57. Common Nodding. — *L. campanula*.

Cups on the stem angular, fruitstalks branched. — Curtis. I. 59.

Cornfields, July, A. Common.

Burrs. Cat's-ear. — *G. pilosella*.

Receptacle somewhat chaffy. Cup smooth. Feather mostly with 5 teeth, imperfect.

Obs : Receptacle smooth.

Smooth Cat's-ear. — *H. glabra*, leaves naked.

Cornfields, July, A. Common. *H. glabra*.

Obs : Receptacle smooth.

Fiducia. July, A. Common.

Equal Polygamy. Polyg. AEq.

567. Smooth Cat's-ear. — *H. glabra*.

Cups oblong, tiled, stem branched, naked,
leaves tooth-waved. — Curtis. III. 53. Sowerby. 575.

Heaths, June, A. Rare.

Ampthill, — Sandy, — Potton.

Obs: Rosy-rhizome, purple at the point, flowers not bigger than a silver penny.

Nipplewort. *LAPSANA*. Gen. Pl. 1247.

Receptacle naked. Cup with a smaller one,
each of the inner scales channelled.

568. Common Nipplewort. — *L. communis*.

Cups of the fruit angular, fruitstalks thin,
much branched. — Curtis. I. 59. Sowerby. 844.

Cornfields, July, A. Common.

Succory. *CICHORIUM*. Gen. Pl. 1251.

Receptacle somewhat chaffy. Cup double.
Feather mostly with 5 teeth, imperfectly
hairy.

569. Wild Succory. — *C. Intybus*.

Flowers in pairs, sitting, leaves notched. —
Curtis. IV. 56. Woodville. 248. Martyn.

144. Sowerby. 539.

Fields, July, B. Common.

Burr.

Equal Polygamy. Polyg. Eq.

Burr. ARCTIUM. Gen. Pl. 1253.

Cup globular, with hooked scales bent in at
the point.

570. *Clot Burr.* — A. Lappa.

Leaves heart-shaped, thornless, on leafstalks. —
Curtis. IV. 55. Woodville. 15.

Road-sides, July, B. Common.

Sawwort. SERRATULA. Gen. Pl. 1264.

Cup somewhat cylindrical, tiled, thornless.

571. *Common Sawwort.* — S. tinctoria.

Leaves harp-shaped, winged, terminal wing
the largest; florets all alike. — Sowerby.

38.

Woods and Fields, July, P. Common.

Obs:

Stem 3 to 4 feet high, with prominent angles: bottom leaves
often very entire, finely serrated. Cup closely tiled, smooth, with
scales of an elegant brown color.

572. *Way Thistle.* — S. arvensis.

Leaves toothed, thorny. — Curtis. VI. 57.
(*Carduus arvensis.*) Sowerby. 975.

Cornfields, Way-sides, July, P. Common.

Thistle. CARDUUS. Gen. Pl. 1254.

Cup ovate, tiled, with thorny scales? Recep-
tacle hairy.

Spear

Equal Polygamy. Polyg. Eq.

573. *Spear Thistle*. — *C. lanceolatus*.

Leaves running down the stem, winged, rough with hair, segments straddling, cups ovate, thorny, woolly : stem hairy. — Sowerby. 107.

Fields and Way-sides, July, B. Common.

574. *Nodding Thistle*. — *C. nutans*.

Leaves half running down the stem, thorny, flowers nodding, scales of the cup open above. — Flora Danica. 675. Sowerby. 1112.

Dry Pastures, July, B. Common.

575. *Slender Thistle*. — *C. tenuiflorus*.

Leaves running down the stem, prickly at the edge, branches straight, cups in clusters, sitting, oblong-conical, scales upright, expanding, prickly. — (Curtis.) Curtis. VI. 55. Sowerby. 412.

Borders of Fields, July, A. Rare.

Alm's Houses, -- Cox's Pits.

576. *Thistle-upon-Thistle*. — *C. polyacanthos*.

Leaves running down the stem, prickly at the edge, branches expanding, cups nearly round, loose, scales awl-shaped, expanding, rather harmless. — (Curtis.) Curtis. VI. 54. Sowerby. 973.

Hedges, July, A. Common.

Marsh

Equal Polygamy. Polyg. Eq.

577. *Marsh Thistle.* — *C. palustris.*

Leaves running down the stem, toothed, prickly at the edge, flowers in bunches, upright,

Sowerby 974. fruitstalks thornless. — Curtis. VI. 56.

Wet Places, July, B. Common.

578. *Milk Thistle.* — *C. marianus.*

Leaves embracing the stem, halberd-shaped, winged. Cups leafless, thorns channelled,

Sowerby 976. with double rows of thorns. — Curtis. III. 54.

Rubbish, June, B. Common.

579. *Woolly Thistle.* — *C. eriophorus.*

Leaves sitting, with winged divisions, pointing 2 ways, segments alternate, upright, cups globular, woolly. — Sowerby. 386.

Dry Pastures, July, B. Common.

580. *Dwarf Thistle.* — *C. acaulis.*

Stemless, cup smooth. — Sowerby. 161.

Hilly Pastures, July, P. Common.

Obs:

Cup closely tiled: Root turning inwards, with a brown bark, whitish red within.

Cotton Thistle. ONOPORDUM. Gen. Pl. 1256.

Receptacle like a honey-comb. Cupscales pointed.

Common

Equal Polygamy. Polyg. Æq .

581. Common Cotton Thistle. — O. Acanthium.

Cups scurfy, scales expanding, leaves oblong
and waved. — Curtis. V. 57. Sowerby 977--.

Borders of Fields, July, B. Common.

Carline Thistle. CARLINA. Gen. Pl. 1258.

Cup with a ray, consisting of long marginal
colored scales.

582. Common Carline Thistle. — C. vulgaris.

Stem with many flowers in a flat topped spike,
flowers terminal, cups with a white ray. —

Johnson's Gerard. 1159. I. Sowerby 1144--.

Hilly dry Pastures, July, B. Common.

Hemp Agrimony. BIDENS. Gen. Pl. 1267.

Receptacle chaffy. Feather with straight
awns, rough. Cup tiled. Blossom with
sometimes one or two radiated florets.

583. Trifid Hemp Agrimony. — B. tripartita.

Leaves in threes, cups somewhat leafy, seeds
upright. — Curtis. IV. 57. Sowerby 1113.

Banks and wet Places, August, A. Common.

584. Nodding Hemp Agrimony. — B. cernua.

Leaves spear-shaped, embracing the stem,
flowers nodding, seeds upright. — Curtis.

III. 55. Sowerby 1114.

A a

Wet

Equal Polygamy. Polyg. Æq .

Wet Places, August, A. Rare.

Clapham, — *Clophill*.

585. *Least Hemp Agrimony*. — *B. minimus*. SD.

Leaves spear-shaped, sitting, flowers as well
as the seeds upright. — Ray's Synopsis.
VII. 2.

Wet Places, August, A. Rare.

Kempston Pits.

Dutch Agrimony. EUPATORIUM. Gen. Pl. 1272.

Receptacle naked. Feather downy. Cup ob-
long. Stile cloven halfway down, long.

586. *Common Dutch Agrimony*. — *E. cannabi-
num*.

Leaves fringed. — Sowerby. 428.

Woods, July, P. Common.

Superfluous Polygamy. Polyg. Super.

The hermaphrodite florets of the center are
furnished with a female organ, and produce
seeds. The female florets also in the cir-
cumference produce seeds.

Tansy. TANACETUM. Gen. Pl. 1280.

Receptacle naked. Feather with sometimes
a border. Cup tiled, hemispherical. Blos-
soms of the ray obscurely divided into three.

Common

Bryum
No. 1. sent to
D. Smith Sowerby
Jan. 26. 1801.
Jan. 22 1801.
No. 2.

Sent to
Dickson *R. purpureum*.

No. 2.
1801. Feb. 10.

179
Superfluous Polygamy. — Polyg. Super.

587. Common Tansy. — *T. vulgare*.
Leaves doubly winged, jagged, serrated. —
Woodville. 115.

Pastures and Road-sides, July, P. Rare.
Bromham Grange.

Southernwood. *ARTEMISIA*. Gen. Pl. 1281.

Receptacle somewhat woolly or nearly bare.
Feather none. Cup tiled with roundish
scales, approaching. No blossoms in the
ray.

588. Wormwood. — *A. Absinthium*.
Leaves compound, in many divisions, flowers
nearly round, hanging down, receptacle
woolly. — Woodville. 120.
Road-sides, August, P. Common.

589. Common Mugwort. — *A. vulgaris*.
Leaves winged, flat, jagged, downy under-
neath, bunches simple, bent back, florets in
the ray five. — Woodville. 121. Sowerby. 978—
Road-sides, August, P. Common.

Cudweed *GNAPHALIUM*. Gen. Pl. 1282.
Receptacle naked. Feather downy. Cup ti-
led with rounded scales at the edge, skinny,
colored.

590. Upright Cudweed. — *G. rectum*.

Superfluous Polygamy. Polyg. Super.

Stem upright, terminating in a leafy compound spike, leaves strap-spear-shaped, almost naked on the upper side.—(English Botany.) Sowerby. 124. excellent.

Heaths and Woods, August, B. Rare.

Clophill,—*Aspley*,—*Soutbill*.

Obs:

Root fibrous, leaves hoary underneath, for the most part sitting in the bosoms of the leaves.

591. *Marsh Cudweed*. — *G. uliginosum*.

Stem herbaceous, branched, spreading, flowers crowded, terminal.—Flora Danica. 859.

Inundated spots, August, A. Common.

Spikenard. *CONYZA*. Gen. Pl. 1286.

Receptacle naked. Feather simple. Cup tiled, nearly round. Blossoms of the ray trifid.

592. *Plowman's Spikenard*. — *C. squarrosa*.

Leaves spear-shaped, pointed, stem herbaceous, bearing loose topped spikes, cups scurfy. — Blackwell. 102.

Road-sides, August, B. Common.

Fleabane. *ERIGERON*. Gen. Pl. 1287.

Receptacle naked. Feather hairy. Blossoms of the ray strap-shaped, very narrow.

Purple ~

Common Fleabane. — *P.*193. *Purple Fleabane*. — *E. purpurea*.

Frustulate, slender, bearing one flower.

Stems 1 foot. Hoffman. Gen. Flora. 1287.

August, B. Rare.

194. *Common Fleabane*. — *P. adonis*.

Stems 1 foot. Hoffman. Gen. Flora.

August, B. Common.

195. *Common Fleabane*. — *E. canina*.

Herbaceous, naked. Stems branched, long.

Leaves lanceolate, linear, pointed, smooth,

rayed, strap-shaped.

196. *Common Fleabane*. — *E. canina*.

Stems branched, hairy, leaves wither

anthers, annual.

Common. Sowerby. 1287.

Clay soil, March, P. Common.

197. *Butter Bur*. — *P. officinalis*.

Stems obovate-shaped, leaves broad, pointed,

— Sowerby. 1287. Sowerby. 1287.

Banks of Mill-streams, April, P. Common.

Common. Sowerby. Gen. Pl. 1287.

Leaves broad, pointed, simple. Cup

shaped, smaller top at the base.

Scars of attachment at the point.

Superfluous Polygamy. Polyg. Super.

593. Purple Fleabane. — E. acre.

Fruitstalks alternate, bearing one flower. —

Curtis. I. 60. Hoffman. Ger. Flo. 11. Sowerby. 1158. —

Pastures, August, B. Rare. —

Thurlegh, — Ampthill, — Biddenham. —

Obs.

Our specimens are 6 inches high, leaves alternate, points of the cup purple. Outer row of florets a diluted purple.

Colt's-foot. TUSSILAGO. Gen. Pl. 1289.

Receptacle naked. Cup-scales equal, long as the center, somewhat membranaceous. Feather simple.

594. Common Colt's-foot. — T. Farfara.

Stem tiled, bearing one flower, leaves rather heart-shaped, angular, finely toothed. —

Curtis. II. 60. Sowerby. 429. —

Clay soil, March, P. Common.

595. Butter Burr. — T. Petasites.

Panicle egg-shaped, female florets few, naked. — Curtis. II. 59. Sowerby. 431. —

Banks of Mill-streams, April, P. Common.

Groundsell. SENECIO. Gen. Pl. 1290.

Receptacle naked. Feather simple. Cup cylindrical, with a smaller cup at the base. Scales as it were dead at the point. —

Purple

Common

Superfluous Polygamy. Polyg. Super.

596. Common Groundsell. — *S. vulgaris.*

— Flowers without rays, leaves winged and waved, embracing the stem, flowers scattered. —

Curtis. I. 61. Sowerby. 747.

Gardens, Fields, May, A. Common.

597. Bushy Groundsell. — *S. sylvaticus.*

Blossoms in the ray bent back, leaves winged, finely toothed, stem branching to a flat topped spike, upright. — Johnson's Gerard.

278. 2. Sowerby. 748.

Heaths, June, A. Common.

598. Hoary Groundsell. — *S. erucæfolius.*

Blossoms radiate, leaves winged, toothed, somewhat hairy, stem upright. — Curtis.

V. 64. Sowerby. 574.

Hedges, August, P. Common.

599. Common Ragwort. — *S. Jacobæa.*

Blossoms radiate, leaves winged, lyre-shaped, segments subdivided, stem upright. — Martyn. 85. excellent. Sowerby. 1130 —

Barren Spots, July, P. Common.

600. Water Ragwort. — *S. aquaticus.*

Blossoms radiate, leaves toothed, root leaves ovate, stem leaves winged, terminal segment the largest. — Flora Danica. 784.

Sowerby. 1131.

Wet

Superfluous Polygamy. Polyg. Super.

Wet Meadows, July, P. Common.

Golden-rod. SOLIDAGO. Gen. Pl. 1292.

Receptacle naked. Feather simple. Blos-
soms of the ray about 5. Scales of the cup
tiled, closed.

601. *Common Golden-rod.* — S. Virgaurea.

Stem somewhat zigzag, bunches panicled,
upright, crowded. — Sowerby. 301. excel-
lent.

Woods, Heaths, August, P. Rare.

Aspley, — Birchmore, — Potton Wood.

Found by J. PARKER, M. D.

Obs:

*Stem 2 feet or more in height, leaves slightly notched, ellipti-
cal, turned back, flowers of the whole ray 6 to 9.*

Elecampane. INULA. Gen. Pl. 1295.

Receptacle naked. Feather simple. Anthers
ending at the base in 2 bristles.

602. *Common Elecampane.* — I. Helenium.

Leaves embracing the stem, ovate, wrinkled,
cottony underneath, cup-scales ovate. —
Woodville. 108. Sowerby 1546.

Pastures, July, P. Rare.

Ravensden, — Stevington.

Middle

decipiens.

Superfluous Polygamy. Polyg. Super.

603. Middle Fleabane. — *I. dysenterica*.

Leaves embracing the stem, heart-oblong,
stem woolly, panicled, capsules bristle-
shaped. — Curtis. III. 56. Sowerby. 1115.

Road-sides, August, P. Common.

604. Least Fleabane. — *I. Pulicaria*.

Leaves embracing the stem, stem prostrate,
flowers almost globular with a very short
ray. Curtis. III. 57.

Inundated Spots, August, A. Rare.

Goldington, — Ravensden.

Obs :

Root descending obliquely. Flowers button-shaped, florets of
the ray small and as it were abridged.

Daisy. *BELLIS*. Gen. Pl. 1300.

Receptacle naked. Feather none. Cup hemispherical with equal scales. Seeds inversely ovate.

605. Common Daisy. — *B. perennis*.

Stalk naked. — Sowerby. 424. Curtis. I. 62.

Pastures, May, P. Common.

Marigold. *CHRYSANTHEMUM*. Gen. Pl. 1307.

Receptacle naked. Feather bordered. Cups hemispherical, tiled with marginal membranaceous scales.

Ox-eye

Superfluous Polygamy. Polyg. Super.

606. *Ox-eye Daisy*. — *C. leucanthemum*.

Leaves embracing the stem, oblong, serrated above, toothed below. — Curtis. V. 62.
Martyn. 109. **Sowerby. 609.**

Pastures, July, P. Common.

607. *Chamomile Ox-eye*. — *C. inodorum*.

Leaves winged in many divisions, stem branched, spreading. — Flora Danica. 696. **Sowerby. 676.**
Cornfields and Rubbish, July, A. Common.

608. *Corn Marigold*. — *C. segetum*.

Leaves embracing the stem, jagged above, tooth-serrated below. — Curtis. VI. 60.
• Martyn. 110. **Sowerby. 540.**

Cornfields, June, A. Common.

Feverfew. **MATRICARIA**. Gen. Pl. 1308.

Receptacle naked. Feather none. Cup hemispherical, tiled. Marginal scales rather pointed, solid.

609. *Common Feverfew*. — *M. parthenium*.

Leaves compound, flat, leaflets ovate, jagged, fruitstalks branched. — Woodville. 249.

Road-sides, June, P. Common.

610. *Corn Feverfew*. — *M. Chamomilla*.

B b Receptacles

Superfluous Polygamy. Polyg. Super.

Receptacles conical, rays expanding, cup-scales even at the edge. — Curtis. V. 63.
Martyn. 74.

Cornfields, June, A. Common.

Chamomile. ANTHEMIS. Gen. Pl. 1312.

Receptacle chaffy. Feather none. Cup hemispherical, nearly equal. Florets of the ray more than five.

611. *Fætid Chamomile.* — A. Cotula.

Receptacles conical, chaff bristle-shaped, seeds without a border. — Curtis. V. 61.

Cornfields, May, A. Common.

612. *Corn Chamomile.* — A. arvensis.

Receptacles conical, chaff bristle-shaped, seeds crowned with a border. — Petiver.

XIX. 8. Sowerby. 602.

Cornfields, July, A. Rare.

Oakley West Field.

Obs :

Chaff broader than in the foregoing species.

Milfoil. ACHILLEA. Gen. Pl. 1313.

Receptacle chaffy. Feather none. Cup ovate, tiled. Florets of the ray about four.

613. *Common Milfoil.* — A. Millefolium.

Leaves

Superfluous Polygamy. Polyg. Super.

Leaves doubly winged, naked, segments strap-shaped, toothed, stem leaves furrowed upwards. — Curtis. VI. 61. Woodville. 64. Martyn. 123. excellent. Sowerby. 758.

Pastures, July, P. Common.

614. Sneezewort. — A. Ptarmica.

Leaves spear-shaped, pointed, finely serrated. — Curtis. V. 60. Sowerby. 757.

Banks of Rivers, July, P. Common.

* * *

Frustraneous Polygamy. Polyg. Frustra.

The hermaphrodite florets of the center are furnished with a stigma, and so bear seed: but the florets, which form the circumference, have no stigma and are unable to bear seed.

Centaury. CENTAUREA. Gen. Pl. 1331.

Receptacle bristly. Feather simple. Blossoms of the ray funnel-shaped, longer and irregular.

615. Blue Bottle. — C. Cyanus.

Cups serrated, leaves strap-shaped, very entire, lower ones toothed — Curtis. VI. 62. Sowerby. 277.

Cornfields, July, A. Common.

616. Least Knapweed. — C. nigra.

Frustraneous Polygamy. Polyg. Frustra.

Cups fringed, with a small ovate scale at the end : fringes hair-like, upright, leaves lyre-shaped, angular, flowers without rays. — Sowerby. 278. Martyn. 130.

Pastures, July, P. Common.

617. Great Knapweed. — C. Scabiosa.

Cups fringed, leaves winged, wings spear-shaped. — Sowerby. 56.

Cornfields, July, P. Common.

618. Star Thistle. — C. Calcitrapa.

Cups almost doubly thorned, sitting, leaves winged, strap-shaped, toothed, stem hairy. — Sowerby. 125. excellent.

Road-sides, July, A. Rare.

Biddenham,—Eaton-Soccon.

* * * *

Necessary Polygamy. Polyg. Necess.

The hermaphrodite florets, from some defect in the stigmas, are unable to produce seeds, but the female florets in the ray perfect their seeds.

Cat's-foot. FILAGO. Gen. Pl. 1345.

Receptacle naked. Feather none. Cup tiled : female florets placed between the scales of the cup.

Common

Necessary Polygamy. Polyg. Necess.

619. Common Cat's foot. — F. germanica.

Stems several, leaves scattered, from the bases of the leaves, the flowers perched on long slender pedicels.

Road-sides, July, Common.

620. Mountain Cat's foot. — F. montana.

Stems several, naked, pedicels slender and mostly erect, from the axils of the leaves.

Road-sides, July, Common.

Common Cat's foot.

I House. I. Male. Monoec. Monand.

Wet Ditches, June, A. Common.

646. *Brittle Stone-wort*. — *C. tomentosa*.

Prickles on the stem ovate. — Johnson's Gerard. 1115.

Ditches, Bogs, June, A. Rare.

Clapham Springs, — *Stevington Bogs*.

Obs:

Plant more compact in its habit than the former, when living, flesh-colored; dead, ashy white.

I House. II Males. Monoec. Diand.

Duckweed. *LEMNA*. Gen. Pl. 1400.

Male. — Cup of 1 leaf. Blossom none.

Female. — Cup of 1 leaf. Blossom none.

Style 1. Capsule 1 cell.

647. *Ivy-leaved Duckweed*. — *L. trisulca*.

Leaves on leafstalks, spear-shaped. *Sowerby*. 926.

Ditches, August, A. Common.

648. *Least Duckweed*. — *L. minor*.

Leaves sitting, flattish on both sides, root single. — Ray's Synopsis. IV. 1. *Sowerby* 1095.

Ditches, June, A. Common.

Obs:

The single root and flat leaves at once demonstrate this species.

Hunched

I House. II Males.

646. one of Dr. Smith's desiderata.

647* *Chara hispida* —

Bedford River

Sep. 10. 1800.

I House. III Males. Monoec. Trand.

Rock Plant. — *Chara*. Gen. Pl. 1400.

Male. — Ovate cylindrical, composed entirely of 3-leaved, blossoms none.

Female. — Ovate cylindrical, below the male flowers. Cup-like & hollowed out. Blossoms none. Seed of nothing but a hair-like down.

648. *Broad-leaved Rock Plant*. — *C. latifolia*.

Leaves somewhat sword-shaped, narrow and pointed, closely close together. — Curtis. III. 61. *Sowerby*. 1455.

Sergeant Water Park. P. Common.

649. *Smooth Rock Plant*. — *C. glabra*.

Leaves smooth, narrow,

10. June 1800.

Bedford River. — *Chara*.

I House. II Males. Monoec. Diand.

649. *Hunched Duckweed*. — *L. gibba*.

Leaves sitting, hemispherical underneath,
roots single.

Ditches, July, P. Rare.

Goldington.

650. *Great Duckweed*. — *L. polyrrhiza*.

Leaves sitting, roots crowded. — Ray's Sy-
nopsis. IV. 2.

Obs:

Leaves red underneath.

I House. III Males. Monoec. Triand.

Reed Mace. *Typha*. Gen. Pl. 1401.

Male. — Catkin cylindrical. Cup indistinct-
ly 3-leaved. Blossom none.

Female. — Catkin cylindrical, below the male
flowers. Cup with a hair-like wool.
Blossom none. Seed 1, sitting on a hair-
like down.

651. *Broad-leaved Reed Mace*. — *T. latifolia*.

Leaves somewhat sword-shaped, male and
female spikes close together. — Curtis. III.

61. *Sowerby*. 1455.

Stagnant Water, June, P. Common.

652. *Small Reed Mace*. — *T. angustifolia*.

Leaves

10th Line add

Watery Pits. P. June. Rare
Ford-End

I House. III Males. Monoec. Triand.

Leaves half-cylindrical, male and female spikes distinct from each other. — Curtis.
III. 62. Sowerby. 1456:

Pools, July, P. Rare.

Knotting-Green.

Bur-reed. SPARGANIUM. Gen. Pl. 1402.

Male. — Catkin nearly round. Cup of 3 leaves. Blossom none.

Female. — Catkin nearly round. Cup of 3 leaves. Blossom none. Stigma divided. Fruit juiceless, with 1 seed.

653. *Branched Bur-reed.* — S. erectum.

Sowerby. 744. Leaves upright, 3-square. — Curtis. V. 68.

Wet Ditches, July, P. Common.

654. *Simple Bur-reed.* — S. simplex.

Leaves 3-square at the base, the sides flat,

Sowerby. 745. fruitstalks simple. — Curtis. V. 67.

Ditches, July, P. Rare.

Elstow.

Obs :

Upper flowers sitting, lower ones on fruitstalks.

Sedge. CAREX. Gen. Pl. 1407.

Male. — Catkin tiled. Cup of 1 leaf. Blossom none.

Female. —

I House. III Males. Monoec. Triand.

Female. — Catkin tiled. Cup of 1 leaf.
Blossom none. Nectary inflated, with
3 teeth. Stigmas three. Seeds 3-square,
beneath the nectary.

655. *Small Sedge*. — *C. dioica*.

Spike simple, male and female catkins on
distinct plants. — Flora Danica. 369. Sowerby. 543.

Turfy Bogs, June, P. Rare.

Amptbill.

656. *Naked Sedge*. — *C. leporina*.

Spike compound, spikelets ovate, sitting, ap-
proaching, alternate, naked, composed of
barren and fertile flowers. — Sowerby. 306. *C. ovalis* —
Bogs, June, P. Rare.

Potton Marshes.

657. *Great Sedge*. — *C. vulpina*.

Spike more than doubly compound, less com-
pact in the lower part, spikelets ovate,
clustered, composed of fertile and barren
flowers, the barren flowers above. — Sower-
by. 307.

Wet Places, June, P. Common.

Obs:

Spikelets (when ripe) brown and permanent.

658. *Spiked Sedge*. — *C. spicata*. Sowerby. 1097 —

D d

Spikelets

I House. III Males. Monoec. Triand.

Spikelets nearly round, containing both fertile and barren florets, contiguous; capsules ovate, pointed.

Ditch Banks, June, P. Common.

Obs:

Very like the preceding, but smaller.

659. *Soft Sedge.* — *C. disticha.*

Spike compound, nearly 2-rowed, spikelets egg-shaped, somewhat tilted, with barren and fertile flowers; floral leaves longer than the spikelets, straw 3-square. — (*Hudson.*)

Leers. XIV. 2.

Ditches and Bogs, May, P. Rare.

Eversholt.

660. *Prickly Sedge.* — *C. muricata.* SD. sent-

Spikelets nearly ovate, sitting, remote, both fertile and barren, capsules pointed, diverging, thorny. — Leers. XIV. 8. Flora Danica. 284. *Sowerby.* 806.

Marshes, June, P. Common.

661. *Loose Sedge.* — *C. distans.*

Spikes very remote, nearly sitting, floral leaf sheathing, capsules angular, sharp-pointed.

Ditches, June, P. Rare.

Stevington.

Remote

I House. III Males. Monoec. Triand.

662. *Remote Sedge*. — *C. remota*.

Spikes ovate, nearly sitting, remote, both fertile and barren, floral leaves as long as the straw. — Leers. XVI. Flora Danica. 307.

Ditch Banks, June, P. Common. **Sowerby** 832.

663. *Turfy Sedge*. — *C. cæspitosa*.

Spikes upright, cylindrical, three, nearly sitting, barren one terminating, straw 3-square. — Curtis. IV. 62.

Meadows, May, P. Common.

664. *Flea Sedge*. — *C. pulicaris*.

Spike simple, with fertile and barren flowers, male flowers on the upper part, capsules diverging, bent back. — Leers. XIV. 1. Flora Danica. 166. **Sowerby** 1051.

Meadows, Bogs, June, P. Rare.

Eversholt.

665. *Grey Sedge*. — *C. divulsa*.

Spikelets roundish, barren and fertile, lower ones distant, capsules pointed, upright.

Woods — *Bogs*, May, P. Rare. **Sowerby** 629.

Clapham, — *Everton*, — *Potton*. *Rutnæ Wood*.
Twin Wood.

666. *Panicled Sedge*. — *C. paniculata*. **S.D.**

Spikes barren and fertile, bunch compound.

Leers. XIV. 4. **Sowerby** 1064.

D d 2

Moors,

I House. III Males. Monoec. Triand.

Moors, June, P. Common.

667. Mountain Sedge. — C. montana. S.D.

Fertile spikes sitting, mostly single, egg-shaped, near to the barren spike, straw naked, capsules downy. — Leers. XVI. 6.

Mountainous Meadows, June, P. Rare.

Clapham-Park Pastures.

668. Yellow Sedge. — C. flava.

Spikes crowded, nearly sitting, roundish, male spike strap-shaped, capsules pointed, bowed back. — Leers. XV. 6. Flora Danica. 371. Sowerby. 1294.

Bogs, June, P. Rare.

Amptbill, — Stevington.

669. Pill-bearing Sedge. — C. pilulifera. S.D.

Spikes terminal, crowded, roundish, male spike oblong. — Flora Danica. 444.

Sowerby. 885.
Bogs, June, P. Rare.

Eversholt.

Found by the Rev. J. HEMSTED.

670. Pale Sedge. — C. pallescens.

Spikes hanging down, male upright, females ovate, tiled, capsules crowded, blunt. — Leers. XV. 4.

Woods,

* 669 *Larex extensa*

Flora Brit.
Amptbill-bogs. 1005.

I House. III Males. Monoec. Triand.

Woods, June, P. Rare.

Putnoe Wood. 1001.

671. *Early Sedge.* — *C. præcox.*

Spikes upright, nearly sitting, contiguous,
male spike oblong, larger, capsules downy. —
Flora Danica. 403. *Sowerby* 1099.

Heaths, Meadows, April, P. Common.

672. *Pink Sedge.* — *C. panicea.*

Spikes on fruitstalks, upright, remote, female
spikes strap-shaped, capsules swollen, blun-
tish. — *Leers. XV. 5.* *Sowerby* 1505.

Wet Meadows, June, P. Common.

673. *Wood Sedge.* — *C. sylvatica.*

Spikes on fruitstalks, hanging down, male
spike upright, females thread-shaped, loose,
shorter than the fruitstalk, capsules ovate,
awned, forked. — *Leers. XV. 2.* *Sowerby* 995.

Woods, June, P. Common.

674. *Slender-eared Sedge.* — *C. strigosa.*

Spikes on fruitstalks, pendant, the barren one
upright, fertile ones thread-shaped, loose,
capsules roundish, taper pointed. *Sowerby* 994.

Woods, Hedges, May, P. Rare.

Putnoe Wood, — *Renhold.*

Pendulous

I House. III Males. Monoec. Triand.

675. *Pendulous Sedge*. — C. pendula. **SD.** *sent*

Spikes nearly sitting, drooping, male spike upright, females cylindrical, very long, capsules roundish, pointed. — Curtis. III. 63.

Woods, June, P. Common.

676. *Bank Sedge*. — C. riparia.

Male spikes many, 3-square, blackish, pointed, scales awn-pointed, capsules somewhat swollen, with 2 horns. — Curtis. IV. 60.

Sowerby. 579.

Ditches, April, P. Common.

677. *Sharp Sedge*. — C. acuta.

Male spikes many, blunt, scales rather blunt, stem with sharp angles. — Curtis. IV. 61.

Wet Places, April, P. Common.

678. *Heath Sedge*. — C. recurva. **SD.** *sent*

Spikes crowded, on fruitstalks, cylindrical, somewhat pendant, the barren one terminal, capsules tilted, bluntnish. *Sowerby*. 1506.

Pastures, May, P. Common.

679. *Beaked Sedge*. — C. rostrata. **SD.**

Male spikes numerous, females mostly on fruitstalks, capsules swollen, round, pointed, straddling. — Leers. XVI. 2. *Sowerby*. 780.

Moors, June, P. Rare. *Campu llacea*.

Amptbill, — *Potton*.

Hairy

I House. III Males. Monoec. Triand.

680. *Hairy Sedge*. — *C. hirta*.

Spikes remote, males numerous, females mostly on fruitstalks, upright, capsules hairy. — Leers. XVI. 3. *Sowerby* 685.

Meadows, June, P. Common.

Obs:

Sheaths of the leaves hairy.

I House. IV Males. Monoec. Tetrand.

Birch. *BETULA*. Gen. Pl. 1419.

Male. — Cup of 1 leaf, 3 segments, 3 flowers. Blossom 4 segments.

Female. — Cup of 1 leaf, mostly 3 segments and 3 flowers. Seeds with a winged membrane on each side.

681. *Common Birch*. — *B. alba*.

Leaves ovate, pointed, serrated. — Blackwell.

240.

Woods in a sandy Soil, April, T. Common.

682. *Alder-tree*. — *B. Alnus*.

Fruitstalks branched. — Johnson's Gerard.

1477. 1. *Sowerby* 1508 —

Moist Places, March, T. Rare.

Fenlake.

Nettle.

I House. IV Males. Monoec. Tetrand.

Nettle. URTICA. Gen. Pl. 1422.

Male. — Cup of 4 leaves. Blossom none.
Honey-cup central, in the shape of a
drinking glass.

Female. — Cup of 2 valves. Blossom none.
Seed 1, shining.

683. *Small Stinging Nettle.* — U. urens.

Leaves opposite, oval. — Curtis. VI. 70.
Dunghills, July, A. Common.

684. *Great Stinging Nettle.* — U. dioica.

Leaves opposite, heart-shaped, bunches in
pairs. — Curtis. VI. 69.
Road-sides, July, P. Common.

I House. Many Males. Monoec. Polyand.

Milfoil. MYRIOPHYLLUM. Gen. Pl. 1440.
Male. — Cup of 4 leaves. Petals 4, falling
off.

Female. — Cup of 4 leaves. Petals 4, fal-
ling off. Style none. Seeds 4, naked.

685. *Spiked Milfoil.* — M. spicatum.

Flowers in spikes, spikes naked, interrupted,
terminal. — Sowerby. 83. excellent.

Slow Streams, July, P. Rare.

Ford-End,—Castle Mills.

Whorled

I House. Many Males. Monoec. Polyand.

686. *Whorled Milfoil.* — *M. verticillatum.*

Flowers in whorls, in the bosoms of the leaves. — Sowerby. 218. excellent.

Ditches, July, P. Common.

Arrow-head. *SAGITTARIA.* Gen. Pl. 1441.

Male. — Cup of 3 leaves. Blossom of 3 petals.

Female. — Cup of 3 leaves. Blossom of 5 petals. Seeds many, naked.

687. *Common Arrow-head.* — *S. sagittifolia.*

Leaves pointed, arrow-shaped. — Sowerby. 84.

Rivers, July, P. Common.

Burnet. *POTERIUM.* Gen. Pl. 1445.

Male. — Cup of 4 leaves. Blossom of 4 segments.

Female. — Cup of 4 leaves. Blossom of 4 segments. Berry formed of the hardened tube of the blossom.

688. *Common Burnet.* — *P. Sanguisorba.*

Without thorns, stems somewhat angular. — Curtis. II. 64. Martyn. 69. Sowerby. 860.

Hilly Pastures, June, P. Common.

I House. Many Males. Monoec. Polyand.

Oak. *QUERCUS.* Gen. Pl. 1447.

Male. — Cup of 5 segments. Blossom none.

Female. — Cup of 1 leaf, very entire, rough.

Blossom none. Seed 1, ovate.

689. *Common Oak, or Britain's Glory.* — *Q. Robur.*

Leaves falling off, oblong, broadest towards the end, waves rather sharp, angles blunt.— Blackwell. 487. Sowerby. 1342.

Woods, April, T. Common.

690. *Long-stalked Oak.* — *Q. pedunculata.* SD.

Acorns on long fruitstalks. — Martyn. 10. excellent.

Woods, May, T. Rare.

Clapham Park Wood.

Beech. *FAGUS.* Gen. Pl. 1448.

Male. — Cup 5 segments, bell-shaped.

Blossom none.

Female. — Cup with 4 teeth. Blossom none. Capsule now, what was the cup before, 4-valved, 2-seeded.

691. *Common Beech.* — *F. sylvatica.*

Leaves ovate, indistinctly serrated. — Johnson's Gerard. 1444.

Woods, April, T. Common.

Horn Beam. *Carpinus.*

Male. Blossom none. Cup 1 leaf: a fringed scale, stamens 18 to 20. Female. Gums 2 with 2 styles on each, nut eggshaped.

* 691. Common Hornbeam Tree. *C. Betulus.*

Scales of the cones flat.

Woods & Hedges T. May. Rare. Oakley. Hawnes. Chicksands.

I House. Many Males. Monoec. Polyand.

692. *Common Chesnut.* — F. *Castanea.*

Leaves spear-shaped, serrated, with tapering points, naked underneath.—Blackwell. 330. Sowerby. 886.

Woods, May, T. Rare.

Silsoe.

Hazel. *CORYLUS.* Gen. Pl. 1450.

Male. — Cup of 1 leaf, 3 segments like a scale, 1-flowered. Blossom none.

Female. — Cup of 2 leaves, torn. Blossom none. Styles 2. Nut ovate.

693. *Hazel-nut.* — C. *avellana.*

Leafscales ovate, blunt. — Blackwell. 293. Sowerby. 723.

Woods, Hedges, March, S. Common.

C L A S S. XXII.

Character of the XXIId. Class.

Male and female organs on separate plants.

II Houses. II Males. Dioec. Diand.

Willow. *SALIX.* Gen. Pl. 1493.

Male. — Catkin with scales. Blossom none. A honey-cup gland at the base.

Female. — Catkin with scales. Blossom none. Capsule 1-celled, 2-valved. Seeds downy.

in II Houses. of II Males. Dioec. Diand.

694. Three-threaded Willow. — *S. triandra*.

Leaves serrated, smooth, flowers with three males. — Curtis. VI. 72. Sowerby 1435.

~~Osier-holts, May, S. Rare.~~

above Bedford. Thurleugh, — Fenlake.

1799. Crack Willow. — *S. fragilis*.

Leaves serrated, smooth, ovate-spear-shaped, leafstalks with glandular teeth. — Woodville. 198.

Banks and Hedges, April, S. Common.

696. One-threaded Willow. — *S. monandra*.

Sowerby 1343. Leaves serrated, smooth, strap-spear-shaped, upper ones oblique. — Curtis. VI. 71.

~~Osier holts. Banks and Hedges, May, S. Rare.~~

above Bedford. — Thurleugh.

1799. 697. Eared Willow. — *S. aurita*.

Leaves very entire, woolly on both sides, somewhat ovate, with appendages. — Hoff-

Sowerby 1487. man. His. Salic. IV. 1. 2.

Woods and Hedges, April, S. Rare.

Warden.

698. Common Sallow. — *S. caprea*.

Sowerby 1488. Leaves ovate, wrinkled, woolly underneath, waved, with little teeth towards the end. — Hoffman. His. Salic. III. 1. 2.

Hedges, April, S. Common.

Osier

694. Obs.

The central thread in the flower is the longest —

obs. 696.

Catkin at first purplish red, growing yellower as it expands.

Bugs. June 8.

*: Pointed Sallow. *S. acuminata*.

Leaves ovate-oblong, woolly underneath, upper ones entire, those below crenated.

Sowerby 1434. Hoffman. His. Salic. VI. 1. 2. & XXII. f. 2.

Hedges, Woods. *S. April*. Rare.

Clapham Woods.

*699. Wood Willow - *S. caprea*.

Leaves ovate, wrinkled, woolly beneath,
waved, denticulated above.

Hoffman His. Salic. III. 1. 2.

Woods S. April - Rare.

Clophill & Clapham Park Woods.

*699 Almond-leaved
Peach-leaved Willow. *S. amygdalina*.

Leaves serrated, smooth, spear-shaped
on leafstalks, leafscales shaped like a
blunt square.

Osier holtz & hedges. T. April. Rare.
Bromham Lane

213

II Houses. II Males. Dioec. Diand.

699. Osier Willow. — *S. viminalis*.

Leaves for the most part entire, spear-shaped,
very long, sharp; silky underneath, branch-
es rod-like.—Hoffman. His. Salic. II. 1. 2.

Osier-holts, April, S. Common.

700. Common Willow. — *S. alba*.

Leaves spear-shaped, pointed, downy on both
sides, lower serratures glandular.—Hoff-
man. His. Salic. VII. 1.

Banks of Rivers, May, T. Common.

701. Creeping Willow. — *S. repens*.

Leaves very entire, spear-shaped, naked on
both sides, stem creeping.—Sowerby. 183.
excellent.

Bogs, June, S. Rare.

Amptbill.

Obs:

Root woody, creeping. Stems little-branched, somewhat up-
right, leaves white underneath. Style ovate, blunt at the end,
simple. Stigma divided into 4, yellowish.

Mistletoe. *VISCUM*. Gen. Pl. 1504.

Male. — Cup of 4 segments. Blossom none.

Threads none. Anthers growing to the
cup.

Female. — Cup of 4 leaves, superior. Style
none. Blossom none. Berry 4 seeds.
Seeds heart-shaped.

Common

II Houses. II Males. Dioec. Diand.

702. Common Mistletoe. — *V. album.*

Leaves heart-shaped, blunt, stem forked,
spikes from the bosoms of the leaves. —
Woodville. 270. excellent. Sowerby. 1470

Parasitical on Trees, May, S. Common.

II Houses. V Males. Dioec. Pentand.

Hop. HUMULUS. Gen. Pl. 1523.

Male. — Cup of 5 leaves. Blossom none.

Female. — Cup of 5 leaves, expanding obliquely, entire. Blossom none. Seed 1,
within a leafy cup.

703. Common Hop. — *H. Lupulus.*

Sowerby. 427.

Hedges, June, P. Common.

II Houses. VI Males. Dioec. Hexand.

Ladies' Seal. TAMUS. Gen. Pl. 1527.

Male. — Cup of 6 segments. Blossom none.

Female. — Cup of 6 segments. Blossom none. Style 3-cleft. Berry 3-celled, inferior. Seeds 2.

704. Black Briony. — *T. communis.*

Leaves heart-shaped, entire.—Sowerby. 91.

Hedges, June, P. Common.

Poplar.

II Houses. VIII. Males. Dioec. Octand.

Poplar. *POPULUS.* Gen. Pl. 1531.

Male. — Cup of the catkin a torn scale.
Blossom top-shaped, oblique, entire.

Female. — Cup of the catkin and blossom
as in the male. Stigma 4-cleft. Cap-
sule 2-celled. Seeds many, downy.

705. *White Poplar.* — *P. alba.*

Leaves nearly round, tooth-angular, woolly
underneath. — Johnson's Gerard. 1486. 1. *Sowerby.* 1610.

Hedges, March, T. Rare.

Fenlake, — *Stevington.*

706. *Aspen Tree.* — *P. tremula.*

Leaves nearly round, tooth-angular, smooth
on both sides. — Johnson's Gerard. 1487. 3.

Woods, March, T. Common.

707. *Black Poplar.* — *P. nigra.*

Leaves trowel-shaped, pointed, serrated. —
Blackwell. 248. 1.

Woods, Hedges, March, T. Rare.

Sheerbatch.

II Houses. IX Males. Dioec. Enneand.

Mercury. *MERCURIALIS.* Gen. Pl. 1534.

Male. — Cup of 3 segments. Blossoms
none. Anthers round, double.

Female. — Cup of 3 segments. Blossom
none. Capsule 1 berry, 2 cells.

Dog's

II Houses. IX Males. Dioec. Enneand.

708. *Dog's Mercury*. — *M. perennis*.

Stem very simple, leaves rough. — Curtis.

II. 65.

Woods, Hedges, April, P. Common.

Frog-bit. *HYDROCHARIS*. Gen. Pl. 1535.

Male. — Sheath of 2 leaves. Cup of 3 segments. Blossom 3 petals. Threads, 3 innermost bearing styles.

Female. — Cup of 3 segments. Blossom 3 petals. Capsule 6 cells, many seeds, inferior.

709. *Common Frog-bit*. — *H. Morsus Ranae*.

Curtis. III. 64. *Sowerby*. 808.

Ditches near Bigleswade, on the way to Northill.
Slow Streams, July, P. Rare.

— *Castle Mills*, — *Fentake*.

II Houses. Anthers united. Dioec. Syngen.

Bryony. *BRYONIA*. Gen. Pl. 1480.

Male. — Cup of 5 teeth. Blossom with 5 divisions. Threads 3. Anthers 5.

Female. — Cup of 5 teeth. Blossom with 5 divisions. Style with 3 segments. Berry roundish, with many seeds.

White

10

Knee Holly. *Ruscus*. Gen. Pl.

Male. — Cup of 6 leaves. Blossom none. Nectary central, ovate, ^{open} perforated at the point. top.

Female. Cup, Blossom and Nectary as in the male. Style simple. Berry 3-celled. Seeds. 2.

*710. Common Knee Holly. *R. aculeatus*. bearing the flowers surface. Leaves bare, flowering on the upper side. naked ~~channel~~ 159. Woodville. 237.

Hedge Rows and Spinneys. S. February. Rare.
Oakley Raynes. *Sowerby*. 560.

II Houses. Anthers united. Dioec. Syngen.

710. *White Bryony*. — *B. dioica*.

Leaves hand-shaped, rough on both sides, with callous points; flowers, male and female on separate plants. — Woodville. 189.
Sowerby. 439.

Woods, Hedges, July, P. Common.

C L A S S. XXIII.

Character of the XXIIId. Class.

This Class consists of plants in which some of the flowers in the same species have both male and female; some only male, and others only female organs.

Various dispositions. Polygamia.

I House. Polyg. Monoec.

Soft Grass. *HOLCUS*. Gen. Pl. 1565.

Hermaphrodite. — Cup a chaff, containing 2 florets. Blossom an awned chaff. Seed 1.

Male. — Cup a chaff of 2 valves. Blossom none.

711. *Creeping Soft Grass*. — *H. mollis*.

Root creeping, joints woolly, awn lengthened out beyond the spikelet. — Curtis. V. 8. Martyn. 119. Sowerby. 1170.

F f Woods,

I House. IV Males. Monoec. Tetrand.

Nettle. URTICA. Gen. Pl. 1422.

Male. — Cup of 4 leaves. Blossom none.
Honey-cup central, in the shape of a
drinking glass.

Female. — Cup of 2 valves. Blossom none.
Seed 1, shining.

683. *Small Stinging Nettle.* — U. urens.

Leaves opposite, oval. — Curtis. VI. 70.
Dunghills, July, A. Common.

684. *Great Stinging Nettle.* — U. dioica.

Leaves opposite, heart-shaped, bunches in
pairs. — Curtis. VI. 69.
Road-sides, July, P. Common.

I House. Many Males. Monoec. Polyand.

Milfoil. MYRIOPHYLLUM. Gen. Pl. 1440.
Male. — Cup of 4 leaves. Petals 4, falling
off.

Female. — Cup of 4 leaves. Petals 4, fal-
ling off. Style none. Seeds 4, naked.

685. *Spiked Milfoil.* — M. spicatum.

Flowers in spikes, spikes naked, interrupted,
terminal. — Sowerby. 83. excellent.

Slow Streams, July, P. Rare.

Ford-End,—Castle Mills.

Whorled

I House. Many Males. Monoec. Polyand.

686. *Whorled Milfoil.* — *M. verticillatum.*

Flowers in whorls, in the bosoms of the leaves. — Sowerby. 218. excellent.

Ditches, July, P. Common.

Arrow-head. *SAGITTARIA.* Gen. Pl. 1441.

Male. — Cup of 3 leaves. Blossom of 3 petals.

Female. — Cup of 3 leaves. Blossom of 5 petals. Seeds many, naked.

687. *Common Arrow-head.* — *S. sagittifolia.*

Leaves pointed, arrow-shaped. — Sowerby. 84.

Rivers, July, P. Common.

Burnet. *POTERIUM.* Gen. Pl. 1445.

Male. — Cup of 4 leaves. Blossom of 4 segments.

Female. — Cup of 4 leaves. Blossom of 4 segments. Berry formed of the hardened tube of the blossom.

688. *Common Burnet.* — *P. Sanguisorba.*

Without thorns, stems somewhat angular. — Curtis. II. 64. Martyn. 69. Sowerby. 860.

Hilly Pastures, June, P. Common.

I House. Many Males. Monoec. Polyand.

Oak. *QUERCUS.* Gen. Pl. 1447.

Male. — Cup of 5 segments. Blossom none.

Female. — Cup of 1 leaf, very entire, rough.

Blossom none. Seed 1, ovate.

689. *Common Oak, or Britain's Glory.* — *Q. Robur.*

Leaves falling off, oblong, broadest towards the end, waves rather sharp, angles blunt.— Blackwell. 487. Sowerby. 1342.

Woods, April, T. Common.

690. *Long-stalked Oak.* — *Q. pedunculata.* SD.

Acorns on long fruitstalks. — Martyn. 10. excellent.

Woods, May, T. Rare.

Clapham Park Wood.

Beech. *FAGUS.* Gen. Pl. 1448.

Male. — Cup 5 segments, bell-shaped.

Blossom none.

Female. — Cup with 4 teeth. Blossom none. Capsule now, what was the cup before, 4-valved, 2-seeded.

691. *Common Beech.* — *F. sylvatica.*

Leaves ovate, indistinctly serrated. — Johnson's Gerard. 1444.

Woods, April, T. Common.

Horn Beam. Larpius.

Male. Blossom none. Cup 1 leaf: a fringed scale, stamens 18 to 20. Female. Gums 2 with 2 styles on each, nut eggshaped.

* 691. Common Hornbeam Tree. *C. Betulus.*

Scales of the cones flat.

Woods & Hedges T. May. Rare. Oakley. Hawnes. Chicksands.

I House. Many Males. Monoec. Polyand.

692. *Common Chesnut.* — F. *Castanea.*

Leaves spear-shaped, serrated, with tapering points, naked underneath.—Blackwell. 330. Sowerby. 886.

Woods, May, T. Rare.

Silsoe.

Hazel. *CORYLUS.* Gen. Pl. 1450.

Male. — Cup of 1 leaf, 3 segments like a scale, 1-flowered. Blossom none.

Female. — Cup of 2 leaves, torn. Blossom none. Styles 2. Nut ovate.

693. *Hazel-nut.* — C. *avellana.*

Leafscales ovate, blunt. — Blackwell. 293. Sowerby. 723.

Woods, Hedges, March, S. Common.

C L A S S. XXII.

Character of the XXIId. Class.

Male and female organs on separate plants.

II Houses. II Males. Dioec. Diand.

Willow. *SALIX.* Gen. Pl. 1493.

Male. — Catkin with scales. Blossom none. A honey-cup gland at the base.

Female. — Catkin with scales. Blossom none. Capsule 1-celled, 2-valved. Seeds downy.

in II Houses. of II Males. Dioec. Diand.

694. Three-threaded Willow. — *S. triandra*.

Leaves serrated, smooth, flowers with three males. — Curtis. VI. 72. Sowerby 1435.

~~Osier-holts~~, May, S. Rare.
above Bedford.

1799 Thurleugh, — Fenlake.

695. Crack Willow. — *S. fragilis*.

Leaves serrated, smooth, ovate-spear-shaped, leafstalks with glandular teeth. — Woodville. 198.

Banks and Hedges, April, S. Common.

696. One-threaded Willow. — *S. monandra*.

~~Sowerby 1343.~~ Leaves serrated, smooth, strap-spear-shaped, upper ones oblique. — Curtis. VI. 71.
~~Osier holts.~~ Banks and Hedges, May, S. Rare.

above Bedford — Thurleugh.

1799. 697. Eared Willow. — *S. aurita*.

Leaves very entire, woolly on both sides, somewhat ovate, with appendages. — Hoff-

~~Sowerby 1487.~~ man. His. Salic. IV. 1. 2.

Woods and Hedges, April, S. Rare.

Warden.

698. Common Sallow. — *S. caprea*.

~~Sowerby 1488.~~ Leaves ovate, wrinkled, woolly underneath, waved, with little teeth towards the end. — Hoffman. His. Salic. III. 1. 2.

Hedges, April, S. Common.

Osier

694. Obs.

The central thread in the flower is the longest —

obs. 696.

Catkin at first purplish red, growing yellower as it expands.

Bugs. June 8.

*: Pointed Sallow. *S. acuminata*.

Leaves ovate-oblong, woolly underneath, upper ones entire, those below crenated.

~~Sowerby 1434.~~ Hoffman His. Salic. VI. 1. 2. & XXII. f. 2.

Hedges, Woods. *S. April*. Rare.

Clapham Woods.

*699. Wood Willow - *S. caprea*.

Leaves ovate, wrinkled, woolly beneath,
waved, denticulated above.

Hoffman His. Salic. III. 1. 2.

Woods S. April - Rare.

Clophill & Clapham Park Woods.

*699 Almond-leaved
Peach-leaved Willow. *S. amygdalina*.

Leaves serrated, smooth, spear-shaped
on leafstalks, leafscales shaped like a
blunt square.

Osier holtz & hedges. T. April. Rare.
Bromham Lane

213

II Houses. II Males. Dioec. Diand.

699. Osier Willow. — *S. viminalis*.

Leaves for the most part entire, spear-shaped,
very long, sharp; silky underneath, branch-
es rod-like.—Hoffman. His. Salic. II. 1. 2.

Osier-holts, April, S. Common.

700. Common Willow. — *S. alba*.

Leaves spear-shaped, pointed, downy on both
sides, lower serratures glandular.—Hoff-
man. His. Salic. VII. 1.

Banks of Rivers, May, T. Common.

701. Creeping Willow. — *S. repens*.

Leaves very entire, spear-shaped, naked on
both sides, stem creeping.—Sowerby. 183.
excellent.

Bogs, June, S. Rare.

Amptbill.

Obs:

Root woody, creeping. Stems little-branched, somewhat up-
right, leaves white underneath. Style ovate, blunt at the end,
simple. Stigma divided into 4, yellowish.

Mistletoe. *VISCUM*. Gen. Pl. 1504.

Male. — Cup of 4 segments. Blossom none.

Threads none. Anthers growing to the
cup.

Female. — Cup of 4 leaves, superior. Style
none. Blossom none. Berry 4 seeds.
Seeds heart-shaped.

Common

II Houses. II Males. Dioec. Diand.

702. Common Mistletoe. — *V. album.*

Leaves heart-shaped, blunt, stem forked,
spikes from the bosoms of the leaves. —
Woodville. 270. excellent. Sowerby. 1470

Parasitical on Trees, May, S. Common.

II Houses. V Males. Dioec. Pentand.

Hop. HUMULUS. Gen. Pl. 1523.

Male. — Cup of 5 leaves. Blossom none.

Female. — Cup of 5 leaves, expanding obliquely, entire. Blossom none. Seed 1,
within a leafy cup.

703. Common Hop. — *H. Lupulus.*

Sowerby. 427.

Hedges, June, P. Common.

II Houses. VI Males. Dioec. Hexand.

Ladies' Seal. TAMUS. Gen. Pl. 1527.

Male. — Cup of 6 segments. Blossom none.

Female. — Cup of 6 segments. Blossom none. Style 3-cleft. Berry 3-celled, inferior. Seeds 2.

704. Black Briony. — *T. communis.*

Leaves heart-shaped, entire.—Sowerby. 91.

Hedges, June, P. Common.

Poplar.

II Houses. VIII. Males. Dioec. Octand.

Poplar. *POPULUS.* Gen. Pl. 1531.

Male. — Cup of the catkin a torn scale.
Blossom top-shaped, oblique, entire.

Female. — Cup of the catkin and blossom
as in the male. Stigma 4-cleft. Cap-
sule 2-celled. Seeds many, downy.

705. *White Poplar.* — *P. alba.*

Leaves nearly round, tooth-angular, woolly
underneath. — Johnson's Gerard. 1486. 1. Sowerby. 1610.

Hedges, March, T. Rare.

Fenlake, — Stevington.

706. *Aspen Tree.* — *P. tremula.*

Leaves nearly round, tooth-angular, smooth
on both sides. — Johnson's Gerard. 1487. 3.

Woods, March, T. Common.

707. *Black Poplar.* — *P. nigra.*

Leaves trowel-shaped, pointed, serrated. —
Blackwell. 248. 1.

Woods, Hedges, March, T. Rare.

Sheerbatch.

II Houses. IX Males. Dioec. Enneand.

Mercury. *MERCURIALIS.* Gen. Pl. 1534.

Male. — Cup of 3 segments. Blossoms
none. Anthers round, double.

Female. — Cup of 3 segments. Blossom
none. Capsule 1 berry, 2 cells.

Dog's

II Houses. IX Males. Dioec. Enneand.

708. *Dog's Mercury*. — *M. perennis*.

Stem very simple, leaves rough. — Curtis.

II. 65.

Woods, Hedges, April, P. Common.

Frog-bit. *HYDROCHARIS*. Gen. Pl. 1535.

Male. — Sheath of 2 leaves. Cup of 3 segments. Blossom 3 petals. Threads, 3 innermost bearing styles.

Female. — Cup of 3 segments. Blossom 3 petals. Capsule 6 cells, many seeds, inferior.

709. *Common Frog-bit*. — *H. Morsus Ranae*.

Curtis. III. 64. *Sowerby*. 808.

Ditches near Bigleswade, on the way to Northill
Slow Streams, July, P. Rare.

— *Castle Mills*, — *Fentake*.

II Houses. Anthers united. Dioec. Syngen.

Bryony. *BRYONIA*. Gen. Pl. 1480.

Male. — Cup of 5 teeth. Blossom with 5 divisions. Threads 3. Anthers 5.

Female. — Cup of 5 teeth. Blossom with 5 divisions. Style with 3 segments. Berry roundish, with many seeds.

White

100

Knee Holly. *Ruscus*. Gen. Pl.

Male. — Cup of 6 leaves. Blossom none. Nectary central, ovate, ^{open} perforated at the point. top.

Female. Cup, Blossom and Nectary as in the male. Style simple. Berry 3-celled. Seeds. 2.

*710. Common Knee Holly. *R. aculeatus*. bearing the flowers surface. Leaves bare, flowering on the upper side. naked ~~channel~~ 159. Woodville. 237.

Hedge Rows and Spinneys. S. February. Rare.
Oakley Raynes. *Sowerby*. 560.

II Houses. Anthers united. Dioec. Syngen.

710. *White Bryony*. — *B. dioica*.

Leaves hand-shaped, rough on both sides, with callous points; flowers, male and female on separate plants. — Woodville. 189.
Sowerby. 439.

Woods, Hedges, July, P. Common.

C L A S S. XXIII.

Character of the XXIIId. Class.

This Class consists of plants in which some of the flowers in the same species have both male and female; some only male, and others only female organs.

Various dispositions. Polygamia.

I House. Polyg. Monoec.

Soft Grass. *HOLCUS*. Gen. Pl. 1565.

Hermaphrodite. — Cup a chaff, containing 2 florets. Blossom an awned chaff. Seed 1.

Male. — Cup a chaff of 2 valves. Blossom none.

711. *Creeping Soft Grass*. — *H. mollis*.

Root creeping, joints woolly, awn lengthened out beyond the spikelet. — Curtis. V. 8. Martyn. 119. Sowerby. 1170.

F f Woods,

I House. Polyg. Monoec.

Woods, Hedges, July, P. Common.

712. Meadow Soft Grass. — *H. lanatus*.

Chaff with 2 flowers, woolly, hermaphrodite floret awnless, male one awned, awn bowed back. — Curtis. IV. 11. Martyn. 118.

Pastures, June, P. Common. *Sowerby. 1169.*

Crosswort. VALANTIA. Gen. Pl. 1575.

Hermaphrodite. — Cup none. Blossom with 4 divisions. Style 2-cleft. Seed 1.

Male. — Cup none. Blossom 3 or 4 divisions. Female organ indistinct.

713. Common Crosswort. — *V. Cruciatum*.

Male flowers with 4 segments, fruitstalks with 2 leaves. — Sowerby. 143. excellent.

Woods, Hedges, May, P. Common.

Pellitory. PARIETARIA. Gen. Pl. 1576.

Hermaphrodite. — Cup with 4 segments. Blossom none. Threads 4. Style 1. Seed 1, superior, lengthened out.

714. Common Pellitory. — *P. officinalis*.

Leaves spear-ovate, fruitstalks forked, cups with 2 leaves. — Curtis. IV. 63. *Sowerby. 879.*

Walls, June, P. Common.

Orach.

I House. Polyg. Monoec.

Orach. ATRIPLEX. Gen. Pl. 1577.

Hermaphrodite. — Cup of 5 leaves. Blossom none. Threads 5. Style with 2 segments. Seed 1, depressed.

Female. — Cup of 2 leaves. Blossom none. Threads none. Style with 2 segments. Seed 1, compressed.

715. *Spear Orach.* — *A. hastata.*

Stem herbaceous, valves of the cup in the female florets large, trowel-shaped, waved. — Curtis. II. 66.

716. *Spreading Orach.* — *A. patula.*

Stem herbaceous, spreading, leaves somewhat trowel-spear-shaped, cups of the fruit toothed in the disc. — Johnson's Gerard. 326. Sowerby 936.

Dunghills, August, A. Common.

Maple. ACER. Gen. Pl. 1590.

Hermaphrodite. — Cup 5-cleft. Blossom 5 petals. Capsule 2 or 3-celled, with one seed ended by a wing.

Male. — Cup 5-cleft. Blossom 5 petals.

717. *Sycamore.* — *A. pseudo-platanus.*

Leaves with 5 lobes, unequal, serrated, flowers in bunches. — Sowerby. 303.

Woods, May, T. Rare.

Renhold Wood.

F f 2

Common

I House. Polyg. Monoec.

718. Common Maple. — *A. campestris*.

Leaves lobed, blunt, nicked.—Sowerby. 304.

Woods, June, T. Common.

II Houses. Polyg. Dioec.

Ash. *FRAXINUS*. Gen. Pl. 1597.

Hermaphrodite. — Cup none. Blossom
none. Seed 1, spear-shaped.

Female. — Female organ 1. Seed 1, spear-
shaped.

719. Common Ash-tree. — *F. excelsior*.

Leaves serrated, flowers without petals. —

Johnson's Gerard. 1472. Sowerby. 1692.

Hedges, April, T. Common.



CRYPTOGAMIA

CLASS 1.

Character of the ANTHOCYANAE.

CLASS 2. FERN. CRYPTOGAMS.

Division 1. POLYPSORIACE. Gen. Pl. 1594.

Fronds simple, supported in various ways,
club-shaped stalk, variegated, or not.

Wood Ferns. — *E. sylvaticus*.

Stern hairy, starchy, fronds compound.

Boston fern.

Maidenhair. — *Adiantum*.

False Maidenhair. — *Asplenium*.

False Maidenhair. — *Asplenium*.

C R Y P T O G A M I A.

Conjugate Nucleus, R. Common

Spores of the genus may have 1 or 2 eyes

— *Leaf-wall shrub*. 22

— *Bolton*. 22

— *Class. XXIV.* 22

Character of the XXIVth. Class.

Flowers not visible.

Stems scarcely jointed near the base

XXIV. Ferns. Crypt. Filices.

Horse-tail. EQUISETUM. Gen. Pl. 1614.

Fructifications disposed in an ovate-oblong, club-shaped spike, whorled, on a stem.

720. *Wood Horse-tail.* — *E. sylvaticum.*

Stem bearing a spike, fronds compound. — Bolton. 33.

Woods, April, P. Rare.

Hawnes Wood, — Ampthill Park.

Obs: *can be broken*

Fronds simple in the seedling plants, in the more adult much whorled and branched.

Corn

XXIV. Ferns. Crypt. Filices.

721. *Corn Horse-tail*. — *E. arvense*.

Fertile stalk naked, barren one leafy. —

Curtis, IV. 64.

Cornfields, March, P. Common.

*Obs:**Sheaths of the joints with from 4 to 5 teeth.*722. *Marsh Horse-tail*. — *E. palustre*.

Stem angular, fronds undivided. — Bolton.

35.

Marshes, June, P. Rare.

Ampthill Moor, — *Stevington Bogs*.723. *River Horse-tail*. — *E. fluviatile*.

Stem scored, fronds nearly simple. — Bolton.

36.

Marshy Places, May, P. Common.

724. *Smooth Horse-tail*. — *E. limosum*.Stem somewhat naked, smooth. — Ray's Synopsis. V. 2. *Sowerby*. 929.

Rivers, May, P. Common.

*Obs:**Sheaths of the joints more sharply and narrowly toothed than in the foregoing.*725. *Rough Horse-tail*. — *E. hyemale*.

Stem naked, rough to the touch, somewhat branched at the base. — Bolton. 39.

Sowerby. 915.

Bogs,

XXV. Ferns. Crypt. Filices.

Bogs, July, P. Rare.

Sporangia numerous, scattered over surface.

XXIV. Ferns. Crypt. Filices.

Bogs, July, P. Rare.

*Potton Marshes, — Ampthill Bogs.**Obs.:*

Spike very slender. Sheaths of the joints pale, not infrequently growing in my specimens at 3 inches distance from each other.

Adder's-tongue. *OPHIOGLOSSUM.* Gen. Pl. 1621.

Capsules numerous, growing together on a membrane into a double rowed spike, nearly round; when ripe opening transversely. Ring none. Seeds many, very small.

726. *Common Adder's-tongue.* — *O. vulgatum.*

Frond ovate. — Sowerby. 108.

Meadows, May, P. Common.

Obs.:

Root fibrous, leaf indistinctly heart-shaped, and bearing a 2-rowed spike on a long fruitstalk.

Moonwort. *OSMUNDA.* Gen. Pl. 1622.

Capsules distinct, disposed in a bunch, crowded, sitting, nearly round, opening transversely. Ring none. Seeds many, very small.

727. *Common Moonwort.* — *O. Lunaria.*

Bunch lateral, leaf winged, solitary. — Sowerby. 318.

Pastures, May, P. Rare.

Oakley

XXIV. Ferns. Crypt. Filices.

Oakley West Field.

Obs :

*Root fibrous, seeds in panicles, leaflets kidney-shaped.*728. Royal Osmund. — *O. regalis.*Leaves with 2 rows of wings, with bunches
at the end. — Sowerby. 209.

Boggy Places, July, P. Rare.

Aspley Wood.

Found by Mr. GEORGE DIXON, of Eversbott.

729. Rough Moonwort. — *O. Spicant.*Leaves lance-shaped, winged, segments run-
ning into each other, very entire, parallel. —

Curtis. II. 67. Sowerby. 1159.

Heaths and Marshes, July, P. Rare.

Potton Marshes.

Obs :

Barren leaves broader, fertile fronds narrowest.

Brakes. PTERIS. Gen. Pl. 1626.

Capsules disposed in a line on the under side
along the margin of the frond.730. Common Fern. — *P. aquilina.*Leaves more than double compound, leaflets
winged, wings spear-shaped, lower ones
with winged clefts, upper ones smaller. —
Bolton. 10. Sowerby. 1679.

Heaths, Commons, August, P. Common.

Spleen-wort.

XXIV. Ferns. Crypt. Filices.

Spleenwort. *ASPLENIUM.* Gen. Pl. 1631.

Capsules disposed in straight lines on the under side of the frond.

731. *Hart's-tongue.* — A. *Scolopendrium.*

Leaves undivided, heart-shaped, very entire,

— stalks hairy. — Curtis. I. 67. *Sowerby*. 1150.

Old Walls and Wells, August, P. ~~Rare~~ Common —
Bromham, — Eversholt. *Oakley - Puddington*.

732. *English Maiden-hair.* — A. *Trichomanes.*

Leaves winged, wings nearly round, scollopéd. — Woodville. 204. *Sowerby*. 576.

Old Walls, May, P. Rare.

Stafford Bridge, Oakley.

733. *Wall Rue.* — A. *Ruta muraria.*

Leaves doubly compound, in alternate order, leaflets wedge-shaped, scolloped. — Flora Danica. 190. *Sowerby*. 150.

Old Walls, June, P. Common.

734. *Black Maiden-hair.* — A. *Adiantum nigrum.*

Leaves almost trebly winged, leaflets alternate, wings spear-shaped, between cut and serrated. — Blackwell. 220.

Old Walls, April, P. Rare.

Stafford Bridge, Oakley.

G g

Polypody.

XXIV. Ferns. Crypt. Filices.

Polypody. POLYPODIUM. Gen. Pl. 1632.

Capsules disposed in roundish dots over the under surface of the frond. *to this tabn.*

735. *Common Polypody.* — P. vulgare.

Fronds with winged divisions, wings oblong, somewhat serrated, blunt, root scaly. —

Curtis. I. 68. Woodville. 271. Sowerby 1149.

Old Walls, Roots of Trees, June, P. Common.

Obs:

Fructifications in 2 rows, parallel.

736. *Crested Polypody.* — P. cristatum.

Fronds almost doubly winged, leaflets ovate-oblong, wings bluish, sharply serrated at the point. — Bolton. 23. Sowerby 11461.

Moist Woods and Bogs, June, P. Rare.

Potton Marshes, — Aspley Wood.

Obs:

Leaves paler than in the other species of this genus.

737. *Male Polypody.* — P. Filix mas.

Fronds doubly winged, wings blunt, finely scolloped, stalk chaffy. — Woodville. 49. Sowerby 1458.

Woods, June, P. Common.

738. *Female Polypody.* — P. Filix fæmina.

Frond doubly winged, smaller wings spear-shaped, with winged divisions, pointed. — Blackwell. 325. Sowerby 1459.

Moist

XXIV. Ferns. Crypt. Filices.

Moist Places, June, P. Rare.

*Potton Marshes.**Obs :*

Plant extremely elegant, divisions very fine, fructifications ovate, somewhat fringed.

739. *Marsh Polypody.* — P. *Thelypteris.*

Frond doubly winged, wings with other winged divisions, very entire, quite covered with impregnating dust on the under side. —

Bolton. 43. Sowerby. 1010.

Marshes, July, P. Rare.

*Potton Marshes.**Obs :*

The outline of the whole frond, taken collectively, is spear-shaped.

740. *Prickly Polypody.* — P. *aculeatum.*

Fronds doubly winged, wings spear-shaped, with fringe-like teeth, stalk chaffy. —

Bolton. XXVI. 1. and 3. Sowerby. 1562.

Springs in the Shade, June, P. Rare.

Eversholt.

Obs :

The lobe of that part of the leaf, which runs along the midrib, is shaped like an ear, and terminated by an awn.

G g 2

Club-moss.

XXIV. Mosses. Crypt. Musci.

Club-moss. Lycopodium. Gen. Pl. 1615.

Fruit disposed in the bosom of the scales, which (being tiled closely) form a spike, sometimes sitting among the leaves.

741. Common Wolf's-claw. — *L. clavatum.*

Leaves scattered, ending in threads, spikes round, in pairs, on fruitstalks. — Dillenius.

LXVIII. i. Sowerby. 224.

Heaths, July, P. Rare.

Potton Heath.

Obs :

Shoots creeping to a great extent at the roots of heath; fruit-stalks and spikes paler than the leaves and shoots.

742. Marsh Club-moss. — *L. inundatum.*

Leaves scattered, very entire, spikes terminating, leafy. — Dillenius. LXII. 7.

Bogs, June, P. Rare.

Amptbill.

Bog-moss. SPHAGNUM. Gen. Pl. 1637.

Capsule with a lid. Mouth smooth. Veil none.

743. Grey Bog-moss. — *S. palustre.*

Branches bent downwards. — Dillenius.

XXXII. i. Sowerby. 1405.

Bogs, July, P. Common.

Tree

* 745

Silver Earth-moss. *P. argenteum.*

Fertile stems dwarf, barren ones longer. Leaves egg-shaped, concave, con-
cavement forming cylindrical shoots,
silvery, ending in a fine hair.

Enclosure Banks. Jan'y Rare
Caldwell Jan'y 1800.

* 745 Thread moss Earth-moss.

P. bryoides,

Capsules exserted, upright, oval, beaked, leaves pointed somewhat erect, stems nearly simple — Dickson. Fasc. 4.

Tab. X. f. 3.

Wet places, on the edge of Stones
Marsh. Rare.

Clapham Springs —

*746. Creeping Earth-moss. *P. Holocistifera*.
Shoots thread-shaped, branched, geniculated, leaves which surround the capsule spear-shaped, toothed. *Dickson.*

Dickson. tab. VII. f. 2.

Play banks. Dec? rare
Caldwell Inclosures.

Obs.
like the *P. serratum*, but the capsule twice as large, root creeping by suckers, no herbage distinct from the shoots which bear the capsules.

*746 Serrated Earth-moss. *P. serratum*.

Shoots capillary, jointed, branched, capsule surrounded by lanceolate serrated leaves, rising above it.

Sowerby. 460. *English Botany.*

Banks. December. Rare.

Caldwell inclosures. Rec'd. J. H.

Obs.
Capsule a bright red, in which circumstance alone it differs from the figure above cited.

226

XXIV. Mosses. Crypt. Musci.

744. Tree Sphagnum. — *S. arboreum*.

Branched, creeping, capsules on the sides of the branches, flowering all one way. — *Dillenius.* XXXII. 6. *Sowerby.* 1180.

Trees, March, P. Rare.

Clapham Park Wood.

Obs:

This very rare plant grows imbedded in other mosses.

Earth-moss. *PHASCUM.* Gen. Pl. 1636.

Veil very small. Lid none.

745. Pointed Earth-moss. — *P. cuspidatum*.

On a stem, leaves ovate, pointed, open, uppermost erect, converging. — *Curtis.* IV. 66.

Pastures, January, A. Common.

746. Awl-leaved Earth-moss. — *P. subulatum*.

Stemless, capsule sitting, leaves awl-bristle-shaped, open. — *Curtis.* IV. 67.

Heaths, March, A. Common.

747. Alternate-leaved Earth-moss. — *P. alternifolium*. S.D.

Fertile stems dwarf, barren ones taller, upright, leaves alternate, awl-shaped. — *Dickson.* Fasc. II. tab. II. fig. 1.

Bogs, April, A. Rare.

Stevington Bogs.

Crooked

XXIV. Mosses. Crypt. Musc.

sent 748. Crooked Earth-moss. — P. curvicollum. S.D.

Sowerby. 905. Stemless, capsules on fruitstalks, bowed downwards, leaves spear-shaped, pointed, open. — (Dickson.) Dickson. I. 3.
Dec 23. 1880. Banks, April, A. Rare.

Caldwell Inclosures. ~~Bryum~~ Bryum lanceolatum.

749. Hairy Phascum. — P. piliferum. S.D.

On a stem, capsules on fruitstalks, leaves oblong, erect, with hairs at the ends. —

Schreber. I. 6. 7.

Banks, Hillocks, March, P. Rare.

Clapham Hill.

Obs.:

Capsule smaller than usual in the genus. Root-leaves fresh green, under the capsules somewhat paler, ending in a hair, sometimes bent downwards, often erect.

750. Round-capsuled Earth-moss. — P. sphæ-rocarpon. Sowerby. 610.*

Stemless, capsules sitting in the center of a rose of leaves; leaves strap-spear-shaped, spit-pointed.

Cart-ruts in Woods, October, A. Rare.

Clapham Park Wood.

Water-moss. FONTINALIS. Gen. Pl. 1655.

Capsule sitting on a short bristle, oblong, surrounded by a tiled fence. Fringes two, outer one with 16 broadish teeth, inner reticulated.

Great

748. Obs:

Capsule reddish brown, globular fruitstalk incurved, a little longer than the spear-shaped leaves.

*748. curled clustered Earth-moss. — P. crispum.

on a stem, leaves minute, scattered, floral leaves in clusters, elongated, pointed, incurved, entire — ~~Dickson. 610.~~
Sowerby. 1680. ~~excellens~~

Molehills. March. P. Rare.

Clapham Pastures.

*749 Wurfs Earth-moss. P. muticum

Stemless: leaves egg-shaped, concave, converging, not ending in hairs. soothening.

Dillenius XXXII. 12.

Garden Walks Autumn - W. Common.

Obs.:

Plant more diminutive than the cuspidatum, leaves not so deep a green. They want also the terminal hair, the true diagnostic of h? 745 —

*749 Many-capsuled Earth-moss. P. multicapitulare.

On a stem, leaves spear-shaped, concave, expanding, not contracted in a green state, capsules round, many on one shoot — Sowerby. 610. excellens

Molehills and arable Land.
Winter and Spring, Common.

XXIV. Mosses. Crypt. Musci.

751. Great Water-moss. — *F. antipyretica.*

Capsules nearly cylindrical, from the sides of the branches, leaves spear-shaped, loosely tiled, shoot branched, straggling. — Sowerby. 359.

Rivulets, June, P. Common.

Obs:

Leaves in three rows, fringes red, veil pointed, lid conical.

752. Least Water-moss. — *F. minor.*

Capsules terminating, leaves ovate, pointed, concave, pointing 3 ways; ~~away~~.

~~D.~~ — Dillenius. XXXIII. 2. Sowerby. 557.

Mill Walls, June, P. Rare.

Duck and Oakley Mills.

Obs:

Leaves more closely tiled than in the preceding species.

Bottle-moss. *SPLACHNUM.* Gen. Pl.

Capsule sitting on a large colored tubercle.

Veil falling off. Star of the female on a distinct plant.

753. Purple Bottle-moss. — *S. ampullaceum.*

Receptacle cruet-shaped, inversely ovate. — Sowerby. 144.

Bogs, March, A. Rare.

Ampbill.

Hair-moss.

XXIV. Mosses. Crypt. Musci.

Hair-moss. POLYTRICHUM. Gen. Pl.

Capsule with a lid, sitting on a very minute tubercle. Veil woolly. Star on a distinct plant.

754. *Common Hair-moss.* — P. commune.

Capsules growing in parallel lines from the base, veils woolly, leaves strap-shaped, finely scolloped. — Dillenius. LIV. 1.

Moors, May, P. Common.

755. *Juniper Hair-moss.* — P. juniperinum.

Capsules growing in parallel lines from the base, veils woolly, leaves strap-spear-shaped, very entire, pointed. — Dillenius. LIV. 2.

Hills, June, P. Common.

756. *Dwarf Hair-moss.* — P. subrotundum.

Capsules roundish, nodding, veils woolly, leaves indistinctly serrated. — Curtis. II. 68.

Heaths, May, P. Common.

Marsh-moss. MNIUM. Gen. Pl. 1593.

Capsule with a lid, veil smooth. Barren flowers in a small head, naked, powdery, remote.

757. *Fountain Marsh-moss.* — M. fontanum.

Capsules roundish, lid short, conical, leaves spear-serrated, tiled, bent inwards at the points. — Sowerby. 390.

Turfy

XXIV. Mosses. Crypt. Musci.

757. *Turfy Bogs*, May. P. Rare.

Amphibill. — *Sowerby*.

Obs:

Capsule supported by a fruitstalk, somewhat twisted, growing out of the fork of the shoot. Tuberle more splendid than usual in the genus.

758. *Common Marsh-moss*. — *M. palustre*.

Stem forked, leaves awl-shaped. — *Sowerby*. 391.

Bogs, June, P. Rare.

Amphibill, — *Potton*, — *Aspley*.

759. *Yellow Marsh-moss*. — *M. hygrometricum*.

— Stemless, capsule nodding, veil 4-square, bent back. — *Dillenius*. LII. 75. *Sowerby*. 342.

Walls, Banks, March, A. Common.

Obs:

Leaves rolled inwards, transparent, concave: fruitstalk bent inwards, yellowish green; veil slanting.

760. *Purple Marsh-moss*. — *M. purpureum*.

Stem forked, fruitstalks from the bosoms of the leaves, capsule erect, leaves keeled. — *Dillenius*. XLIX. 51.

Walls, April, P. Common.

761. *Swan's-neck Marsh-moss*. — *M. hornum*.

H h

Capsules

XXIV. Mosses. Crypt. Musci.

Capsules oblong, nodding, fruitstalk bowed down, leaves oblong-spear-shaped, slightly serrated, shoot nearly simple. — Dillenius.

LI. 71.

Bogs, February, P. Rare.

Potton, — Amptbill.

762. *Capillary Marsh-moss.* — *M. capillare.*

Capsules hanging down, leaves ovate, keeled, with bristles at the end, fruitstalks very long.

Walls, April, P. Rare.

Amptbill.

763. *Dotted Marsh-moss.* — *M. punctatum.*

Capsules ovate, hanging down, leaves inversely ovate, blunt, dotted, very entire. — Dillenius. LIII. 81.

Groves in a sandy soil, April, P. Rare.

Amptbill, — Southill.

Obs :

Shoots creeping, leaves roundish, alternate.

764. *Pointed Marsh-moss.* — *M. cuspidatum.*

Capsules ovate, hanging down, leaves sharp-ovate, pointed, slightly serrated. — Dillenius. LIII. 79. Sowerby. 1474.

Groves in a sandy soil, April, P. Common.

Obs :

Leaves keeled, fringed. Shoots rather upright.

Proliferous

Annual Marsh-moss. *M. annotinum.*

Leaves eggshaped, tapering, pellucid, fruitstalks from near the base of the shoot, capsules pendulous.

Dillenius I. 68.

Garden Walks, and wet places
P. March. June — Rare.

In Cuthbert's Bedford —
Road to Milton Grays.

Obs:

Younger shoots, barren, bulbiferous.
Bulbs red, roundish, pellucid,
separately arranged, sitting in
the wings of the leaves.

Leers.

XXIV. Mosses. Crypt. Musci.

765. *Proliferous Marsh-moss*.—*M. proliferum*.

Capsules ovate, hanging down, leaves crowded, in a rose, spear-shaped, pointed. — Dillenius. LII. 77.

Woods, April, P. Common.

Obs :

Shoots upright, branched, leaves keeled.

766. *Waved Marsh-moss*. — *M. undulatum*.

Capsules ovate-oblong, hanging down, leaves oblong, waved, slightly serrated. — Dillenius. LII. 76. Sowerby. 1449.

Woods, April, P. Common.

Obs :

Shoots very much branched, resembling a small tree, leaves waved.

767. *Triangular Marsh-moss*.—*M. triquetrum*.

Leaves in three rows, open, awl-spear-shaped, keeled. — Dillenius. LI. 72.

Bogs, April, P. Common.

768. *Turfy Marsh-moss*. — *M. cæspiticium*.

Capsules hanging down, leaves spear-shaped, between pointed and ending in a bristle, fruitstalks very long. — Curtis. III. 67. 1.

Walls, February, P. Common.

Obs :

The lower half of the fruitstalks red, the upper yellowish-green.

H h 2

Thread-moss.

XXIV. Mosses. Crypt. Musci.

Thread-moss. BRYUM. Gen. Pl. 1194.

Capsule with a lid. Veil smooth. Thread arising from a tubercle at the end of the shoot.

769. *Sessile Thread-moss.* — B. apocarpum.

Capsules sitting, at the ends of the branches, veil very small. — Dillenius. XXXII. 4. Flora Danica. 480. Sowerby. 1134.

On Tiles in a north aspect, November, P. Rare.
Amphill, Clapham, St. Mary's Church, Bedford.

Obs:

This species is soon determined by the extreme smallness of the veil and the scarlet color of the lid.

770. *Scored Thread-moss.* — B. striatum.

Capsules ovate-cylindrical, scored, leaves spear-shaped, margin bent back, open, upper ones gnawed at the point. — (Sibthorpe.) Dillenius. LV. 8.

Trees, February, P. Common.


Obs:

Capsules sitting, shoots small, much branched.

771. *Curled Thread-moss.* — B. crispum.

Capsules ovate, on fruitstalks thickening upwards, leaves strap-spear-shaped, curled up in drying. — (Sibthorpe.) Dillenius. LV. 11. Sowerby. 996.

Trees,



XXIV. Mosses. Crypt. Musci.

Trees, May, P. Common.

Obs :

Leaves more brilliant than in the preceding.

772. Bearded Thread-moss. — B. barbatum. S.D. sent

Capsules upright, oblong, bearded, lid pointed, slanting, fruitstalks from the sides of the shoots. — Dillenius. XLVIII. 48.

Walls, February, P. Common.

773. Shining Thread-moss. — B. pellucidum. S.D. sent

Capsules egg-oblong, small; lid taper-pointed, bent, shoot hairy, leaves spear-shaped, reflected, acute. — (Wilbering.) Dillenius. XLVI. 23. 24. Sowerby. 1020--.

Marshy spots, June, P. Rare.

Bog in Aspley Wood.

Obs :

Roots naked and covered with rust-colored capillary fibres.

774. Small Heath Thread-moss. — B. cerviculatum.

Capsules egg-shaped, unequal, toothed, leaves very slender. Sowerby. 1661.

Heaths, August, P. Rare.

Ampbill.

775. Pea-green Thread-moss. — B. convolutum. S.D. sent

Capsules

* 772 Beardless Thread-moss. B. imberbe.
 capsules oblong, beak slanting, leaves slender-spear-shaped, keeled, bent outwards.

Dillenius XLVIII. 46.

Ditch banks and mud walls. January, Rare.

Wall opposite the Limekiln St. Lloyd's
Bedford.

^{Obs.}
 Shoots more branched than in the barbatum, and of a deeper green. Lid very sharply pointed, veil placed on obliquely not occupying more than one third of the capsule —

Bryum infuscans
 between Milton & Radwell. 1799
 Pickering's Closes 1801. —

775. ^{Obs.}

Capsules in the younger specimens watery white, almost transparent. Fruitstalks rather spiral, when they first appear.

XXIV. Mosses. Crypt. Musci.

Capsules cylindrical, leaves of the fence blunt, rolled up so as to form a cylinder, leaves spear-shaped. — Dillenius. XLVIII.

44.

Base of brick walls, March, A. Common.

776. *Tendril Thread-moss.* — *B. cirratum.*

Capsules egg-oblong, lid taper-pointed, veil reaching but half way, shoots branched, leaves strap-spear-shaped, midribbed, acute. — (Withering.) *Flora Danica.* 538. 4.

Boggy woods, March, P. Rare.

Aspley Wood.

777. *Trifid Thread-moss.* — *B. trifarium.* S.D.

Capsules egg-oblong, mouth bearded, shoots seldom branched, leaves spear-awl-shaped, expanding. (Dickson.)

Stones, March, P. Rare.

Clapham Springs.

778. *Pale Thread-moss.* — *B. androgynum.*

Capsules cylindrical, lids taper-pointed, fruit-stalks terminating, shoot branched, upright, leaves spear-shaped, rather expanding. — Dillenius. XXXI. 1.

Woods, June, A. Rare.

Aspley Wood.

Simple

775.

Elstow Churchyard Wall
aspect N.E. facing the stile as
you enter from the Swan.
Decr. 2nd 1790.

*775

Small green Thread-moss. *B. viridulum.*

~~Capsules egg-shaped, leaves spear-shaped
taper-pointed, tilted, but expanding.~~

~~Dillenius XL. VIII. 43.~~

~~Mud walls, December, Rare
walls of Oakley Churchyard.~~

~~Fruit stalk generally single, some-
times two from the same shoot.~~

777. *Obs.*

leaves dark green, fruit stalks
crimson, base of the tubercle pale
yellow, upper part crimson.

XXIV. Mosses. Crypt. Musci.

779. *Simple Thread-moss*. — *B. simplex*.

Capsules ovate-oblong, nodding, lid conical,
leaves awl-shaped, scattered. — (*Sibthorpe.*)
Dillenius. L. 59. *Sowerby*. 1215.

Marshy spots, March, P. Rare.

Clapham Springs.

780. *Seagreen Thread-moss*. — *B. glaucum*.

Capsules oblong, lid pointed, bent back,
leaves erect, tiled, stem branched, upright. —
Dillenius. XLVI. 20.

Heaths and woods, October, P. Rare.

Ampthill Heath, — *Aspley Wood*.

781. *Equal Thread-moss*. — *B. decipiens*. S. D. *but*

Capsules slanting, lid conical, fruitstalks as
high as the leaves, leaves strap-spear-shaped,
keeled.

Walls, February, P. Rare.

Clapham, — *Ampthill*.

Obs:

Like the barbatum, but smaller in all its parts.

782. *Extinguisher Thread-moss*. — *B. extincto-*
rium.

Capsule erect, oblong, smaller than the veil ;
veils loose, equal at the base. — Dillenius.
XLV. 8. *Sowerby*. 550.

Walls, October, P. Common.

Spear-shaped

782. Obs:

Fruitstalks red towards the base
pale yellow under the capsule
which is bright green and conicab.

XXIV. Mosses. Crypt. Musci.

783. Spear-shaped Thread-moss. — B. lanceolatum.

Capsule erect, ovate, toothed, shoots rather upright, leaves spear-shaped, concave, pointed. — (Dickson.)
Sowerby. 1408.

Grassy spots, March, P. Common.

784. Heath Thread-moss. — B. heteromallum.

Sowerby. 1272. Capsules erect, leaves bristle-shaped, pointing one way. — Dillenius. XLVII. 37.

Heaths, March, A. Common.

Obs:

Fruitstalks rising from the back of the shoots.

785. Wall Thread-moss. — B. murale.

Capsules upright, leaves with a hair at the end, straightish, shoots simple, in a turf. — Dillenius. XLV. 14.

Walls, March, P. Common.

786. Egg Thread-moss. — B. ovatum.

Capsules upright, ovate, leaves ovate, concave, with a hair at the end. — (Dickson.)

Mud walls, April, A. Common.

Obs:

Shoots tiled in the closest manner, with leaves that are not only concave, but converging.

787. Apple Thread-moss. — B. pomiforme.

Capsules erect, spherical. — Dillenius.
XLIV. 1. Sowerby. 990.

Sandy

* 783. Dickson's Thread-moss. B. Dicksonii.

Capsule erect, between ovate and cylindric, shoots branched, leaves strap-shaped, keeled, rather curled in drying.

Dickson. Fasc. 3. tab. 7. f. 7. Dickson.

Rotten wood & Trunks of Trees.
February. Rare.

Amphill. — .

785. Obs:

leaves oval, semi-convolute.
Fruit at first yellowish green, brown in decay.

XXIV. Mosses. Crypt. Musci.

Sandy Banks, February, A. Rare.

Clophill, — Southill, — Potton.

Obs:

Leaves growing in a thick turf, very finely scalloped towards the points. Capsule round, yellow-green.

788. Pear Thread-moss. — *B. pyriforme.*

Leaves expanded, flat, oblong, pointed, capsules shaped like a pear. — Sowerby. 413.

Banks, February, A. Common.

789. Stiff Thread-moss. — *B. rigidum.*

Capsules oblong, upright, shoots simple, very short, leaves oblong, stiff, rolled in at the edge. — Sowerby. 180. *spec.*

Walls, November, P. Common.

Obs:

Leaves curl up in drying, lid red at the base, fringe twisted up cylindrically.

790. Grey Thread-moss. — *B. rurale.*

Capsules nearly upright, leaves bent back, ending with a hair. — Dillenius. XLV. 12.

Roofs of Cottages, December, P. Common.

791. Broom Thread-moss. — *B. scoparium.*

Capsules nearly upright, fruitstalks incorporated, leaves pointing one way, bowed back, stem declining. — Dillenius. XLVI. 16. Sowerby. 354.

Heaths, June, P. Common.

789.
specimens in the Gratus ad Parn.
at Page. 394

XXIV. Mosses. Crypt. Musci.

792. Awl Thread-moss. — *B. subulatum*.

Capsules upright, awl-shaped, shoots stemless. — Curtis. III. 66. Sowerby. 1101.

Walls, Banks, January, P. Common.

Obs:*The size of the leaves particularizes the species at first sight.*793. Brown Thread-moss. — *B. truncatulum*.

Capsules upright, roundish, lid ending in a sharp point. — Curtis. II. 70. 2.

Ditches, February, A. Common.

Obs:*Capsule lopped when ripe.*794. Crowded Thread-moss. — *B. hypnoides*.Capsules egg-shaped, shoots ascending, branches lateral, short, fruit-bearing. — (With Sowerby. 1348. ^{erung.}) Dillenius. XLVII. 32.

Fir Groves, Autumn, Rare.

Warden.

795. Bird's-claw Thread-moss. — *B. unguiculatum*. Specimens in British 3rd vol.

Capsules upright, oblong, lid pointed, oblique, fruitstalks thickening upwards, leaves strap-spear-shaped, keeled. — Dillenius. XLVIII. 47.

Walls, March, A. Common.

Green

Red Thread-moss. *B. rufescens*.

Capsules egg-shaped, toothed, leaves strap-shaped, pellucid, flaccid, se: :cundorous — Dickson.

— " — Dickson. Fasc. III. VIII. 3.

Cart Ruts in wet Places A. March Rare.

Pickings by a double Hedge.
Between Milton Brays & Radwell.Obs.

Plant grows in patches, but is very sparing in the produce of capsules: I find the fruitstalk longer in proportion to the rest of the plant in the plate above quoted, and tho' I have perceived many specimens in fruit, could never discover any veil.

792. Calcareous Thread-moss. *B. fallax.*
Capsules oblong, mouth with a twisted
beard, shoots branched upwards, leaves
strap spear-shaped, bending outwards.

Hedwig. I. 24.

Barbula fallax.

Joints of Brickwork in damp places.
March. P. Rare.

Bromham House.—

793. Obs.:

Leaves altho' closely tiled, visibly
terminated each by a fine hair
towards the point of the shoot—

Obs.:

794. Leaves deep green, convolute.
whole plant a cushion shaped msp.
extremely soft, the fruit almost
buried in the herbage—

795. Silky Thread-moss. *B. sericeum.*

Capsules cylindrical, lids conical,
dagger-pointed, shoot simple, leaves
egg-spear-shaped, closing.

Dillenius L. 61.

Foggy places. P. Marsh July. Rare.
Aspley Wood.

797. Brittle Thread-moss. *B. fragile.*
Capsules upright, toothed, obovate, fruitstalk
zigzag, leaves between hair & strap-shaped
rather cleaning on one side. Dillenius XLVII. 33.
Dry Places. April. Rare.
Aspley Wood. 243 F.G.

XXIV. Mosses. Crypt. Musci.

796. Green Thread-moss. — *B. virens.*

Capsules upright, oblong, leaves awl-shaped,
quite straight. — Dillenius. XLVIII. 43.

Ditch Banks, March, A. Common.

797. Silver Thread-moss. — *B. argenteum.*

Capsules hanging down, shoots cylindrical,
tiled, smooth. — Curtis. III. 67. 2. Sowerby. 1602.
Walls, Banks, February, P. Common.

798. Cushion Thread-moss. — *B. pulvinatum.*

Capsules roundish, fruitstalks bowed back,
leaves ending in hairs. — Dillenius. L. 65.

Walls, March, A. Common.

799. Curled Thread-moss. — *B. undulatum.*

March

Capsule cylindrical, nodding, veils smooth,
leaves spear-shaped, waved, serrated.—Curtis. I. 70.

Woods, February, P. Common.

800. Chalky Thread-moss. — *B. calcareum.*

Capsules upright, inversely conical, toothed,
leaves upright, round, bluntnish.—Sowerby.

191. excellent.

Chalk Pits, May, A. Rare.

Barton Hill.

Obs.:

Fruitstalk scarcely taller than the leaves. Veil covering the
upper half of the capsule.

I i 2

Feather-moss

XXIV. Mosses. Crypt. Musci.

Feather-moss. HYPNUM. Gen. Pl. 1656.

Capsule with a lid; veil smooth, thread from the side of the stem, surrounded by an empalement.

801. *Upright Feather-moss.* — H. adiantoides.

Leaf winged, branched, upright, bearing fruitstalks in the middle. — Sowerby. 264.

Commons, April, P. Common.

Obs:

Leaf stiffish, leaflets alternate. X

802. *Flat Feather-moss.* — H. complanatum.

Leaf winged, branched, leaflets tiled, pointed, folded, compressed.—Dillenius. XXXIV. 7.

Sowerby. 1492
Trunks of Trees, March, P. Common.

Obs:

Fruitstalks rise sometimes from the midrib, often out of the fork.

803. *Yew-leaved Feather-moss.* — H. taxifolium.

Leaf not branched, winged, spear-shaped, bearing the fruitstalk at its base. — Dillenius. XXXIV. 2. Sowerby. 426.

Woods, February, A. Common.

804. *Small Feather-moss.* — H. bryoides.

Fruitstalks terminating, capsules oblong, lid acute, leaflets spear-shaped, sharp-pointed. — Dillenius. XXXIV. 1. Sowerby. 625.

Woods,

X leaves of the empalement about
4. strap - spear shaped, base of the
spiral fruit stalk a deep crimson.

*804. Two rowed Feather moss.

H. denticulatum.

Seldom branched, fruit stalks from near the base, capsules cylindrical, lid blunt, leaves wedge-shaped, pointed in 2 rows.

Dillenius XXXIV. 5.

Boggy Woods at the roots of Trees
P. March. April. Rare.

Aspley Wood.

Obs.:
Leaves in the alternate row
shorter.

Small Plants in P. Woods
Frustulatae by no means separate

245

XXIV. Mosses. Crypt. Musci.

Woods, May, A. Rare.

Aspley.

805. *Round-leaved Feather-moss.* — *H. rotundifolium.* S.D. sent

Branched, fruitstalks lateral, capsules ovate.
Mill wheels, April, P. Very rare elsewhere,
but common in Bedfordshire.

806. *Floating Feather-moss.* — *H. fluitans.* S.D.

Shoots straggling, branched, leaves spear-awl-shaped, scattered. — Dillenius. XXXVIII.

33. Sowerby 1448.

Stagnant Waters, September, A. Rare.

Ponds on the Goldington Road.

Obs.:

Where it grows it is found in profusion, but never fructifies here.

807. *Yellow Feather-moss.* — *H. lutescens.*

Shoots straggling, branched, leaves awl-shaped, capsules oblique. — Dillenius. XLII.

60. Sowerby 1301.

Chalky Hills, January, P. Common.

808. *Common Feather-moss.* — *H. rutabulum.* Sowerby. 1647.

Shoots straggling, branched, leaves ovate-pointed, open, capsules slanting, lid conical. — Dillenius. XXXVIII. 29.

Woods, September, P. Common.

Butcher's

XXIV. Mosses. Crypt. Musci.

- Sowerby. 1275.* 809. *Butcher's broom Feather-moss.* — *H. ruscifolium.*
Shoots creeping, branched, leaves oval, concave, transparent, capsules roundish, bent inwards. — Dillenius. XXXVIII. 31.
Mill Wheels, January, P. Rare.
- Oakley and Duck Mills.*
- Sibthorpe. 1302.* 810. *Starry Feather-moss.* — *H. stellatum.* S.D.
Shoots branched, starry at the points, leaves spear-shaped, with long points, open. — (Sibthorpe.) Dickson. I. 7. *Sowerby. 1b22.*
Turfy Bogs, April, P. Rare.
- Ampthill.*
811. *Triangular Feather-moss.* — *H. triquetrum.* *Sowerby. 1b22.*
Branches straggling, bent back, leaves ovate, bent back, open. — Dillenius. XXXVIII. 28.
Woods, September, P. Common.

*Obs:**Leaves not keeled.*

812. *Fir Feather-moss.* — *H. abietinum.*
Shoots winged, roundish, distant, unequal. — Dillenius. XXXV. 17.
Chalkpits, March, P. Rare.

*Barton Hills.**Obs:**I have not seen it in fructification.**Crested*

813. *H. molluscum*. Stem procumbent, pectinate,
leaves ovate and shaped, curved, without ribs, capsule
ovate, drooping, lid conical.
English Botany.

Bryum Dicksoni.

To Sowerby
Dr. Smith
Jan 26.
1801.

March 30 1799

No. 3
Oct 28th 1799

To Dickson
No. 1. Feb. 10
1801.

To Sowerby.

No. 4.
Feb 10. 1801.

247

XXIV. Mosses. Crypt. Musci.

813. Crested Feather-moss. — *H. exstacans*
~~erensis~~ *molluscum*. Sowerby. 1327.

Shoots winged, little branches close together,
with the points bowed back. — Dillenius.
XXXVI. 20.

Hills, March, P. Common.

Obs :

Plants often crowded into great masses, procumbent, soon
perceived by their silky gloss.

814. Bog Feather-moss. — *H. filicinum*. S. D.

Shoots winged, little branches distant, leaflets
tiled, bowed inwards, sharp, pointed one
way. — Dillenius. XXXVI. 19. Sowerby. 1570.

Bogs, March, P. Rare.

Stevington Bogs.

Obs :

Leaves branched, often proliferous at the sides.

815. Wall Feather-moss. — *H. parietinum*. Sowerby 1424.

Shoots flat and winged, lengthened out, fruit-
stalks in clusters. — Dillenius. XXXV. 13.

Heaths, February, P. Common.

Obs :

The shoots of the new year do not arise from the end of the
old one, but near to it.

816. Feathery Feather-moss. — *H. plumosum*.

Shoots

XXIV. Mosses. Crypt. Musci.

Shoots winged, creeping, branches crowded, leaves tiled, awl-shaped, capsules erect. — Dillenius. XXXV. 16.

Trees, January, P. Rare.

Sheerhatch Wood.

817. *Trailing Feather-moss.* — *H. praelongum.*

Shoots simply winged, leaves spear-shaped, loosely tiled, capsules ovate, lid like a reaping hook. — Dillenius. XXXV. 15.

Trees, February, P. Common.

818. *Proliferous Feather-moss.* — *H. proliferum.*

Sowerby. 1494 Shoots bearing others, flat, winged, fruit-stalks in clusters. — Dillenius. XXXV. 14.

Shady Woods, December, P. Common.

Obs :

Shoot produced from the end of the leaf.

819. *Hooked Feather-moss.* — *H. aduncum. S.D.*

Shoot branched, upright, leaves strap-spear-shaped, like a reaping hook, crowded, hooked at the end. — Dillenius. XXXVII. 26.

Marshes, April, P. Common.

820. *Cypress Feather-moss.* — *H. cupressiforme.*

Shoots somewhat winged, leaves pointing one way, bowed back, awl-shaped at the end. — Dillenius. XXXVII. 23.

Woods, March, P. Common.

Scorpion's

XXIV. Mosses. Crypt. Musci.

821. *Scorpion's-tail Feather-Moss.* — H. scorpioides. S.D.

Branches straggling, trailing, bowed back, leaves pointing one way, pointed. **Sowerby.** 1039.

Turfy Bogs, November, P. Rare.

Amptbill, — *Potton*.

822. *Scurfy Feather-moss.* — H. squarrosum.

Shoots branched, leaves awl-shaped, bowed back in all directions, capsules oblong, rather nodding, lid short, conical. — Dillenius. XXXIX. 38.

Moist Meadows, March, P. Common.

823. *Twig Feather-moss.* — H. viticulosum. S.D.

Shoots creeping, branches straggling, round, leaves open, taper-pointed. — Dillenius. XXXIX. 43. **Sowerby.** 265.

Trees, March, P. Common.

Obs :

Lid pointed, capsule mahogany-colored.

824. *Fox-tail Feather-moss.* — H. alopecurum.

Shoots upright, branches in bundles, terminating, subdivided, capsules rather nodding. Dillenius. XLI. 49. **Sowerby.** 1182.

K k

Woods,

XXIV. Mosses. Crypt. Musci.

Woods, April, P. Common.

*Obs:**Stem scales tapering to a point.*825. *Tree Feather-moss.* — *H. dendroides.*

Shoot upright, branches in bundles, terminating, mostly simple, capsules upright. — Dillenius. XL. 48. *Sowerby*. 1565.

Woods, February, P. Rare.

*Amphill, — Southill, — Clophill.**Obs:**Stem scales blunish.*826. *Pointed Feather-moss.* — *H. cuspidatum.*

Shoots straggling, pointed at the ends by the edges of the leaves rolled inwards. — Dillenius. XXXIX. 34. *Sowerby*. 1425.

Wet Pastures, March, P. Common.

827. *Meadow Feather-moss.* — *H. purum.*

Shoots winged, scattered, awl-shaped, leaves ovate, blunt, approaching. — Curtis. III. 65.

Pastures, February, P. Common.

828. *Water Feather-moss.* — *H. riparium.*

Shoots cylindrical, branched, leaflets pointed, open, distant. — Dillenius. XL. 44.

Bogs, April, P. Common.

Squirrel's-tail

#830 - Bob-tail Feather-moss.

H. curtipedunculum.

Shoots upright, branched, leafy.
leaves oblong egg-shaped, acute, tilted.
capsules oval, lid tapering.

Dillenius XLIII. 69.

Beech tree Bark. Feby Rare.

Milton Hill. Coburn Park.

Obs: Feb. 7 1799.

Fructification axillary, leaves
of the fence very long, fruit:
stalk very short. — .

XXIV. Mosses. Crypt. Musci.

829. Squirrel's-tail Feather-moss.—*H. sciuroides.*

Shoots upright, branched, bowed inwards. —
Dillenius. XLI. 54.

Trees, February, P. Common.

830. Mouse-tail Feather-moss.—*H. myosuroides.*

Shoots very much branched, branches awl-shaped, round, tapering each way. — Dillenius. XLI. 51. Sowerby. 1567.

Trees and Pales, March, P. Common.

831. Silky Feather-moss.—*H. sericeum.*

Shoot creeping, branches crowded, upright,
leaves awl-shaped, capsules upright. —
Dillenius. XLII. 59. Sowerby. 1445.

Walls, Trees, April, P. Common.

832. Creeping Feather-moss.—*H. serpens.*

Shoots creeping, branches thread-shaped,
leaves very minute. — Dillenius. XLII. 64.

Trees, March, P. Common. Sowerby. 1037 —

833. Velvet Feather-moss.—*H. velutinum.*

Shoot creeping, branches crowded, upright,
leaves awl-shaped, capsules somewhat nodding. — Dillenius. XLII. 61. Sowerby. 1568 ..

Trees, Autumn, P. Common.

XXIV. Thongs. Crypt. Algae.

Star-tip. JUNGERMANIA. Gen. Pl. 1662.

Capsule solitary, with 4 valves. Seeds attached to a thread.

834. *Spleenwort Star-tip.* — *J. asplenoides.*

Sowerby. 1788. Leaves simply winged, leaflets ovate, somewhat fringed. — Dillenius. LXIX. 6.

Ditch Banks near Woods, April, P. Common.

835. *Forked Star-tip.* — *J. bicuspidata.*

Leaves simply winged, flowering in the middle, leaflets with two teeth. — Sowerby. 281.

Wet shady Places, March, P. Common.

Obs:

The two nicks in the leaves form an acute angle.

836. *Cloven Star-tip.* — *J. bidentata.*

Sowerby. 606. Leaves simply winged, flowering at the point, leaflets with two teeth. — Dillenius. LXX. 11.

Wet shady places, October, A. Common.

Obs:

The two nicks of the leaves resemble the pointed horns of the new moon.

837. *Straggling Star-tip.* — *J. viticulosa.*

Leaves simply winged, leaflets roundish, very entire. — Dillenius. LXIX. 6.

Walls, April, P. Common.

Flat

In? Sowerby deserves any species of this genus.

* 835 Spear-shaped Star-tip. *J. lanceolata.*

Shoots spear-shaped, fruit stalks terminating, leaflets very entire.

Dillenius LXX. 10.

Shady Banks near Springs.
Rare.

Lversholt

Ayde hill near Luton.

Owing on the frond.—

v. 41. *Sowerby. 771.*

P. Rare.

Amphill, — Potton.

Obs:

not growing to the soil.

Strap-shaped

XXIV. Thongs. Crypt. Algæ.

§38. Flat Star-tip. — *J. complanata.*

Shoots creeping, leaflets doubly tiled, with ear-shaped segments underneath, branches of equal breadth. — Curtis. IV. 68. Dillenius. LXXII. 26.

Trees, January, P. Common.

§39. Scaly Star-tip. — *J. dilatata.*

Shoots creeping, leaflets doubly tiled, with ear-shaped segments underneath, branches broader at the point. — Dillenius. LXXII. 27.

Tress, March, P. Common.

§40. Tamarisk Star-tip. — *J. tamariscifolia.*

Leaves tiled in a double row, upper ones roundish, convex, blunt, four times as large as the other. — Dillenius. LXXII. 31. Sowerby. 108b.

Trees, February, P. Common.

Obs :

Fructifications yellow, sometimes erect, oftener drooping.

§41. Broad-leaved Star-tip. — *J. epiphylla.*

Stemless, a little leaf growing on the frond. —

Dillenius. LXXIV. 41. Sowerby. 771.

Bogs, March, P. Rare.

Ampthill, — Potton.

Obs :

Lower leaf growing to the soil.

Strap-shaped

*840. Fern-like Star-tips *J. Tomentella*

Shoots creeping, downy on the upper surface, leaves doubly winged, very downy Dillenius LXXIII. 35.

Springs & damp Woods. April, Rare Clapham Springs.

XXIV. Thongs. Crypt. Algae.

842. Strap-shaped Star-tip. — *J. furcata*.

Stemless, frond strap-shaped, branched, the ends forked and bluntish. — Dillenius. LXXIV. 45. Sowerby 1632.

Trees, April, P. Rare. ^{in Amptbill-Park Plantations.}

Amptbill-Park Plantations.

Obs :

Small branches adhering firmly to the bark of trees, strap-shaped, forked.

843. Wall Star-tip. — *J. platyphylla*.

Sowerby 798. Shoots trailing, tiled underneath, leaflets heart-shaped, pointed. — Dillenius. LXXII. 32.

Walls, March, P. Common.

Obs :

Spreading like a carpet over the brink of old walls.

844. Dwarf Star-tip. — *J. multifida*.

Stemless, shoots with double wings. — Sowerby. 186. excellent.

Woods, April, P. Common.

Obs :

Leaves very finely divided, fructifications rising nearly from the center of the leaves, yellowish, white at the base, numerous.

845. Jagged Star-tip. — *J. pinguis*.

Stemless, frond oblong, waved, greasy. — Sowerby. 185.

Bogs, April, P. Common.

Obs :

* 843.

Pale Star-tip. — *J. cinerea*.

Shoots creeping, doubly winged above, fruit stalk central, sheath round leaves blunt.

Dillenius LXXII. 28.

On damp sticks and mosses in wood winter & spring

Clapham between Woods. rare.

XXIV. Thongs. Crypt. Algæ.

Obs:

Leaves broader than in the other star-tips, paler towards the ends.

§46. *Shining Star-tip.* — *J. pusilla.*

Stemless, frond mostly winged, lobes tiled, empalement plaited. — Dillenius. LXXIV.

46.

Shady Lanes, March, P. Rare.

Hazel-wood Lane.

§47. *Powdered Star-tip.* — *J. fissa.*

Fronds simply winged, leaflets in two rows, nicked at the point. — Dillenius. XXXI. 6.

Wet Places, March, P. Rare.

Ampthill Moor.

Obs:

Fronds in two rows, nicked, small branches few, 2 or 3 only, so that they can hardly be called winged.

Liver-green. MARCHANTIA. Gen. Pl. 1661.

Receptacle target-shaped, bearing seeds underneath, in many capsules. Capsules with 4 valves. Seeds attached to a thread.

§48. *Star-headed Liver-green.* — *M. polymorpha.*

Common cup with 10 divisions. — Dillenius. LXXVI. 6. Sowerby. 210.

*. *M. conica.* — *Conical Liver-green.* — Wet Cup of the female flowers ovate, pointed with 5 marginal notches; male flowers in separate warts.

Sowerby. 504.

Shady rills Spring Rare Clapham Between woods.

* 848

M. conica

Sowerby 504.

XXIV. Thongs. Crypt. Algae.

Wet Places, July, P. Rare.

Eversb.^{olt.}

Obs:

Heads shaped like the skeleton of an umbrella.

Liver-wort. LICHEN. Gen. Pl.

Male. — Receptacle roundish, rather flat, shining.

Female. — A meal sprinkled on the leaves.

* Powdery Lichens.

849. *White Liver-wort.* — L. albus.

Very white, between powdery and crustaceous. — Dillenius. I. 2, Flora Danica.

840. 4. Sowerby. 1349.

On decayed Mosses, January, P. Common.

850. *Hoary Liver-wort.* — L. incanus.

Hoary, powdery, like scattered meal. — Dillenius. I. 3.

Bark of Trees, June, A. Common.

851. *Yellow Liver-wort.* — L. flavus.

Yellow, powdery, growing on wood. — Dillenius. I. 4. Sowerby. 1350.

Old pales, Oak bark, September, A. Common.

* * Crustaceous,

849. *White Liver-wort.* — L. albus.

850. *Hoary Liver-wort.* — L. incanus.

851. *Yellow Liver-wort.* — L. flavus.

852. *Crustaceous Liver-wort.* — L. crustaceum.

853. *Dark Liver-wort.* — L. nigra.

854. *Crustaceous Liver-wort.* — L. crustaceum.

855. *Crustaceous Liver-wort.* — L. crustaceum.

856. *Crustaceous Liver-wort.* — L. crustaceum.

857. *Crustaceous Liver-wort.* — L. crustaceum.

858. *Crustaceous Liver-wort.* — L. crustaceum.

859. *Crustaceous Liver-wort.* — L. crustaceum.

860. *Crustaceous Liver-wort.* — L. crustaceum.

XXIV. Thongs. Crypt. Algæ.

* * * Crustaceous, with black lines.

852. Wrinkled Liver-wort. — *L. rugosus.*

Crust whitish, simple, black lines and dots crowded together. — Dillenius. XVIII. 2.

Bark of Trees. January, P. Common.

Obs :

Scarcely different from *L. scriptus*.

853. Lettered Liver-wort. — *L. scriptus.*

Crust whitish, with black lines branched like Asiatic characters. — Dillenius. XVIII. 1.

Bark of Trees, January, P. Common.

854. Flea Liver-wort. — *L. pulicaris.*

Crust whitish, lines black, very short and simple. — Hoffman's Enum. Lich. III. 2. E.

Bark of Trees, December, P. Rare.

Amphill Park.

* * * Crustaceous, with tubercles.

855. Beech Liver-wort. — *L. fagineus.*

Crust white, tubercles white, mealy. — Dillenius. XVIII. 11.

Bark of Trees, December, P. Common.

856. Whitish-black Liver-wort. — *L. atro-albus.*

Crust black, tubercles black and white, mixed together. — Jacquin. Collect. II. 14. 1.

L 1

Walls,

x 853. Hebrew Liverwort *L. hebraica*.

Leprous, whitish, fructifications consisting of small black lines diverging.

Hoffmanni Num. Lich. III. 2. F.

Bark of Trees, December P. Rare.
Clapham Park Wood.

London Museum. January P. Common.

* 853

L. inquinans.

English Botany —

Dillenius. L. 14. 1. 1753
old names. Collected September A. Cam-

Commonwealth

XXIV. Thongs. Crypt. Algæ.

Walls, January, P. Common.

857. *Dyer's Liver-wort.* — *L. calcareus.*
Crust white, tubercles black. — Dilleniæ.
XVIII. 8.

Walls, January, P. Common.

858. *Sunk Liver-wort.* — *L. immersus.*
Tubercles black, sunk in the stones. — Sowerby. 193. excellent.
Chalk or Limestone, January, P. Rare.

Barton, — Elstow.

Obs :

Tubercles black, very minute, color of the crust very white, substance very thin.

859. *Black-knobbed Liver-wort.* — *L. sanguinarius.*
Crust ashy-green, tubercles black. — Sowerby. 155. excellent.

Trees, January, P. Common.

860. *Moss Liver-wort.* — *L. muscorum.*
Crust ash-colored, tubercles black. — Relhan's plate at page 424. Sowerby. 626.
On Mosses, January, P. Rare.

Newenham Wall.

Obs :

The crust is a mass united and growing amongst various species of the *Hypna*, tubercles globular.

Hoary

which, as said before, containing two
other parts, the one, which, however,
parted largely — and Cetariae

- * 858. *Rock Liverwort.* *L. rupestris.*

Tubercles sitting, bare, dark brown,
Limestone walls. Winter, Rare.
Friars walls.

Obs :
Tubercles at first completely convex
in a dry and advanced state they become
flat, and have a cartilaginous border.

- * 859. *Granulated Liverwort.* *L. graniformis*

Tubercles black, crowded: crust
greenish grey, with granulations.
Hagen. 1. 2 ?

In the clefts of old willows. Common
all the year —

* 861 Grey and red Liverwort. *L. ciliaris*.

Crust granulated, lobed, pale grey, shields tawny-red, flattish, with a thick white undulated border. — English Botany.

Sowerby. 1040.

Bricks walls
861. in a sloping position.

* Confluent Liverwort. *L. confluens*.

Tuberules very black, distinct when young, confluent when old, crust blue-grey, pale brown, or white. — Withering

Hoffman Lich. XIX. 1.

Walls - Winter - Rare.

Garden walls, Bedford.

* 864. Mosaic Lichen. *L. tessellatus*.

Crust gray, cracked, whitish about the tuberules, which are immersed, black prominent, smooth. — English Botany.

Sowerby. 533.

Brick walls, December, Rare.
Bedford, my own garden.

259

XXIV. Thongs. Crypt. Algæ.

861. Hoary Liver-wort. — *L. canescens*.

Crust whitish, tubercles very black, crowded. Dillenius. XVIII. 17. A.

Trees, Walls, December, P. Common.

862. Bluish Liver-wort. — *L. cæruleo-nigricans*.

Crust bluish, tubercles black. — Hoffman, Enum. Lich. XXXII. 3. Sowerby. 1139 — .

Hills, January, P. Rare.

Barton Hill, — Kempston.

Obs:

Younger tubercles greenish, almost sitting, older ones on stems, at length turning back.

863. Wall Liver-wort. — *L. fusco-ater*.

Crust brown, tubercles black. Jacquin. Collect. II. 14. 3.

Walls, January, P. Common.

864. Oak Liver-wort. — *L. querneus*.

Crust yellowish brown, tubercles black, nearly sunk in the crust. — Dickson. Fasc. Crypt. I. 2. 3. Sowerby. 405.

Posts and Gates, January, P. Common.

865. Sulphureous Liver-wort. — *L. sulphureus*.

Crust brimstone-colored, cracked, uneven, tubercles brown-black, irregular, very small. Hoffman. Enum. Lich. VI. 3.

L 1 2

Brick

XXIV. Thongs. Crypt. Algæ.

Brick Walls, March, P. Rare.

Elstow.

866. Horn-beam Liver-wort. —
- L. carpineus.*

Crust ash-colored, tubercles whitish, wrinkled. — Dillenius. XVIII. II. C.

Trees, January, P. Common.

Obs:

Crust very thin.

867. Heath Liver-wort. —
- L. ericetorum.*

Crust the color of verdigris, tubercles nearly sitting, flat, flesh-colored. — Sowerby. 372. excellent.

Heaths, December, P. Common.

Obs:

At first sight like a slice of sage cheese.

868. Mushroom Liver-wort. —
- L. byssoides.*

Crust ashy-green, tubercles brown, mostly on fruitstalks. — Sowerby. 373.

Heaths, December, P. Rare.

Ampthill.

Obs:

Tubercles on long stems like a small species of Agaric.

869. Large Heath Liver-wort. —
- L. Baomyces.*

Crust whitish, tubercles on stems, flesh-colored. — Sowerby. 374. excellent.

Heaths,

#866 Quartz Liverwort. *L. dendriticus.*

Crust between ash-colored and black, border branching like a tree, shields convex, of the same colour — Hoffman —

Hoffman. Pl. Lich. XIX. 4.

Bricks in an Eastern aspect.
all the year - Rare -Friars Farm — St. John's Hospital walls
Bedford.

Obs:

This Liverwort when most perfect is the smallest of the tribe I ever yet met with, it is not therefore to be wondered that it should have so long escaped notice in Britain — we may however venture to pronounce that upon a minute and microscopic inspection it will amply repay the study of the scrutinizing Botanist — Crust rising, never, border foliaceous tubercles very prominent, darker than the crust.

* 879. to follow L. ater.

~~Bleak Liverwort~~ *L. fragilis.*

Expanded, soul-like, white. Shields flat,
Ochre: flesh colored, borders white.

Winter *Dickson.*

~~Bleak Liverworts~~ rare

~~Barton Hills.~~

* 879.2 Globular Lichen. *L. sphaeroides.*

Crust greenish grey, tubercles roundish
flesh: coloured. *Dickson.*

Dickson. June, Oct: Woods Common.

Fasc. I. tab. 2. f. 2.

871. a single elm tree on the
cross road at the entrance into Elstow.
also behind Kempston-Bury-house.

to follow Liverwort after

* 874 White Liverwort. *L. leptigerus.*

Crust whitish somewhat lobed. Shields
yellowish when full grown, white at the
edge *Rellian* at p. 430. *Withering*

Hoffman IX. 4.

Heaths and dry walls. *P. Janz rare*
Bedford old walls.

* 874. Chart Liverwort. *L. chartaceum.*

Crust white, cracked, shields black within,
externally white, with geographical lines.

Walls, winter. *P. Rare.*

Amphill —
crustaceus villosus, ringens, testellis nigris,
extus albus, lineis chartaceis — intus

261

XXIV. Thongs. Crypt. Algæ.

Heaths, January, P. very Rare.

Amphill.

Obs:

Crust very uneven.

870. Pierced Liver-wort. — *L. pertusus.*

Crust with smooth warts, set chequer-wise,
pierced with 1 or 2 cylindrical perforations. — Dillenius. XVIII. 9. Flora Danica.

766. *Sowerby* 677.

Trees, January, P. Common.

871. Yellow-red Liver-wort. — *L. flavo-rubescens.* *S. D.*

Crust greenish ash-color, tubercles orange-red. — Flora. Danica. 955. 1.

872. Trees, January, P. Common.

872. Rusty Liver-wort. — *L. vernalis.*

Crust whitish, tubercles roundish, rust-colored. — Dillenius. LV. 8. *Sowerby*. 845.

Rails, Posts, January, P. Common.

Obs:

Crust ashy-white.

* * * * Liver-worts bearing shields.

873. Rugged Liver-wort. — *L. ater.*

Crust whitish, wrinkled, shields black, margin white, rather scolloped. — Dillenius.
XVIII. 15. A. *Sowerby*. 949.

Walls,

XXIV. Thongs. Crypt. Algæ.

Walls, Bricks, January, P. Common.

Obs:

Shields of various sizes, black with ash-colored borders.

Sent 874. Golden Liver-wort. — L. candelarius. S.D.

Crust yellow, shields gold-colored. — Dillenius. XVIII. 18. B.

Stones on chalky hills, January, P. Common.

875. Pale Liver-wort. — L. pallescens.

Crust whitish, shields pale. — Dillenius. XVIII. 17.

Walls, January, P. Common.

876. Crab's-eye Liver-wort. — L. parellus.

Crust white, shields concave, pale, with blunt edges. — Dillenius. XVIII. 10.

Stones, January, P. Rare.

Barton Hill.

877. Hollowed Liver-wort. — L. scruposus.

Crust ash-colored, granulated, shields immersed, black, with a scolloped border. — Sowerby. 266.

Walls, January, P. Common.

Obs:

Shields imbedded in the crust.

878. Brownish Liver-wort. — L. subfuscus.

Crust

* 879. Notched Liverwort. — L. emarginatus.

Crust ash-colored, uneven, shields white on the exterior surface black within, very deeply notched.

Walls, January, Rare. Bricks walls.

* 873 Contracted Liverwort. — L. coarctatus.

Crust gray, cracked. Shields black, flat, with an elevated, white, mealy margin, when young drawn close over the centre.

English Botany.

Sowerby. 534.

Brick Walls, Autumn, Rare.

Gardens in Bedford.

Obs:

Saunders dark slate colored with white borders.

** 873. Bloody-spotted Liverwort. — L. hematommumCrustaceous, white, mealy with a fibrous border. — Shields imbedded, scarlet, concave: when old, convex: their margin thick white, elevated, mealy — L. sanguinaria. English Botany.

Sowerby. 486.

Brickwalls, Winter, Rare.

Gardens in Bedford.

Obs:

The fructifications of this elegant Liverwort, at first and to the naked eye appear more like wounds in the brickwork, than the marks of vegetation —

mealy bordered Liverwort.

* 870. Black and yellow Liverwort.
Powdery, brownish, shields flat, dull yellow
with a very white mealy elevated border
Sowerby. 432. English Botany
Walls, October, Rare
Friars Wall, on Limestone

* 870. Waxen Liverwort. *L. cerinus*.

Saucers pale yellow, smooth, border and
under-side whitish, crust grey-white.
Dickson III. 14.

Bark of trees in Woods. Winter, Rare
Clapham between Woods. 1799.
Hempston Wood. 1797. +

** Crevice

870. Crink Lichen. *L. cerinus*
remosus

Saucers sea-green, with a white border,
crust whitish, cracked into roundish
angular pieces.

Flora Danica. 168. 3.

Church Walls. Winter, Common Rare
Dapley-Church-Walls.

*** 870. Crisp Liverwort. *L. crenulatus*.
Somewhat tiled and lobed, lobes abridged,
depressed, somewhat rising, rounded, thickish:
shields bordered, between flat and convex.
Dilleni. XIX. 23.

walls, winter not uncommon.
Friars walls.

* 870. Mealy-flesh colored Liverwort
Sowerby 20. 057. *L. Turneri*.
Wilhamstead Wood 1002. ²⁶³ on bark.
XXIV. Thongs. Crypt. Alga.

Crust whitish, shields brown, with an ash-
colored somewhat scalloped border. — Dill-
eni. XVIII. 16.

Trees, January, P. Common.

Obs:

This is a very sportive and variable species.

879. Crested Liver-wort. — *L. cristatus*.

Tiled, leaves between toothed and fringed,
shields larger than the leaves. — Dilleni.
XIX. 26.

Damp Walls, October, A. Common.

880. Olive Liver-wort. — *L. olivaceus*.

Tiled, leaves lobed, olive-colored, shields of
the same color, scalloped. — Dilleni.
XXIV. 77.

Trees, January, P. Common.

881. Common Liver-wort. — *L. parietinus*.

Tiled, leaves curled, deep yellow, shields of
the same color. — Sowerby. 194.

Trees, January, P. Common.

Obs:

Leaves in damp spots of a vivid hue, in exposed and dry
places, deep yellow. Shields always corresponding.

882. Inflated Liver-wort. — *L. physodes*.

Tiled, segments blunt, inflated. — Sowerby.
126.

Oak

+ Obs. to *L. cerinus*.

In the early stage of the plant, its shields
are of the color of beeswax, in its middle
state they become convex, and in its
dearray it loses somewhat of its original
color. The margins are always greyish white.

XXIV. Thongs. Crypt. Algæ.

Oak pales, January, P. Common.

883. Stone Liver-wort. — *L. saxatilis*. S.D.

Tiled, leaves waved, rough, pitted, shields of
Sowerby. 603. a deep bay color. — Dillenius. XXIV. 83.

Oak pales, January, P. Common.

884. Wall Liver-wort. — *L. muralis*.

Shields green yellow, changing to full yellow, border paler, crust greenish, with a tinge of yellow, somewhat tiled. — (Withering.) Hoffman. Enum. Lich. XI. 1.

Sowerby. 1695. Walls, January, P. Rare.

Oakley Church-yard.

885. Starry Liver-wort. — *L. stellaris*.

Tiled, leaflets oblong, jagged, narrow, ash-colored, shields blackish-brown. — Dillenius. XXIV. 70.

Sowerby. 1697. Trees, March, P. Common.

886. Channel'd Liver-wort. — *L. calicaris*.

Leafy, upright, strap-shaped, branched, pitted, convex, ending in a point. — Dillenius. XXIII. 62. Sowerby. 890. *fastigiatus*.

Trees, January, P. Common.

887. Rose Liver-wort. — *L. caperatus*. S.D.

Pale green, wrinkled, waved at the edge. — Dillenius. XXV. 97. Sowerby. 654.

Trees,

* 883. Bordered Liverwort *L. limitatus*. Crust nearly ashcolor, shields black, enclosed by a serpentine line.

Hoffmann Ger. Flora.

Hoffman Num. Lich. V. 3.

Bark of trees, all the Year. Common.

Obs:

This differs especially from the *L. Samiguanus*, with which it has been too much confounded, in not being red without and the fructifications being more shieldlike.

* 885. grey shielded Liverwort. *L. pulverulentus*.

Tiled, leaflets lobed, scalloped, blunt, green, sprinkled with pollen, shields black, with scalloped margins.

Ribthorpe page. 325

Dillenius. XXIV. 71.

Trunks of trees. Nov? Common.

Obs:

The permanency of this variety, for some will not admit it to be a species requires from the Botanist something more than a transient observation and I am persuaded that when the Student shall have contrasted the universally grey tint of this plant with the deeper black of the *stellaris*, he will pronounce it to be distinct —

XXIV. Thongs. Crypt. Algæ.

Trees, January, P. Rare.

Everton,—Eversholt.

888. *Self-colored Liver-wort.* — L. concolor. S. D. ~~seal~~
 Shields reddish-yellow, foliage yellow, leaves
 minute, upright, crowded, curled. — (With- Sowerby. 1794.
 ering.) Dickson. IX. 8.

Trees, May, P. Rare.

Milton-Ernys.

889. *Fringed Liver-wort.* — L. ciliaris.

Leafy, somewhat upright, segments strap-shaped, fringed, shields on fruitstalks, scolopod. — Dillenius. XX. 45. Sowerby. 1352.

Trees, January, P. Common.

Obs :

Small branches fringed.

890. *Little-horned Liver-wort.* — L. corniculatus.

Leafy, somewhat upright, somewhat pitted, smooth, segments narrow, ending in small horns. — Dillenius. XXI. 54.

Trees, January, P. Common.

891. *Mealy Liver-wort.* — L. farinaceus.

Leafy, upright, flat, branched, mealy, with warty edges. — Dillenius. XXXIII. 63. Sowerby. 889.

Trees, January, P. Common.

M m

Asb

The margins of the shields sometimes compleat, but they often go off to powder and render the border indistinct.

Obs.
 This grows on the trunk and branches
 of "Salix latifolia rotunda" Ray., the
 tree that nourishes the larva of Pap. Ina.
 The branches by its growing in great
 profusion exhibit a most grotesque appearance
 looking like a ^{so many} gouty limbs and nearly
 equaling the whole trunk of the Salix
 in size —

XXIV. Thongs. Crypt. Algæ.

892. *Ash Liver-wort*. — *L. fraxineus*.

Leafy, upright, oblong, spear-shaped, pitted, smooth, shields on short fruitstalks. — Dillenius. XXII. 59.

Trees, December, P. Common.

893. *Least Liver-wort*. — *L. tenellus*.

Shields dark-brown, nearly sitting, leaves grey-green, segments blunt, nearly upright, fringed, the ends tubular, when old. — Dillenius. XX. 46. Sowerby. 1351.

Trees, January, A. Common.

894. *Bat's-wing Liver-wort*. — *L. nigrescens*.

Leafy, skinny, roundish, lobed, wrinkled, blackish-green, shields crowded, reddish. — Sowerby. 345.

Willows, October, A. Rare.

Clapham, — Kempston.

Obs :

There are no shields ever found on our plants.

895. *Black-thorn Liver-wort*. — *L. prunastri*.

Leafy, rather upright, pitted, white and downy underneath. — Dillenius. XXI. 55.

Trees, January, P. Common.

Obs :

Shields terminal, paler than the leaves, brown with age.

Lung-wort.

Sowerby. 859.

* 894 Hand-shaped Liverwort. L. palmatus.
Leaves saucers tawny, ~~foliage~~, purplish sea-green, transparent, rather jelly-like, tender, scalloped and indistinctly hand-shaped.
Dillenius XIX. 30.
Heaths. Jane. Sep? Rare.
Amphithe Heath.

XXIV. Thongs. Crypt. Algae.

896. *Lung-wort*. — *L. pulmonarius*.

Leafy, jagged, blunt, smooth, pitted above, downy underneath. — Dillenius. XXIX.

113. Sowerby. 572-

Trees, January, P. Rare.

Maulden Wood.

* * * * * Leathery, bearing targets.

897. *Thrush Liver-wort*. — *L. apthosus*.

Leathery, creeping, lobed, blunt, flat, with scattered warts, target on the edge, ascending. — Dillenius. XXVIII. 106.

Woods, January, P. Rare.

Ampthill, — Maulden.

898. *Dog's Liver-wort*. — *L. caninus*.

Leathery, creeping, lobed, blunt, flat, veiny underneath, woolly, target on the edge, ascending. — Dillenius. XXVII. 102.

Pastures, January, P. Common.

899. *Flat-shield Liver-wort*. — *L. horizontalis*.

Leathery, creeping, flat, not veiny underneath, targets on the edge, horizontal. — Dillenius. XXVIII. 104. Sowerby. 888.

Heaths, Woods, January, P. Rare.

Clophill, — Sheerbatch.

M m 2 * * * * * Shaped

ogg. *Tremella diversicolor*. L. *Tremella*.

Saucers red brown, numerous, minute leaves dark brown, pellucid, crisp, thin notched. Dillenius XIX. 31.

Amongst heapes. April. P. Rare.

Bronham House Moat.

XXIV. Thongs. Crypt. Algæ.

* * * * * Shaped like a drinking glass.

900. *Cup Liver-wort.* — L. pyxidatus.

Cup-bearing, simple, finely scolloped, tubercles brown. — Dillenius. XIV. 6. A. B.
Sowerby. 1393. Walls, Heaths, January, P. Common.

901. *Tuberculated Liver-wort.* — L. tuberculatus.

Cup-bearing, simple, finely scolloped, tubercles brown, sitting on the borders of the cups. — Dillenius. XIV. 6. C. K. L. M.

Walls, Heaths, January, P. Common.

902. *Fringed Liver-wort.* — L. fimbriatus.

Cup-bearing, simple, finely toothed, stalk of the fruit cylindrical. — Dillenius. XIV. 8.

Heaths, January, P. Rare.

Ampthill. — Sandy.

903. *Cornucopia Liver-wort.* — L. cornu-pioides.

Cup-bearing, simple, shorter than the leaf, tubercles scarlet. — Dillenius. XIV. 9.

Heaths, July, P. Rare.

Ampthill.

904. *Trumpet Liver-wort.* — L. filiformis.

Cup-bearing, simple, very entire, thread-shaped, tubercles brown. — Dillenius. XIV. 10.

Heaths,

XXV. Thongs. Crypt. Algæ.

Heads, Woods, January, P. Common.

Leaves mostly upright, fringed, curved, bear

tubercles, cups at first very short, conical

and narrow, afterwards spreading, the

clock tubercles. — Dillenius. XIV. 11.

Heaths, January, P. Common. *Sowerby. 1393.*

905. *Smooth Liver-wort.* — L. glaberrimus.

Cup-bearing, smooth, very short stalks

and narrow, spreading, the

Heaths, January, P. Common.

Leaves mostly upright,

cups entire. — Dillenius.

Hills, January, P. Rare.

Leaves mostly upright,

cups entire. — Dillenius.

Hills, January, P. Rare.

Leaves mostly upright,

cups entire. — Dillenius.

Hills, January, P. Rare.

Leaves mostly upright,

cups entire. — Dillenius.

Hills, January, P. Rare.

XXIV. Thongs. Crypt. Algæ.

Heaths, Woods, January, P. Common.

*Obs :**Cups very fine, truly thread-shaped.*905. *Elk's horn Liver-wort.* — *L. alcicornis.*

Leaves nearly upright, jagged, curled, bearing cups, cups at first very short, conical and simple: afterwards proliferous, with black tubercles.—Dillenius. XIV. 12. A. B.

Heaths, January, P. Common. **Sowerby. 1392.**

906. *Scarlet Liver-wort.* — *L. cocciferus.*

Cup-bearing, simple, very entire, stem cylindrical, tubercles scarlet.—Dillenius. XIV. 7.

Heaths, January, P. Common.

*Obs :**Cups as if tipped with red sealing wax.*907. *Horned Liver-wort.* — *L. cornutus.*

Cup-bearing, nearly simple, rather bellying, cups entire.—Dillenius. XV. 14.

Hills, January, P. Rare.

*Barton Hill.*908. *Rein-deer Liver-wort.* — *L. rangiferinus.*

Shrubby, perforated, very much branched, branches nodding.—Sowerby. 173.

Heaths, Woods, January, P. Common.

*Obs :**Small branches fistulous.**Sbert*

* 908. *Prickly Liverwort.* *L. hispidus.*
Saucers red brown bordered, terminating
plant firm, branched, branches decurrent,
flattish, corners obtuse, points forked
pointed Dillenius. XVII. 31.

Heaths March, June. Rare
Amphithe Heath.

XXIV. Thongs. Crypt. Algæ.

909. *Short Liver-wort.* — L. uncialis.

Shrubby, perforated, small branches very short, pointed. — Sowerby. 174.

Heaths, January, P. Rare.

Amphibill.

Obs :

Barren shoots acute, fertile ones blunt.

* * * * * Branches hair-like.

910. *Hairy Liver-wort.* — L. hirtus.

Thread-like, very much branched, upright, tubercles mealy, scattered. — Dillenius. XIII. 12. Sowerby. 1354.

Old Rails, January, P. Common.

911. *Wiry Liver-wort.* — L. jubatus.

Thread-like, branched, somewhat hanging down, between matted and zigzag, wings of the shoots compressed. — Dillenius. XII. 7.

Park Pales, January, P. Rare.

Amphibill.

Obs :

Plant like the tail of a horse in miniature.

Star-jelly. TREMELLA.

Fructifications scarcely visible, situated in a gelatinous substance.

912. *Brain-like Star-jelly.* — T. albida.

Sitting,

XXIV. Thongs. Crypt. Algæ.

*913 *Frenella Savine.*
Sowerby.
Eng. Botany. tab. 710.
 On Savine, Oakley Plantation
 behind the house —

914
 Substance of the nature of wax
 plaited into various shapes.

Sitting, jelly-like, of various shapes, whitish. — Bulliard. 386.

Fallen branches of trees, October, A. Common.

Obs:

Plant white when young, when old brownish black.

913. *Tree Star-jelly.* — *T. arborca.*

Sitting, roundish, waved, black. — Dillenius. X. 15.

Fallen branches of trees, October, A. Common.

Obs:

The whole surface studded with very black tubercles.

914. *Yellow Star-jelly.* — *T. mesenterica.*

Sitting, membranaceous, jelly-like, crowded, waved, yellow. — Jacquin. Misc. Aust. I.

13. *Sowerby. 709.*

Pales, Stumps, &c. September, A. Rare.

Ampthill, — Warden.

915. *Common Star-jelly.* — *T. Nostoc.*

Sitting, roundish, plaited, waved, brownish-green. — Sowerby. 461.

Pastures after rain, August, A. Common.

916. *Violet Star-jelly.* — *T. violacea.*

Sitting, jelly-like, wrinkled, violet-colored, smooth underneath.

Fallen

XXIV. Thongs. Crypt. Algæ.

Fallen branches of trees, September, A.
Common.

Obs :

Like the tartar of red wine.

917. *Blackish Star-jelly.* — T. nigricans. S.D.

Fleshy, thickish, with a very irregular surface, at first scarlet, afterwards black. — Bulliard. 455. 1.

Stumps, September, A. Rare.

Southill, — Clophill, — Ampthill.

918. *Ruddy Star-jelly.* — T. cinnabarina.

Jelly-like, fleshy, surface smooth, reddish. — Bulliard. 455. 2.

Pales, October, A. Rare.

Ampthill Park.

Obs :

Plant sitting, hemispherical, at first soft, afterwards gristly.

919. *Deliquescent Star-jelly.* — T. deliquesens.

Nearly round, sitting, jelly-like, deliquescent, smooth, yellow. — (*Sibthorpe.*) Bulliard. 455. 3.

Stumps, November, A. Common.

Obs :

Like the gum that exudes from injured trees.

920. *Glandulous Star-jelly.* — S. glandulosa.

Jelly-like,

XXIV. Thongs. Crypt. Algæ.

Jelly-like, roundish, cushion-shaped, brownish-black, entire, surface breast-shaped. — Bulliard. 420. 1.

Fallen branches of trees, December, A. Common.

921. *Charcoal Star-jelly.* — T. carbonaria. S.D.

Burnt, small, in vascular substances. — Bulliard. 420. 2.

On decayed Lemons, Winter, A. Common.

Obs :

Small globular substances arising from membranaceous cells.

Laver. ULVA.

Fructification in a transparent membrane, watery.

922. *Gut Laver.* — U. intestinalis.

Tubular, simple, membranaceous, green. — Dillenius. IX. 7.

Rivers, Ditches, May, A. Common.

923. *Curled Laver.* — U. crispæ.

On the ground, very tender, green, curled. — Dillenius. X. 12.

Stones, January, A. Rare.

Elstow.

N n

Plumb

XXIV. Thongs. Crypt. Algæ.

924. *Plumb Laver.* — *U. pruniformis.* S. D.Roundish, single, succulent within. — Weiglers obs. II. 4. *Sowerby.* 968.

On Moss, October, A. Rare.

*On the Bath at Wrest Park.**Obs:**Green, not bigger than a Sloe.**River-weed.* *CONFerva.* Gen. Pl. 1672.

Fibres simple, uniform, fine as a hair, thread-like.

925. *Cotton Conferva.* — *C. bullosa.*Threads equal, branched, matted, inclosing bubbles of air. — *Dillenius.* III. 11.

Ditches, March, A. Common.

926. *River Conferva.* — *C. rivularis.*Threads very simple, equal, very long. — *Dillenius.* II. 1. *Sowerby.* 1654.

Ditches, January, P. Common.

*Obs:**Plant floating, silky.*927. *Mill Conferva.* — *C. canalicularis.* S.D.Threads equal, more branched towards the base. — *Dillenius.* IV. 15.

Mill-ponds, January, P. Common.

*Obs:**Threads 3 to 4 inches long, branches matted.**Amphibious*

Line 4
for Weiglers read
Weigel.

Phascum caulinicapsulare

Molehills near

Clapham Springs

March 29. 1799

XXIV. Thongs. Crypt. Algæ.

928. *Amphibious Confervæ*. — C. amphibia. **SD.**

Threads equal, branched, when dry uniting
into points like prickles. — Dillenius. IV.

17.

Damp Walls, January, P. Rare.

*Luton Hooe.**Obs :*

It forms a kind of skin over the mould with it's very numerous small fibres.

929. *Frog-spawn Confervæ*. — C. gelatinosa.

Threads branched, necklace-shaped, joints
round, jelly-like. — Dillenius. VII. 42. **Sowerby**. 689.
Springs, January, A. Rare.

*Eversbott.*930. *Net Confervæ*. — C. reticulata.

Threads net-like, uniting together. — Dille-
nius. IV. 14. **Sowerby**. 1687.

Ditches, May, A. Rare.

*Woburn, — Aspley.**Obs :*

Threads branching in various directions, composing a net.

931. *Clustered Confervæ*. — C. glomerata.

Threads jointed, very much branched, little
branches very short, in many divisions. —
Dillenius. V. 31.

Collingham

N n 2

Springs,

XXIV. Thongs. Crypt. Algæ.

Springs, April, A. Rare.

Eversholt.

Obs:

Midrib thicker than the other parts.

Byssus. *Byssus.* Gen. Pl. 1673.

Fibres simple, downy.

932. *Paper Byssus.* — B. flos aquæ.

Threads feathery, floating.

Stagnant Waters, May, A. Common.

Obs:

In a dry state very like coarse whitey-brown paper.

933. *Violet Byssus.* — B. phosphorea.

Downy, violet colored, adhering to wood. —

Dillenius. I. 6.

Stumps, October, A. Common.

Obs:

Plant velvety, extremely grand when viewed in sunshine.

934. *Velvet Byssus.* — B. velutina.

Hair-like, green, threads branched. — Dille-
nius. I. 14.

On the ground, October, A. Common.

Obs:

Unless seen with a magnifier, this plant has the appearance
of some of the smaller mosses.

935. *Golden Byssus.* — B. aurea.

Hair-like,

XXV. Thongs. Crypt. Algæ.

Hair-like, powdery, fibrofibrilliferous, coarse,
threads simple and branched. — Dillenius.
I. 16. Smirnov. 122.

On trees, January, P. Rare.

Brownish.

Obs:

Wet dark yellowish brownish green.

936. *Yellow Byssus.* — B. flos.

Threads branched, yellow. — Dillenius. I. 6.
Fallen branches of trees, January, P. Com-
mon.

937. *White Byssus.* — B. candida.

Threads very much branched, small branches
bundled, white. — R. Smirnov. 123.
Rotten wood, September, A. Common.

938. *Yellow Byssus.* — B. flos.

Hair-like, very pale, yellowish, very brittle
again. — Dillenius. I. 6.

Culture: Dillenius. B. Common.

939. *Yellow Byssus.* — B. aurea.

Yellow, yellow, golden brownish yellow.
Dillenius. I. 6.

Common: Dillenius. B. Common.

XXIV. Thongs. Crypt. Algæ.

Hair-like, powdery, fructifications scattered, threads simple and branched. — Dillenius.
I. 16. Sowerby. 212.

On Roots, January, P. Rare.

Barton Hill.

Obs :

When fresh, golden; when dry, pale-green.

936. *Tawny Byssus.* — *B. fulva.*

Threads branched, tawny. — Dillenius. I. 17.
Fallen branches of trees, January, P. Common.

937. *White Byssus.* — *B. candida.*

Threads very much branched, small branches bundled, whitish.—Ray's Synopsis. XXIII.
Rotten wood, September, A. Common.

938. *Flaky Byssus.* — *B. septica.*

Hair-like, very soft, parallel, very brittle, pale. — Dillenius. I. 9.

Cellars, December, P. Common.

939. *Yellow Byssus.* — *B. candelaris.* S.D. *Sent*

Powdery, yellow, growing upon wood. — Dillenius. I. 4.

Bark of Trees, September, A. Common.

Obs :

Different from the Lichen candelarius.

Mealy

Byssus. *Byssus.* Gen. Pl. 1613.

Fibres simple, downy.

935. *Flax Byssus.* — *B. fibrosa.*

Threads slender, fibrous.

Southern Waters, Italy, A. Cannon.

Obs :

Threads very much branched, simple.

936. *Vine Byssus.* — *B. phosphorea.*

Young, violet colored, adhering to wood. —

Dillenius. I. 1.

Stems of Trees, Rotten.

Plantain, extremely good eaten raw as a vegetable.

934. *Velvet Byssus.* — *B. velutina.*

Threads very closely bundled —

On the ground, October, A. Common.

Obs :

Outer surface a greyish, irregular, silvery white, rest of the plantaceous.

935. *Golden Byssus.* — *B. aurea.*

Hair-like,

XXIV. Thongs. Crypt. Algae.

940. *Mealy Byssus.* — *B. incana.* **S.D.***Sowerby. 1603.* Powdery, mealy. — Dillenius. I. 3?
Trees, October, A. Common.

XXIV. Mushrooms. Crypt. Fungi.

Merulius.

Cap with veins underneath of the same substance with the rest of the plant.

941. *Minute Merulius.* — *M. squamula.*

Cap convex, flattish, white, with some indistinct wrinkles, radiate on the under side, stem very tender, between awl and bristle-shaped. — Batsch. 84. E. Fungi. 93.

Woods, on poplar leaves, October, A. Rare.

*Clapham Park & Renhold Woods.**Obs :**Beneath the cap instead of gills there are only from 3 to 5 irregular plait.*942. *Clove Merulius.* — *M. caryophylleus.*

Sitting, concave, shaggy, edge scolloped and waved, veins branched underneath. — Batsch.

E. Fungi. 213. 121. Schæffer. 325. Sowerby. 9

Fir Groves, September, A. Rare.

*Amptbill, — Warden.**Obs :**Plant dark cinnamon-colored, tiled, edges feathery.**Chanterelle*

XXIV. Mushrooms. Crypt. Fungi.

943. Chanterelle. — *M. chantarellus.*

On a stem, cap irregularly funnel-shaped, orange-colored, veins branched, anastomosing, stem flattish, of the same color. — E. Fungi. 46. excellent.

Woods, July, A. Common.

Obs:

Stem tapering at the base.

944. Skinny Merulius. — *M. membranaceus.*

Sitting, membranaceous, smooth, lobed, curled at the edge, veins on the underside branched. — Bulliard. 498. i.

Fir Groves, on Moss, January, A. Rare.

Ampthill Park. - Hafocks: walls -*Obs:*

Cap very irregular, with a border much waved and uncommonly curled, veins like net-work.

Agaric. AGARICUS. Gen. Pl. 1674.

Mushroom horizontal, with gills underneath.

* On a stem, solid with decurrent gills.

945. Stick Agaric. — *A. ramealis.*

Cap convex, white, stem white. — Bulliard.

336.

Small hazel sticks, October, Common.

Obs:

Hazel branches, where they shoot out from the old stumps, are almost entirely covered with this species.

Ivory

XXIV. Mushrooms. Crypt. Fungi.

946. Ivory Agaric. — *A. eburneus*.Snowy-white, gills in pairs, stem short. —
E. Fungi. 32. excellent.

Hills, Heaths, September, Common.

Obs:

Cap in its old age irregular; when young, hemispherical.

947. Drinking-glass Agaric. — *A. cyathiformis*.Cap shaped like a drinking-glass, white, stem
of the same color. — Bulliard. 248.

Fir Groves, September, Rare.

Ampthill, — Maulden.

Obs:

Stem zigzag, cap pale brown.

948. Funnel Agaric. — *A. infundibuliformis*.Cap funnel-shaped, waved, stem thickening
upwards. — Bulliard. 286. ~~E. Fungi. 106.~~

Fir Groves, November, Common.

Obs:

Cap brown, leather-like, stem and gills of the same color.

949. Pestle Agaric. — *A. pistillaris*.Gills whitish, in pairs, very short, cap whitish,
convex, stem of the same color, conical,
bent inwards. — Batsch. 62.

Roots

*946. Deep buff Agaric — *A. miniatus*.Gills in pairs, stem tapering, cap
deep buff colour. — E. Fungi. 141.

Barren Heaths — October — Rare.

Ampthill Clapham Springs

*Obs:*Caps often turning up, and sometimes
proliferous, splitting at the edge.*948 Lobed Agaric *A. lobatus*.Cap lobed, cracking, deep bay,
center darker, gills uniform
greyish brown, stem twisted.

E. Fungi. 106.

September Pastures. November, Rare.

Houghton Conquest —

*Obs:*Plants generally larger than in
Mr. Sowerby's figures.

*950. Turfstack Agaric. A. turfosus.

^{brown}
Cap with concentric zones, gills plaitlike
anastomosing, stem fawn colour.

E. Fungi 210.

On Heaths where turf stacks have stood
Winter Rare

Ampathia

Obs:

The zones on the cap are indistinct,
The cap itself cupping up so as
to give it the appearance of a Peccaria
indeed it is not impossible but that
upon a revision it may be re-
moved to another genus.

281

XXIV. Mushrooms. Crypt. Fungi.

Roots of Trees, Winter, Common.

Obs:

Stem not upright.

950. Cheese Agaric. — A. caseus.

Cap pale-grey, margin turned in, gills whitish, stem cylindrical, thickening downwards. — E. Fungi. 61. excellent. Bulliard. 400.

Woods, Autumn, Common.

Obs:

Cap at first conical, when in perfection funnel-shaped.

951. Lister's Agaric. — A. Listeri.

Gills whitish, mostly regular, very thickly set and narrow, cap white, smooth, irregular, flattish, center flattened, stem white, not central, cylindrical. — Bulliard. 300.

Woods, September, Rare. E. Fungi. ~~245~~ 245.

Clapham Park Wood, — Silsoe.

Obs:

The plant is full of an highly acrid juice.

952. Fragrant Agaric. — A. fragrans.

Gills brown-white, 4 in a set, cap brown-white, semi-transparent, stem brown-white. (Withering.) E. Fungi. 10. excellent.

O e

Under

XXIV. Mushrooms. Crypt. Fungi.

Under Chesnuts, October, Rare.

Silsoe.

953. *Umbrella Agaric.* — A. *Umbraculum*.

Cap expanded, stem twisted, ash-colored above, brown below, gills 4 in a set. —
Batsch. II. 4.

Stumps, September, Rare.

Bromham.

Obs:

Stem twisted upwards, channelled downwards.

954. *Milky Agaric.* — A. *lactifluus*.

Cap nearly expanded, but having a central point, nearly brown, gill and stem of the same color. — Bulliard. 224. E. *Fungi*. 204.

Woods, August, Common.

Obs:

Stem slanting, milk not acrid.

955. *Amethystine Agaric.* — A. *amethystinus*.

Cap nearly bell-shaped, smooth, stem and
E. *Fungi*. 187. gills violet-blue. — Bulliard. 198.

Woods, September, Common.

Obs:

The whole plant amethystine.

956. *Heath Agaric.* — A. *ericeus*.

Cap

* 957.

Delicious Agaric. *A. deliciousus.*

Cap rather undulating, Zoned. rich red brown, gills flame coloured watery.
Stem: orange

Sowerby. 202.

Woods & Hills. October, Common

Obs:

Gills distinct where they set off from the Cap, branching twice before they reach the margin —

283

XXIV. Mushrooms. Crypt. Fungi.

Cap white, clammy, margin thin, stem of the same color. — Bulliard. 188.

Heaths, October, Common.

Obs:

Stem obversely conical, bluntish.

957. *Biting Agaric.* — *A. piperatus.*

Gills pale, 2 in a set, cap yellowish-white, woolly, center flattened, stem pale-yellow.

Woods, September, Rare.

Silsoe.

Obs:

The plant abounds with an highly acrid milky fluid.

958. *Bluish-red Agaric.* — *A. livido-rubescens.* SD.

Stiff, brittle, pithy, milky, ashy-livid, cap brownish, spotted, stem whiter, gills very pale-yellow. — Batsch. 202.

Fir Groves, October, Common.

Obs:

Stem solid only above, with a singular hollowness in the lower part.

959. *Glutinous Agaric.* — *A. velatus.*

Cap brownish red, flattish, gills dirty white, few, 4 in a set, stem cylindrical. — E. Fungi. 7. excellent.

Fir Groves, September, Rare.

Oo 2

Warden,

XXIV. Mushrooms. Crypt. Fungi.

Warden, — *Southill*. Cap white.*Obs*:

Cap generally dark-brown.

960. *Slimy Agaric*. — *A. limacinus*.

Gills dirty-watery-white, not numerous, 4 in a set, cap pinky-brown, nearly flat, edge turned down, stem buff, tapering downwards. — E. Fungi. 8.

Fir Groves, October, Rare.

Warden.961. *Yellow Agaric*. — *A. sulphureus*.

Cap at first convex, then flattened, brimstone-colored, gills of the same color. — E. Fungi. 44.

Woods, October, Rare.

Silsoe, — *Warden*.*Obs*:

The whole plant of the same color, border of the cap uneven.

962. *Black-stalked Agaric*. — *A. androsaceus*.

White, cap skinny, plaited, stem black. — E. Fungi. 94.

Woods, October, Common.

Obs:

Stem not quite straight, darkish brown.

963. *Woolly-edged Agaric*. — *A. contiguus*.

Cap

XXIV. Mushrooms. ^{or} Crypt. Fungi.

Cap brownish-red, smooth, brim turned in, woolly, gills narrow, thickish, branched, pale yellow. — E. Fungi. 56.

Fir Groves, September, Common.

Obs :

Stem very short in proportion to the size of the plant.

964. *Ruddy Agaric.* — A. rutilus.

Cap rather conical, ruddy, gills reddish, leather-like, 2 in a set, stem red, bent at the base. — Schaeffer. 55. **E. Fungi. 105.**

Fir Groves, October, Common.

Obs :

The whole exterior surface of the plant nearly of the same color.

965. *Small Orange Agaric.* — A. fibula.

Cap orange, scored, gills orange, in pairs, stem orange. — E. Fungi. 45.

Amongst Moss, August, Common.

Obs :

Cap dimpled in the center.

966. *Spangled Agaric.* — A. micaceus.

Cap bluntly conical, point brown, gills greyish-black, crowded. — Bulliard. 246. excellent.

Banks, Autumn, Common.

Obs :

XXIV. Mushrooms. Crypt. Fungi.

Obs:

Plant in decay dissolving into an ink-like liquor, when young ovate, in old age very ragged.

967. Elm Agaric. — *A. ulmarius.*

Cap pale-yellow, gills white, irregular, stem short, thick, twisted, of the same color. — E. Fungi. 67.

Sides of Elms, Autumn, Rare.

*Ravensden.**Obs:*

The cap by the cracking of the flesh has the appearance of net-work. Stem thickening downwards.

968. Gilliflower Agaric. — *A. odoratus.*

Gills white, 8 in a set, cap glaucous-green, nearly flat, stem white, cylindrical. — E. Fungi. 42.

Fir Groves, October, Rare.

*Warden.*969. Confluent Agaric. — *A. confluens.*

Cap funnel-shaped, deep-brown, gills watery-brown, stem tapering downwards, of the same color. — E. Fungi. 168.

Woods, October, Rare.

*Clapham Park Wood.**Obs:*

The plant grows in large clusters from one spindle-shaped root with caps incorporating.

Meadow

trifarium.

XXIV. Mushrooms. Crypt. Fungi.

970. Meadow Agaric. — *A. elixus.*

Cap uneven, dark-brown in the center, lighter towards the margin, stem leaning, gills watery-white. — E. Fungi. 172.

Meadows, Autumn, Common.

* * On a stem, gills solid and fixed.

971. Broad-gilled Agaric. — *A. araneosus.*

Cap conical, cracking, gills broad, 4 in a set, stem brownish, ring fugacious. — Bulliard.

431.

Under Willows, Autumn, Common.

Obs.:

Cap turned up in an advanced state, seldom entire.

972. Goat Agaric. — *A. Cossus.*

Cap conical, glutinous, white, shining, gills watery-white, stem of the same color, leaning. — E. Fungi. 121.

Woods, October, Common.

973. Caesar's Agaric. — *A. xerampelinus.*

Gills golden-yellow, 4 in a set, cap fine red with shades of buff, stem buff, tapering upwards. — E. Fungi. 31.

Fir Groves, Autumn, Rare.

Warden, — Ampibill.

Dove

971 * opaque Agaric. *A. opacus.*
 Cap color of whitewash, almost flat
 Gills 2 or 4 in a set, stem
~~slits~~ of the same color, with brown.
 Plantations October, Rare
Clophill. 1899. J.H.

XXIV. Mushrooms. Crypt. Fungi.

974. *Dove Agaric.* — *A. columbarius.*

Cap conical, lead-color, gills bluish, stem
lead-color in the middle, whitish at both
ends. — E. Fungi. 161.

Woods, September, Common.

975. *Saffron Agaric.* — *A. croceus.*

Ochry, cap nearly conical, gills whitish,
stem shaggy. — E. Fungi. 19.

Fir Groves, October, Rare.

Ampthill, — Warden.

Obs :

Cap ornamented with marginal ochry tufts.

976. *Spindle-stemmed Agaric.* — *A. crassipes.*

Cap brown, changeable, convex, cracking,
gills brownish, stem grooved, awl-shaped at
the base, thicker above. — E. Fungi. 129.

Pastures, Autumn, Common.

Obs :

Plants growing in clusters, gills distant, 4 in a set.

977. *Whitish Agaric.* — *A. dealbatus.*

Cap whitish with shades of brown, gills and
stem watery-brown. — E. Fungi. 123.

Fir Groves, Autumn, Common.

978. *Elephant Agaric.* — *A. elephantinus.*

Cap

XXV. Mushrooms. Crypt. Fungi.

Cap thick, obovate, chinney,
a few greenish tufts, stem
thin, pale, smooth.

Woods, October, Common.

Small annual species, common.

Cap brownish, conical, gills whitish,

Stems, thin, smooth.

Cap brownish, conical, gills whitish.

XXIV. Mushrooms. Crypt. Fungi.

Cap thick, olive-color, clammy, gills 3 in a set, greenish yellow, stem and cap of one color. — E. Fungi. 36.

Woods, November, Common.

Obs:

The cap of the plant in decay has a black tinge.

979. Small conical Agaric. — A. geophyllus.

Cap bluish, conical, gills dirty brown, in pairs, stem brownish. — E. Fungi. 124.

Woods, Autumn, Common.

980. Cobweb Agaric. — A. glaucopus.

Gills brown, 4 or 8 in a set, cap hemispherical, chesnut-colored, flattish, brim rolled in, stem thick, whitish-pink, ring like a cobweb. — Bulliard. 96. E. Fungi. 223.

Woods, September, Common.

Obs:

Ring fugacious, small particles often adhering to the brims of the cap.

981. Fawn-colored Agaric. — A. hinnuleus.

Gills buff, very broad, 4 in a set, cap fawn-colored, convex, mealy, stem reddish brown at the base. — E. Fungi. 173. excellent.

Fir Groves, October, Rare.

Warden.

P p

Crimson

XXIV. Mushrooms. Crypt. Fungi.

982. *Crimson Agaric.* — A. integer.

Cap clammy, red, rather convex, all the gills uniform and white. — Batsch. 13.

E. Fungi. 201.

Woods, September, Common.

Obs:

The cap varies between rose-color, scarlet, yellow and dark brown.

983. *Fly Agaric.* — A. muscarius.E. Fungi.
286.

Cap blood-colored, warty, warts and gills white, stem bulbous at the base. — (Sibthorpe.) Bulliard. 122.

Fir Groves, September, Common.

Obs:

Cap splendid, hemispherical, the ring turning back after the expansion of the cap.

984. *Spanish-snuff Agaric.* — A. orichalceus.

Cap clammy, rust-colored in the middle, stem ending in a sharp bulb, ring like a cobweb, rust-colored as well as the gills which are elongated, stem bulbous, livid, yellowish, or very pale-green. — Batsch. 184.

Fir Groves, Autumn, Rare.

*Amptbill.**Obs:*

All the cap (except the brim) rust-colored.

Spatterdash

XXIV. Mushrooms. Crypt. Fungi.

985. *Spatterdash Agaric.* — *A. peronatus.*

Cap hemispherical, nearly transparent, gills straw-color, stem thickening downwards, woolly. — (*Sibthorpe.*) E. Fungi. 37.

Fir Groves, Woods, October, Rare.

Clophill, — Maulden.

Obs :

Stem clothed half way up with yellow wool.

986. *Cracking Agaric.* — *A. rimosus.*

Gills olive-brown, 2 or 4 in a set, cap striped, reddish-brown and yellow, conical, bossed, stem yellowish-white, cylindrical.— (*Withering.*) Bulliard. 388.

Woods, August, Common.

987. *Satiny Agaric.* — *A. rufus.*

Gills tawny, not in a regular series, cap dark-brown, glossy, stem red-brown. — Bulliard.

526.

Fields, September, Common.

988. *Stump Agaric.* — *A. stipitis.*

Cap pale-brown, center darker, woolly, gills 4 in a set, white, stem brownish-grey, bulbous at the base, ring white. — (*Withering.*) E. Fungi. 101.

Stumps of trees, Autumn, Common.

P p 2

Obs :

XXIV. Mushrooms. Crypt. Fungi.

*Obs:**Cap cracking, ring permanent.*989. *Sordid Agaric.* — *A. sordidus.*

Gills and cap livid-brown, center dimpled, brim bent downwards, stem thicker at the base. — (*Dickson.*) *Dickson.* I. III. 1.

Fir Groves, October, Rare.

Warden, — Ampthill.

*Obs:**Cap fixed slopingly on the stem.*990. *Earthy Agaric.* — *A. terreus.*

Cap convex, brittle, ashy-black, gills white, stem white, not central. — *E. Fungi.* 76.

Woods, Heaths, October, Common.

*Obs:**Stem not upright, twisted, cap splitting.*991. *Warty Agaric.* — *A. verrucosus.*

Cap convex, ash-colored, warty, warts and gills white, stem bulbous at the base. —

(*Hudson.*) *Curtis.* V. 72.

Woods, September, Common.

*Obs:**Cap brownish, warts crowded.*992. *Green Agaric.* — *A. viridis.*

Gills white, fleshy, 4 in a set, cap bluish-green, stem cylindrical, whitish. — (*Withering.*) *Bolton.* 12.

Fir

XXIV. Mushrooms. Crypt. Fungi.

Fir Groves, Woods, August, Rare.

*Clophill, — Warden.**Obs :**Cap greyish-blue, stem without any ring.*993. *Snuff Agaric.* — *A. violaceus.*Cap cracking, brim violet, woolly, stem blue-
ish, with rust-colored wool.—Bulliard. 250. *E. Fungi 209*

Woods, October, Common.

*Obs :**Brim turned inwards, gills 8 in a set, many, splendid
violet.*994. *Fetid Agaric.* — *A. graveolens.* *SD.*Gills white, very numerous, irregular, cap
white, stem of the same color, tapering.—Bulliard. 585. *E. Fungi. 281.*

Pastures, May, Rare.

Ampthill Park, — Thurleigh.

* * * On a stem, gills solid and loose.

995. *White Agaric.* — *A. albus.*Gills white, many, 4 in a set, cap white,
bluntly conical, brown at the point, stem
white, tapering upwards.— Bulliard. 256.

Woods, Pastures, November, Common.

*Obs :**Root tuberous.**Bulbous*

XXIV. Mushrooms. Crypt. Fungi.

996. *Bulbous Agaric.* — *A. bulbosus.*

Cap hemispherical, glossy, glutinous, white, stem bulbous, of the same color, ring permanent. — Bolton. 48.

Woods, October, Rare.

Silsoe.

Obs:

Cap yellowish-green, quite smooth and free from warts.

997. *Common Agaric.* — *A. campestris.*

Cap convex, scaly, whitish, gills red. — Bulliard. 134. 514. *E. Fungi.* 305.

Pastures, September, Common.

998. *Nail Agaric.* — *A. clavus.*

Gills white, in pairs, cap dimpled, pale-orange, stem of the same color. — Bulliard. 148. *B. C. D.*

Amongst Moss, July, Common.

Obs:

Stem leaning.

999. *Smeared Agaric.* — *A. collinitus.*

Cap bright-bay, conical, gills covered with a veil of gluten, stem whitish, scurfy. — *E. Fungi.* 9.

Fir Groves, Autumn, Rare.

Warden, — Ampthill.

Wafer

*996. *Fishy Agaric.* — *A. vaginatus.*

Cap horizontal, gills white
stem scarcely annulated, thickening downwards — .
Bulliard. f.

~~Obs:~~
Pastures. September — ~~23~~ 1800. Elstow.

Obs:

Smells like stinking fish.

*997 *S. George's Agaric.* — *A. Georgii.*

Cap convex, scaly, brownish white, gills brownish red, border cracking.

E. Fungi. 304.

Under ledges about St. George's Day.


Obs:

Common.

Bark more spongy than in the *campestris*, gills less red, the pileus has been known to exceed 10 inches in diameter — .

X 1001

Bay Agaric. — *A. collinicus*. spp.



295

XXIV: Mushrooms. Crypt. Fungi.

1000. *Wafer Agaric.* — *A. crustuliniformis*.

Gills brown-red, cap red-buff, whitish at the edge, convex, stem white, scurfy. — Bulliard. 308. 546.

Copse, September, Rare.

Putnoe.

1001. *Lurid Agaric.* — *A. luridus*.

Gills blue-grey, numerous, uniform, cap dirty-olive-brown, slimy, bluntly conical, brim irregularly lobed: stem dirty-olive-brown, bent inwards. — (Withering) Bolton. 25.

Fir Groves, Autumn, Rare.

Amphill.

Obs :

The gills approach the stem, but are not fixed to it.

1002. *Parasitical Agaric.* — *A. lycoperdonoides*.

Cap pill-shaped, branny, gills 18 to 20, very entire. — Bulliard. 166.

On rotten Agarics, Autumn, Rare.

Silsoe.

Obs :

Flesh rather firm, taking up almost the whole head of the Agaric.

1003. *Minikin Agaric.* — *A. minutulus*.

Cap thin, scored, brownish, gills few, yellowish, stem whitish, naked. — Schæffer. 308.

Under

XXIV. Mushrooms. Crypt. Fungi.

Under Black-thorns, Autumn, Common.

1004. *Fairy's Agaric.* — A. orcadensis.

Gills brown-watery-white, 2 or 4 in set, cap pale-brown, convex, irregular, stem whitish, browner with age, very tough, rarely central. — (Witbering.) Bulliard. 144. E. Fungi.

Pastures, November, Common.

247.

Obs:

Plant growing in circles.

1005. *Cushion Agaric.* — A. pulvinatus.

Gills ashy-white, 4 in a set, cap raised, brown mouse-color, stem grey, tapering upwards, ring constant. — Bolton. 49.

Woods, Autumn, Rare.

Clophill, — Silsoe,

Obs:

The edge of the cap deeply scored.

1006. *Hairy Agaric.* — A. pilosus.

Cap light-chesnut, with long brown hairs, stem pale brown, darker at the base, gills watery-white. — E. Fungi. 164.

Mountainous spots, Autumn, Rare.

Warden, — Barton.

1007. *Fibrous-rooted Agaric.* — A. radicosus.

Cap hemispherical, ring torn, permanent, gills ochry, stem shaggy, root slanting. — Bulliard. 160. excellent.

Woods,

XXIV. Mushrooms. Crypt. Fungi.

Woods, Autumn, Rare.

Bishopsgate Park Woods.

Obs:

Cap brownish, with brownish spots.

1008. *Irregular Agaric.* — A. irregularis.

Cap cushion-like shaped, brownish, milk-white,

stem bare, root spindle-shaped. — (Relhan.)

E. Fungi. 42. excellent.

Woods, September, Common.

Obs: — On a stem, hollow with decurrent

gills.

1009. *Milky Agaric.* — A. farinaceus.

Cap rather convex, purplish-green, gills

convex, cream-watery-pink, milky, stem

long. — Bartsch. 106. E. Fungi. 240.

For Groves, September, Common.

Obs:

Cap pale, not very thick, with a thin

1010. *Irregular Agaric.* — A. irregularis.

Cap brown, lobed, wavy, gills 3 in a set,

stem pale-brown, stem hollow, pointed. —

(Relhan.) Bolton. 47.

Woods, October, Rare.

Obs:

Cap brownish, with brownish spots.

1011. *Obscure Agaric.* — A. obscurus.

Cap brownish, with brownish spots.

XXIV. Mushrooms. Crypt. Fungi.

Woods, Autumn, Rare.

Silsoe,—Clapham Park Wood.

Obs :

Cap variegated with brownish spots.

1008. Long-rooted Agaric. — *A. radicatus*.Cap rather bell-shaped, brownish, gills white,
stem bare, root spindle-shaped. — (*Relhan.*)
E. Fungi. 48. excellent.

Woods, September, Common.

Obs :

Cap at first conical, afterwards expanding.

* * * * On a stem, hollow with decurrent
gills.1009. Mealy Agaric. — *A. farinaceus*.Cap rather convex, purplish-brown, gills
convex, distant, watery-pink, mealy, stem
long. — Batsch, 100. *E. Fungi.* 200.
Fir Groves, September, Common.

Obs :

Cap gills and stem very often of one color.

1010. Irregular Agaric. — *A. irregularis*.Cap brown, lobed, waved, gills 3 in a set,
broad, pale-rose, stem hollow, greyish. —
(*Relhan.*) Bolton. 13.

Woods, October, Rare.

Silsoe.

Obs :

Cap fixed slanting on the stem.

Q q

Snow-white

XXIV. Mushrooms. Crypt. Fungi.

1011. Snow-white Agaric. — *A. niveus*.

Cap white, viscid, flattish, gills white, in pairs, stem white, cylindrical. — (Withering.) Schaeffer. 232.

Fir Groves, October, Common.

Obs.:

Gills varnished, not numerous.

1012. Dimpled Agaric. — *A. umbilicatus*.

Gills 4 in a set, cap dimpled, stem hollow, cap, gills and stem leather-like and uniform in color. — Bulliard. 411. 2.

Pastures, October, Rare.

Southill, — Aspley.

Obs.:

Cap between the color of wax and leather.

* * * * * On a stem, hollow and fixed.

1013. Verdegrise Agaric. — *A. aeruginosus*.

Cap convex, bluish-green, clammy, gills purplish-brown. — Curtis. V. 70.

Woods, October, Rare.

Ampthill, — Silsoe.

Obs.:

Gills lilac, ring permanent.

1014. Orange Agaric. — *A. aurantius*.

Cap conical, clammy, orange-color, gills yellow, stem naked. — Curtis. V. 69.

Pastures,

1013. Deformed Agaric. *A. monstrosus*.

Cap flattish, turning in, dead white,
Gills uniform 2 to 4 in a set,
stem tapering upwards and downwards.

S. Fungi. 203.

Roadsides. Octth rare.

Garden, Henlow down

Obs.:

Often growing two from the same root, when the stems and caps coalesce as in the figure of Mr.

Loverby

1015. Branched Agaric. *A. racemosus*.
Cap convex, quaker: colour, Stems
crowded with small pedunculated
buttons of the same. Gills 2 in a
set off the same colour.

Woods. Autumn. Rare. E. Fungi. 287.
Melchtonne Wood

Pastures. October. Rare.

XXIV. Mushrooms. ⁰⁰²⁹⁹ for Crypt. Fungi.

Pastures, October, Common.

Obs:

No Agaric varies more in color and shape than this species.

1015. Filberd Agaric. — *A. auricomus*.

Gills brownish-white, few, in pairs, cap golden-brown, convex, bossed, stem brown, thickening upwards. — Batsch. 21.

Roots of hazels, November, Rare.

Clapham Park Wood.

Obs:

Gills flesh-colored.

1016. Small trumpet Agaric. — *A. buccinalis*.

Gills white, cap funnel-shaped, stem uniform in color, very short. — E. Fungi. 107.

On Moss, Autumn, Rare.

Flitton, — Amphill.

1017. Flat Agaric. — *A. compressus*.

Gills white, fleshy, few, 2 or 8 in a set, cap brown, irregular, stem whitish, flattened. — E. Fungi. 66.

Fir Groves, October, Common.

Obs:

Stem leaning, gills brownish-white.

1018. Shield Agaric. — *A. clypeatus*.

Cap hemispherical, clammy, pointed, gills ashy-black, stem long, white, cylindrical. — (Sibthorpe.) Bolton. 57.

Q q 2

Plantations,

XXIV. Mushrooms. Crypt. Fungi.

Plantations, September, Rare.

*Clophill, — Warden.**Obs :**Cap pale-brown.*1019. *Waxen Agaric.* — *A. ceraceus.*

Gills pale-yellow, in pairs, cap deep-yellow, hemispherical, smooth, stem deep-yellow, cylindrical. — *E. Fungi.* 20.

Dry Pastures, September, Rare.

*Thurleigh, — Aspley.*1020. *Cinnamon Agaric.* — *A. cinnamomeus.*

Cap dirty-yellow, gills yellowish-red. — *Bol-*
ton. 150. *E. Fungi.* 205.

Fir Groves, August, Rare.

*Amptbill.**Obs :**The Linnean character of this plant is admirable.*1021. *Shaggy Agaric.* — *A. floccosus.*

E. Fungi. 284. Clustered, cap and stem tufted with yellow-
ish-brown scales. — *Curtis.* IV. 71.
Stumps, August, Rare.

*Between Bedford & Elstow.**Obs :**Stem first prostrate, then ascending.*1022. *Equilateral Agaric.* — *A. simiputris.*

Cap pale-brown, bluntly conical, gills nearly black, stem whitish, with an irregular ring towards the top. — *Bulliard.* 66.

Pastures,

XXIV. Mushrooms. Crypt. Fungi.

Pastures, all the year, Common.

Obs:

The upper part of the stem bearing the traces of the fugitive ring determines this plant.

1023. Clustered Agaric. — *A. fascicularis.*

Cap convex, yellow, gills greenish, stem yellow. — (*Hudson.*) *Bolton.* 29. *E. F. Engg.* 285.

— Stumps, Autumn, Common.

Obs:

Stem leaning, rarely upright.

1024. Moss Agaric. — *A. Hypni.*

Cup with broad plaits at the brim, bossed, gills 3 in a set, inner ones fixed to the stem, all rust-colored, stem strap-shaped, slender, hollow within, with loose hairs. — *Batsch.* 96.

On Moss, October, Common.

Obs:

Cap conical, stem rather leaning.

1025. Lean Agaric. — *A. macer.*

Cap white, rather leather-like, thin, gills diluted-pink, stem white, hollow, splitting. — *Bulliard.* 518. *F.*

Shady Woods, October, Common.

Obs:

Cap almost transparent.

1026. Honied Agaric. — *A. mellitus.*

Gills

XXIV. Mushrooms. Crypt. VI Fungi.

Gills pale-brown, 4 in a set, cap pale-buff, center deeper, rather conical: stem whitish, crooked, curtain fugacious. — (Withering.) Schaeffer. 45.

Fir Plantations, September, Common.

1027. Parrot Agaric. — *A. psittacinus.*

Cap green and yellow-orange, bossed, edge waved, gills and stem of the same color. — E. Fungi. 82.

Parks, October, Rare.

Amphill Park.

1028. Hemispherical Agaric. — *A. semi-globatus.*

Cap hemispherical, straw-colored, clammy, gills horizontal, ring indistinct. — Curtis.

III. 69. E. Fungi. 248.

Pastures, April, October, Common.

Obs:

Cap pale brimstone, gills blackish.

1029. Congregated Agaric. — *A. striatus.*

Gills white, greyish at the edge, 2 or 4 in a set, cap conical, brown-buff, sides grooved, stem whitish, smooth. — E. Fungi. 166.

Willow stumps, Autumn, Common.

1030. Cap of Liberty Agaric. — *A. tener.*

Cap reddish, rather conical, deeper red at the brim, gills reddish, 4 in a set, dipping below the edge of the cap, stem reddish, smooth. — E. Fungi. 33. excellent.

Pastures,

* 1032. Striped Agaric. *A. polygrammus*.
Gills watery purple, 2 in a set, cap
conical, stem cylindrical, rather stiff
Bulliard. 395. S. Fungi. 222.
Stumps and amongst moss, Autumn Common.
Obs.

Stem distinctly striped, cap splitting
and turning up, a dirty brown, whitish
at the edge —

* 1032.
~~Small Stalked Agaric~~ *A. gibbiculatus*
Gills white, 2 in a set, cap bluntly
conical, not scored, stem cylindrical
spiral. S. Fungi. 165.

* 1032. Tentaculated Agaric. *A. nutans*.
Gills white, 2 in a set, cap convex,
nodding, stem ascending and tapering
upwards — S. Fungi. 92.

Branches of trees. Autumn. Rare
Son Hill.

Obs.

This little parasite is chiefly distinguished
from the *A. varius* by the sharp teeth
at the edge of the caps, which in its
more advanced state form a regular
undulation, not unlike the border
of the second wings in some of the

303

XXIV. Mushrooms. Crypt. Fungi.

Pastures, October, Common.

Obs.

This plant in decay resembles the cap of Liberty.

1031. Trembling Agaric. — *A. umbelliferus*.

Cap plaited, membranaceous, gills broader
at the base. — Bulliard. 519. 1. A.

Dead leaves, October, Common.

Obs.

Plant slender, very fine, trembling very much.

1032. Variable Agaric. — *A. varius*.

Gills white, few, 2 or 4 in a set, cap conical,
scored, stem cylindrical, stiff. — E. Fungi.
165. excellent.

Stumps, November, Common.

Obs.

Brim indistinctly scored.

1033. Wood Agaric. — *A. zylophilus*.

Cap cracking, pale-brown, deeper in the center,
edge turning up, gills brown, stem
leaning, of the same color. — E. Fungi. 167.

Fallen branches, August, Rare.

Clapham between Woods.

* * * * * On a stem, hollow and loose.

1034. Garlick Agaric. — *A. alliaceus*.

Cap open, half transparent, gills straw-color,
stem woolly half way up. — Bulliard. 158.
E. Fungi. 84?

Woods,

Butterflies in the Linnaean Clap
Lyrapholis —

XXIV. Mushrooms, no Crypt. Fungi.

Woods, September, Rare,

Silsœus.

*Obs:**Wool not yellow as in the peronatus, but an unpleasant black.*1035. *Watery Agaric.* — *A. aquosus.**Cap rather conical, gills whitish-grey, stem brownish, cylindrical. — Bulliard. (no number.)*

Rotten wood, April, Rare.

Eversholt.

*Obs:**Cap towards it's center and brims brown, intermediate space whitish.*1036. *Mottled Agaric.* — *A. clypeolarius.**Cap convex, branny, variegated, gills white, thin, 4 in a set, irregular, stem cylindrical. — E. Fungi. 14.*

Grassy spots, August, Rare.

Husborne-Crawley.

*Obs:**Stem bulbous at the base, gills ash-colored.*1037. *Cylindrical Agaric.* — *A. cylindricus.**Cap white, cylindrical, shaggy, gills purplish, uniform, stem cylindrical, white. — (Withering.) Curtis. II. 72. E. Fungi. 189.*

Way-sides, October, Common.

*Obs:*1035. *Small Nodding Agaric.* — *A. acicula**Cap conical, gills watery-grey, Stem pinkish, cylindrical. — E. Fungi. 282.*

Trees in Woods, on the bark December. Common.

*Obs:**Cap sharply conical, Stem zigzag, leaning. —*

XXIV. Mushrooms. Crypt. Fungi.

*Obs:**Cap cylindrico-oval.*1038. *Fleshy Agaric.* — *A. carnosus.*

Gills white, crowded, 4 or 8 in a set, cap convex, dirty-white, with reddish blotches and center reddish, stem dirty-white, blotched, ring none. — Curtis. V. 53. E. Fungi. 246.

Fir Groves, September, Common.

1039. *Wheel-like Agaric.* — *A. Rotula.*

Gills uniform, fixed to a collar surrounding the stem, cap white, skinny, dimpled; stem white above, black below. — E. Fungi. 95.

Dry sticks, August, Common.

1040. *Leaf Agaric.* — *A. dryophyllus.*

Gills pale-buff, 4 in a set, cap dead-white heightened to a pale-brown in the center, stem reddish at the base, whitish and tapering upwards. — E. Fungi. 127.

Shady woods, September, Common.

1041. *Wizard's Agaric.* — *A. hariolorum.*

Cap rather bossed, often horizontal, gills pale-pink, stem of the same color, fibrous at the base. — Bulliard. 56.

Fir Groves, Autumn, Rare.

*Amptbill, — Warden.**Obs :**Boss of the cap central, encircled with black.*

R r

Bald

XXIV. Mushrooms. Crypt. Fungi.

1042. *Bald Agaric.* — *A. impuber.*

Cap brown, yellow at the edge, gills 4 in a set, ochry-brown, stem purplish at the base, tapering downwards. — E. Fungi. 125.

Shady woods, October, Rare.

Lawn Wood.

1043. *Weeping Agaric.* — *A. lachrymabundus.*

Gills dull-red, broad, numerous, 2 or 4 in a set, pileus dirty-brown, conical, woolly, stem dirty-white. — (Withering.) E. Fungi. 41.

Woods, lanes, October, Common.

1044. *Short-lived Agaric.* — *A. momentaneus.*

Cap nearly conical, white, branny, gills ashy-black, stem white. — (Sibthorpe.) Bulliard. 68.

Pastures, October, Common.

1045. *Bordered Agaric.* — *A. marginatus.*

Cap ochry, brim very narrow, stem between livid and ochry, snow-white at the base, gills between rust-colored and ochry. — Batsch. 207.

Hedges, October, Common.

Obs.:

The point of the cap appears lopped or flattened.

1046. *Puckered Agaric.* — *A. ovatus.*

Cap

XXIV. Mushrooms. Crypt. Fungi.

Cap ovate, somewhat plaited, stem bare, tapering at the base, roundish, gills crowded, nearly uniting. — Curtis. II. 71. **E. Fungi. 188.**

Grassy spots, October, Common.

Obs :

Cap puckered, gills grey, when ripe dissolving.

1047. *Tall Agaric.* — *A. procerus.*

Cap bell-shaped, nearly brown, scaly, gills whitish, stem bulbous, with a ring. — *(Light-foot.)* Curtis. IV. 69. **E. Fungi. 190.**

Fir Groves, October, Common.

Obs :

Stem thickening at the base and ending in an uncommon large bulb.

1048. *Pill-shaped Agaric.* — *A. piluliformis.* **S D.**

Cap round, brown, gills whitish, stem slanting, of the same color. — Bulliard. 112.

Amongst moss, November, Rare.

Southill.

Obs :

Plant growing in clusters on living Mosses.

1049. *Plaited Agaric.* — *A. plicatilis.*

Cap bell-shaped or flat, mouse-colored, transparent, plaited. — Curtis. III. 70.

Grassy spots, November, Common.

R r 2

Obs :

XXIV. Mushrooms. Crypt. Fungi.

Obs:

Edge of the cap so deeply channelled as to deserve the name it bears.

1050. *Half-egg Agaric.* — *A. semi-ovatus.*

Gills brown-grey to black, 2 or 4 in a set, cap light-brown, smooth, half-egg-shaped, stem cylindrical, white. — E. Fungi. 131.

Pastures, June, Common.

1051. *Yellow-white Agaric.* — *A. titubans.*

Cap half-ovate, yellow, grooved towards the margin, gills pale-pink, stem very brittle, yellow. — E. Fungi. 128. excellent.

Obs:

Stem very long and thin.

1052. *Velvet-stemmed Agaric.* — *A. velutipes.*

Clustered, cap flattish, tawny, stem bare with a fine sooty woolliness at the base.—Curtis.

IV. 70.

Fallen elms, October, Common.

Obs:

Cap orange-tawny, stem velvety.

* * * * * Stems lateral.

1053. *Styptic Agaric.* — *A. stypticus.*

Cap kidney-shaped, ochry, gills of the same color. — E. Fungi. 109.

Stumps,

XXIV. Mushrooms. *Crypt. Fungi.*

Stumps of trees, October, Rare.

Clophill.

Obs:

Plant fan-shaped.

1054. *Lateral Agaric.* — *A. lateralis.*

Nearly on a stem, smooth, lead-colored, gills nearly single.

On moss, October, Rare.

Warden, — Clophill.

Obs:

Whole plant between tan-leather and grey.

1055. *Oyster Agaric.* — *A. ostreatus.*

Between bundled and tiled, cap ash-colored, obversely ovate, edge rolled in, gills whitish, decurrent, somewhat branching at the base.

Curtis. III. 71. E. Fungi. 241.

Trees, Autumn, Rare.

Kempston, — Clapham.

Obs:

Gills towards the edge diluted-pink, in other parts grey.

1056. *Shell Agaric.* — *A. conchatus.*

Gills ochry, cap of the same color, splitting, stem lateral. — *Bulliard. 298.*

Fissures of trees, November, Rare.

Thurleigh.

Obs:

Like sea-shells in it's young state.

* * * * * Stemless.

XXIV. Mushrooms, &c. Crypt. Fungi.

* * * * * Stemless.

1057. *Soft Agaric.* — *A. mollis.*

Between gelatinous and membranaceous, cap irregular, whitish, gills brimstone-colored.—
E. Fungi. 98. excellent.

Fallen elms, December, Common.

1058. *Sessile Agaric.* — *A. sessilis.*

Cap half-lopped, reversed, gills snuff-colored, rim bent inwards.—E. Fungi. 97. excellent.

Hazel sticks in woods, November, Common.

Obs:

Cap convex, white, gills much crowded.

1059. *Dark-grey Agaric.* — *A. applicatus.*

E. Fungi. 305. Turned inwards, round, ashy-black, gills touching in the center, whitish-azure.—
(Dickson.) Batsch. 125.

Posts, Autumn, Rare.

Clophill, — Silsoe.

1060. *Birch Agaric.* — *A. coriaceus.*

E. Fungi. 182. Proliferous, cap with concentric zones, leather-like, woolly, gills ochry-brown, branching.—Bulliard. 394. excellent.

Woods, October, Rare.

Clapham Park Wood.

Obs:

Cap perforated by grass, sticks, &c.

Oak

add to obs. 1061.
appears to be the connecting Link
between the Genera, *Agaricus*,
Boletus and *Hydnus*.

XXIV. Mushrooms. Crypt. Fungi.

1061. Oak Agaric. — *A. quercinus*.

Gills labyrinth-shaped. — Bulliard. 352.

Fallen elms, Winter, Rare. E. Fungz. 101.
Aspley.

Obs :

*This plant first adheres to the trunks of trees horizontally,
afterwards becomes reversed and grows by its cap.*1062. Petal Agaric. — *A. petalodes*, S.D.Gills brimstone, cap shaped like a petal. —
Bulliard. 226.

Old stumps, December, Rare.

Clapham Park Wood.

Obs :

*Clustered, shaped like the petal of a tulip.**Boletus. BOLETUS. Gen. Pl. 1677.*

— Horizontal, porous underneath, pores united.

† On a stem, stem central.

1063. Sponge Boletus. — *B. bovinus*.Cap olive-brown, pores pale-tawny, unequal,
stem dirty-white. — (Sibthorpe.) Bulliard.

60. E. Fungz. V5.

Woods, October, Common.

Obs :

Stem without a ring.

Olive

XXIV. Mushrooms. Crypt. Fungi.

1064. *Olive Boletus*. — *B. olivaceus*.

Cap olive-colored, bent downwards, flesh changing, tubes minute, yellow. — (*Sibthorpe*.) Schaeffer. 105.

Pastures, woods, October, Common.

Obs:

Cap chesnut-colored, pores very minute.

1065. *Azure Boletus*. — *B. cyanescens*.

Cap fawn-colored, fleshy, flesh changing, stem thick, short, tuberous. — (*Sibthorpe*.) Bulliard. 369.

Woods, September, Rare,

Aspley, — Clapham between Woods.

Obs:

Flesh when bruised changes from white to blue.

1066. *Circular Boletus*. — *B. nummularius*.

Cap horny, varying from brown to chesnut, pores decurrent, stem black at the base. — E. Fungi, 89.

Hazel sticks, Autumn, Rare.

Clapham Park Wood.

Obs:

Cap marked with dark stripes.

1067. *Clustered Boletus*. — *B. gregarius*. **S.D.**

Tubes yellow, pores oblong, unequal, cap thin, flattish, chesnut-colored, stem pink at the base. — Flora Danica. 1018.

Fir

XXIV. Mushrooms. Crypt. Fungi.

Fir Groves, October, Rare.

*Warden.*1068. *Common Boletus.* — *B. luteus.*

Cushion-shaped, rather clammy, pores rounded, convex, very yellow, ring permanent. — Schæffer. 114.

Woods, Autumn, Common.

*Obs :**Pores easily separated.*1069. *Bronze Boletus.* — *B. æreus.*

Cap brown, hemispherical, tubes short, sulphureous, stem marked with purple veins. — Bulliard. 385.

Fir Groves, August, Rare.

*Warden.**Obs :**Stem with net-work veins.*1070. *Pepper Boletus.* — *B. piperatus.*

Stem rather slender, cap yellow, flesh sulphur-colored, firm, pores rather long, flame-colored. — E. Fungi. 34.

Fir Groves, September, Rare.

*Warden.**Obs :*

This Boletus, when chewed in the mouth, leaves a sharp peppery taste.

S s

† † Stems

XXIV. Mushrooms. Crypt. Fungi.

† † Stems lateral.

1071. *Oblique Boletus*. — *B. obliquatus*. S.D.

Cap oblique, marked with concentric zones,
stem short, lateral. — Bulliard. 459. L.
Stumps, December, Rare.

Clapham Park Wood.

1072. *Scaly Boletus*. — *B. squamosus*.

Almost stemless, flattish, between ragged and
scaly, yellowish, pores 4-square. — Schaeffer.

101. 102.

Stumps, August, Common.

Obs:

Stem thinnest at bottom, pores spacious.

1073. *Slipper Boletus*. — *B. elegans*.

Cap ear-shaped, smooth, pores small, irregular, stem full, black at the base. — (Relhan.) Bulliard. 360.

Willows, May, Rare.

Duck Mill.

Obs:

Cap very thin and tough.

1074. *Tiled Boletus*. — *B. frondosus*.

Almost stemless, tiled, pores decurrent, stem
black at the base, very irregular. — E. Fun-
gi. 87.

Stumps,

- * 1071. *Red: brown Boletus*. *B. brevis*.

Cap hemispherical, red: brown, pores
sinuous, very long and pendulous
stem scarcely any. — E. Fungi 191.

Fir groves - October - rare -
Amphill.

Obs:

This may possibly be a variety
of the plant quoted above, as Mr.
S. represents his *Boletus* becoming
woody and hard in a dry state.
My specimens were deliquescent, I
might almost say, gelatinous —

XXIV. Mushrooms. Crypt. Fungi.

Stumps, Autumn, Rare.

Maulden,—Renhold.

Obs:

Cap fawn-colored, pores whitish.

1075. Varnished Boletus. — *B. rugosus.*

Pores white, very small, cap crimson, varnished, stem knotty, slanting. — E. Fungi.

134. excellent.

Hazel stumps, Autumn, Rare.

Clapham Park Wood.

Obs:

Stem darker than the cap.

† † † Stemless.

1076. Spongy Boletus. — *B. suberosus. S.D.*

Cushion-shaped, smooth, pores pointed, irregular. — Bolton. 162. E. Fungi. 280.

Ash trees, November, Rare.

Potton,—Everton.

Obs:

Cap rather scaly.

1077. Crumb of bread Boletus.—*B. medulla panis.*

Crust-like, white, spreading, irregular. — Jacquin. Misc. Aust. I. 11.

Decayed sticks, August, Common.

Obs:

Cap reversed, adhering to rotten wood.

S s 2

Willow

XXIV. Mushrooms. Crypt. Fungi.

1078. Willow Boletus. — B. salicinus.

Somewhat leather-like, soft, sitting, thin, semicircular, somewhat whitish, smooth, pores very short, whitish-red. — Bulliard.

433. i. E. Fungi. 227.

Willows, Winter, Common.

*Obs:**Smells like the bloom of May.*

1079. Liver Boletus. — B. hepaticus.

Convex, nearly round, red, smooth, pores round, yellow. — E. Fungi. 58.

Stumps, September, Common.

*Obs:**Exactly like a piece of raw bullock's liver.*

1080. Sweet-smelling Boletus. — B. suaveolens.

Smooth above, growing on the willow. —

Bulliard. 310. E. Fungi. 228.

Stumps, Autumn, Rare.

*Fenlake.**Obs:**Pores odoriferous, white or straw-colored.*

1081. Dry-rot Boletus. — B. lachrymans.

Leather-like, half-oval, orange, between wrinkled and reticulated, border broad, snow-white, arched. — Jacquin Misc. Aust.

Vol. II. viii. 2. E. Fungi. 113.

Rotting

1078.

Hybrid Boletus. B. hybrida

Somewhat leather-like soft, cushion-shaped, pores pale yellow, decurrent.

8. Fungi. 289.

*Rotten Wood, Autumn Rose.**Goldington, Slophill.*

*1082.

Whitish Boletus. B. albidus.

Sitting, cap tender, white pores very small, distinct.

E. Fungi. 226.

²²⁶
At the edges of pannels in old Gates - winter, rare.

Bedford.

Obs:

Pores often irregular.

*1083.

Rusted Boletus B. ferrugineus.

Cap rust colored at the base pores white, border of the cap of the same color -

Fir Stumps - Autumn Rare Amphitheatre Plantations.

Obs:

Unlike the versicolor in structure for it has neither stripes nor concentric ridges -

317

XXIV. Mushrooms. Crypt. Fungi.

Rotting wainscot boards, Autumn, too common.

Obs:

Border bent back, whitish, dotted, cushion-shaped.

1082. Striped Boletus. — B. versicolor.

Ridges discolored, pores white. — Schaeffer.

267. 268. 269. E. Fungi. 229.

Trees, October, Common.

Obs:

Plant extremely thin.

1083. Self-colored Boletus. — B. unicolor.

Leather-like, thin, semicircular, with woolly zones above, pores ash-colored. — Bulliard. 408. - 501. 3.

Hedges on stumps, April, Common.

Obs:

Pores not a quarter of an inch long.

1084. Sulphur Boletus. — B. sulphureus.

Pores brimstone, cap a bright Aurora, streaked. — E. Fungi. 135.

Old stumps, October, Rare.

Hawthorn.

Obs:

Plant much tiled, color soon fading.

1085. Bristly Boletus. — B. hispidus.

Pores yellowish-red, fringed, cap red-brown, rough with bristly hairs. — Bulliard. 210.

Ash-trees,

XXIV. Mushrooms. Crypt. Fungi.

Ash-trees, Autumn, Common.

Obs :

Younger specimens blood-red, old ones tawny.

1086. Touch-wood Boletus. — *B. igniarius*.

Cushion-like, smooth, pores very fine. — E. Fungi. 132.

Trees, Autumn, Common.

Obs :

Like a horse's hoof.

1087. Tinder Boletus. — *B. fomentarius*.

Pores sea-green, round, equal, cap cushion-like, blunt, unequal. — E. Fungi. 133.

Stumps, October, Rare.

Clapham.

Obs :

Thinner than the preceding.

1088. Fan-shaped Boletus. — *B. flabelliformis*.

Tiled, pores dark-purple, cap pale-flesh-colored. — ? Batsch. 266. E. Fungi. 231.

Stumps, October, Common.

Obs :

Pores lead-colored.

Hydnium. HYDNUM. Gen. Pl. 1676.

Mushroom horizontal, hedge-hogged underneath.

‡ On

* 1088 - Bald Boletus. *B. impuber.*

Cap rusty-brown, smooth, border velvety, pores darker, decurrent, of different lengths.

E. Fungi. 195.

Young willows in damp places, Winter, Rare.

Penholt, by a drain near the Wood* 1088. Narrow pored Boletus. *B. angustans*

Cap hemispherical, whitish, often red, border turned in pores yellowish, long and narrow. — E. Fungi. 193.

Bromham garden on trees.
April. 1802.* 1088 Comb-shaped Boletus. *B. pectinatus*.

Tubes yellowish white, inclining, cap reversed and sitting down —

Bolton. 74.

Ray. tab. I. 5.

Joists in cellars - Winter, Rare.

Goldington.

Obs:

This has no pretension to be ranked with the agarics, the pores towards the border where they run into the cap are often very clear and distinct.

* 1080.

Tanleather Boletus. *B. pellionius*.

Sized, pores yellowish, cap the
colour of tanned Leather.

9. Fungi. 230.

Alder and Birch Stumps, winter, rare
Amphill.

Obs.

The resemblance between this and no
1088, is not so close as some Botanists
have suspected, the pores of this
never turn to a compleat purple
as in *B. flabelliformis*. —



XXIV. Mushrooms. Crypt. Fungi.

‡ On a stem.

1089. Ear-picker Hydnum.—*H. auriscalpium.*

Cap semicircular.—Curtis. III. 68.

Decayed cones of firs, September, Common.

Obs :

Cap rather kidney-shaped.

1090. Tiled Hydnum.—*H. imbricatum.*

Cap convex, scaly.—(See our fig. No. 4.)

Fir Plantations, October, Rare.

Warden.

Obs :

Cylinders pinkish-brown, decurrent.

1091. Incorporated Hydnum.—*H. sublamellatum.*

Snow-white, cylinders shaped like gills, caps incorporating.—E. Fungi. 112. excellent.

Woods, October, very Rare.

Clapham Park Wood.

Obs :

I think myself extremely fortunate in being the first discoverer of this *Hydnum* as a native of England.

† † -Stemless-

1092. Jupiter's beard Hydnum.—*H. barba Jovis.*

Tawny, membranaceous, spreading, the ends of the prickles branched.—Bulliard. 481. 2.

Bases of birchen stumps, October, Rare.

Southill Plantations.

Turban-top.

Before 1092

Jupiter's beard Hydnum
inest † † Stemless.



XXIV. Mushrooms. Crypt. Fungi.

Turban-top. HELVELLA, Gen. Pl. 1677.

Mushroom top-shaped, smooth above and underneath.

1093. *Curled Helvella.* — H. mitra.

Cap bent down, growing together, lobed, irregular. — E. Fungi. 39. excellent.

Woods, October, Rare.

Houghton-conquest, — Clapham between Woods.

Obs :

Stem cellular, waxen.

1094. *Relhan's Helvella.* — H. Relhani. S.D.

Cap conical, brown, stem hollow. — E. Fungi. II. excellent.

Pastures, May, Rare.

Medbury.

Obs :

Cap silky above, brim with a silky fringe.

1095. *Agaric-shaped Helvella.* — H. agariciformis.

Very small, white, stem thread-shaped, solid, cap thin, smooth above and underneath. — (Relhan.) E. Fungi. 57.

Under moss at the roots of trees, October, Common.

Obs :

Stem sometimes branched and proliferous.

Jelly-like

XXIV. Mushrooms. Crypt. Fungi.

1096. *Jelly-like Helvella*. — *H. gelatinosa*.

Stem hollow, swelling at the base, cap arched, smooth above, jelly-like within, waved underneath. — (*Bulliard.*) *E. Fungi.* 70.

Woods, October, Rare.

Southill, — *Silsoe*.

Obs :

Upper part of the stem compressed.

1097. *Battledore Helvella*. — *H. spatulata*.

Cap much waved, flat, nicked at the top, stem hollow. — *E. Fungi.* 35.

Fir Groves, September, Rare.

Ampthill.

Obs :

Plant growing in clusters and extremely variable in color.

1098. *Eatable Helvella*. — *H. esculenta*.

Cap ovate, with cells, stem bare, wrinkled. — *E. Fungi.* 51.

Pastures, April, Common.

Obs :

Cap sometimes conical, oftener semi-ovate.

Auricularia. AURICULARIA. Bulliard.

Mushroom ear-shaped, sitting, gradually reversed, discharging seeds from what was the upper, but which is now become the lower surface.

T t

Gelatinous

XXIV. Mushrooms. Crypt. Fungi.

1099. *Gelatinous Auricularia*. — A. tremelloides.

Between jelly-like and gristly, fringed above with zones, pitted or plaited underneath. — Bulliard. 290.

E. Fungi. 290. Gate-posts, January, Common.

Obs:

Outside greyish-brown, inside purple and pitted.

1100. *Buff Auricularia*. — A. papyrina.

Membranaceous, soft, woolly above with zones, at first smooth underneath, afterwards pitted. — Bulliard. 402.

Trunks of trees, October, Rare.

Sheerhatch Wood.

Obs:

Woolly and fringed at the edges.

1101. *Woolly Auricularia*. — A. reflexa.

Leathery, thin, woolly above, smooth underneath. — E. Fungi. 27.

Stumps, December, Common.

Obs:

Cap woolly, with zones.

1102. *Rusty Auricularia*. — A. ferruginea.

Leathery, brownish-rust-colored, velvety above, with concentric scores, smooth, pimply underneath. — E. Fungi. 26.

Old pales, January, Common.

Tobacco

* 1100. *Powdery Auricularia*. *A. pulvoculata*.
Membranaceous, tender, with concentric zones, proliferous, smooth underneath.
P. Fungi. 214.

Damp walls, on the plaster
winter Rare.



Bedford.

Obs:

~~Having seen the whole process of~~
this curious parasite is extremely singular. It rises in the places I have mentioned above, like a small mucilaginous spot, which gradually enlarges and in its second state gives the promise of a *Boletus* laetiporus. The Botanists expectations are at length dispelled by the peeling off of the uppermost edge from the wall, when the zones begin to rise, it separates from the place where it lately grew, and the fructifying powder breaking forth, it puts on the singular appearance, represented in the plate above quoted —

XXIV. Mushrooms. Crypt. Fungi.

1103. *Tobacco Auricularia*. — *A. tabacina*.

Rust-colored, zigzag, thin, with a yellow border. — (*Sibthorpe*.) E. Fungi. 25.

Branches of trees, February, Rare.

Clophill, — *Amptbill*.

Obs:

Lower surface veined, bare, wrinkled.

1104. *Stick Auricularia*. — *A. corticalis*.

Leathery, thin, between flesh-colored and brownish underneath. — Bulliard. 436. 1.

Dead sticks, August, Common.

Obs:

Greyish-brown, pimply, pitted like a Liverwort.

1105. *Robe Auricularia*. — *A. phylacteris*.

Thin, membranaceous, soft, smooth, folded at the base. — Bulliard. 436. 2.

At the foot of large trees in woods, January, Rare.

Sheerbatch Wood.

Obs:

Pale-straw-colored.

Morel. PHALLUS.

Mushroom with a cellular cap, open at the end, springing from a wrapper.

1106. *Stinking Morel*. — *P. impudicus*.

In a wrapper, on a stem, cap cellular. — Curtis. III. 72.

T t 2

Woods,

XXIV. Mushrooms. Crypt. Fungi.

Woods, Hedges, September, Rare.
 Silsoe, — Market-street. Obs:
 Cap sometimes with two openings.

1107. Dog's Morel. — *P. caninus*.

In a wrapper, on a stem, cap wrinkled, conical, mouth closed. — Curtis. IV. 73.

Woods, August, very Rare.

Silsoe. Obs:

Cap covered with a greenish liquor, red underneath.

Peziza. PEZIZA. Gen. Pl. 1680.

Concave, seeds emitted only from the upper surface.

1108. *Cornucopia Peziza. — P. cornucopioides*.

Funnel-shaped, disc expanding, waved, dotted. — (See our fig. No. 5.)

Woods, September, Rare.

Silsoe. *Aspley Wood.*

Obs: Oct. 7. 1805.

Plant sometimes solitary, oftener in clusters. I am indebted for the discovery of this very rare plant to the researches and assiduity of Mrs. Abbot.

1109. *Yellow-brown Peziza. — P. ochroleuca*.

Stem black at the bottom, cap funnel-shaped, dirty-yellow within. — E. Fungi. 115.

Woods,



XXIV. Mushrooms. Crypt. Fungi.

Woods on decayed sticks, Winter, Rare.

Clapham Park Wood.

Obs:

Substance leather-like, horny in decay.

1110. *Snowy Peziza.* — *P. nivea.*

White as snow, funnel-shaped, on a very short stem. — E. Fungi. 65.

Under old stumps, August, Common.

Obs:

Young specimens clothed with a whitish wool.

1111. *Dotted Peziza.* — *P. punctata.*

Top-shaped, lopped, the disc of the cap dotted. — E. Fungi. 54.

Cow-dung, October, Common.

Obs:

The black dots not always visible.

1112. *Crimson Peziza.* — *P. epidendra.*

Bell-shaped, brittle, wax-colored without, crimson within. — E. Fungi. 13.

Hedges, October, Rare.

Clapham, — Southill.

1113. *Goblet Peziza.* — *P. acetabulum.*

On a short stem, cap goblet-shaped, angular without, with branching veins. — E. Fungi. 59.

Woods,

XXIV. Mushrooms. Crypt. Fungi.

Woods, May, Rare.

Ampthill, — Aspley.

Obs :

Stem cellular.

1114. Tuberous Peziza. — *P. tuberosa*.

Cap nearly bell-shaped, brown without, paler within. — E. Fungi. 63.

Woods, Autumn, Rare.

Renhold Wood.

Obs :

Stem buried in the earth, growing at the base to a blackish tuberous substance.

1115. Scarlet Peziza. — *P. coccinea*.

Growing on the ground, between orange and scarlet, cup expanding. — E. Fungi. 78.

Woods, September, Rare.

Clapham Springs.

Brought together with the *Bryum trifarium* by Mr. T. KIDMAN of Bedford.

Obs :

Fine scarlet within, leather-colored without.

1116. Smooth Peziza. — *P. cyathoides*.

Glass-shaped, border blunt, upright. — Ray's Synopsis. XXIV. 4.

Branches of trees, August, Common.

Little Cup

* 1116. Golden Peziza. — *P. aurea*.Stem short, yellow, cap umbrella-like
gold coloured withering.

Fungi. 150.

Horizontally on wet stumps, Woods
November, Rare

Oakley little wood.

Obs : Cap delicate, trembling, and semi-transparent.

* 1117 Rush Peziza. P. junci.

Black, gelatinous, rather conveved, sitting.

On stems of rotten Juncis pools - October - Rare.
Amphill.

* 1117 Leaning Peziza. P. inclinans

White, slender stemmed, head leaning.
^{November}

Rotten stumps in woods, Rare
Clapham between woods

* 1117. Lobed Peziza. P. sinuata.

Emarginate, colour of Beeswax
Smooth, horny -

Rotten stumps in woods. November, rare

Clapham between woods.

(Cross-shaped border bottom, orange)

Branches of trees, August, Common

XXIV. Mushrooms. Crypt. Fungi.

1117. Little Cup Peziza. — P. calyculus.

On a stem, convex, hemispherical, expanding, stem rather lengthened out, strap-shaped, firm, distinctly inserted. — Batsch.

XII. 57. E. Fungi. W6.

Hedge bottoms, August, Common.

Obs :

Whole plant yellowish-brown.

1118. Hornbeam Peziza. — P. fructigena.

Slender, tapering, brittle, smooth, ending in a slender stem, cup scolloped. — Bulliard.

228. E. Fungi. W7.

On acorns, October, Common.

1119. Eyelet Peziza. — P. marginata.

Concave, brownish, border flat, rather scolloped, snow-white. — (Relhan.) E. Fungi. 16. excellent.

On decayed sticks, October, Common.

Obs :

Like the eyelet holes of a Lady's stays.

1120. Wreathed Peziza. — P. cochlearia.

Top-shaped, wreathed. — E. Fungi. 5.
Woods, April, Common.

Obs :

Above dark-brown, beneath whitish.

XXIV. Mushrooms. Crypt. Fungi.

1121. Cow Peziza. — *P. stercoraria*.

Sitting, nearly straw-colored, yellow, cup scolloped, prominent with blackish points, rough with hair. — (*Sibthorpe.*) E. Fungi. 18.

Cow-dung, September, Common.

1122. Orange Peziza. — *P. fulva*.

Flat, orange, border convex, smooth. — Bui- liard. 438. 3.

Cow-dung, October, Common.

Obs:

Whole plant dull-orange.

1123. Eyelash Peziza. — *P. scutellata*.

Flat, border convex, hairy. — E. Fungi. 24.

Bogs, October, Common.

Obs:

Disc scarlet, border fringed.

1124. Vermillion Peziza. — *P. punicea*.

Sitting, not shining, scarlet above, paler underneath, border cushion-shaped. — Batsch. XXXIX. 220.

Mosses, October, Common.

Obs:

Plant often roundish, between an orange and a scarlet color.

1125. Nipple Peziza. — *P. papillaria*.

Very

Small, sitting, reddish, disc-like, rough with small protuberances, surface

wolly, cap lobed. — Batsch. 220.

Common, October, Common.

Fr. yellowish brown, surface

smooth. — Batsch. 220.

Rotten stumps, February, Common.

Fr. greyish brown. — Batsch. 220.

Smooth, pale brown, surface

smooth. — Batsch. 220.

Mosses in woods, October, Common.

Fr. yellowish brown. — Batsch. 220.

Smooth, pale brown, surface

smooth. — Batsch. 220.

Mosses in woods, October, Common.

Fr. yellowish brown. — Batsch. 220.

Smooth, pale brown, surface

smooth. — Batsch. 220.

Mosses in woods, October, Common.

Fr. yellowish brown. — Batsch. 220.

Smooth, pale brown, surface

smooth. — Batsch. 220.

Mosses in woods, October, Common.

Fr. yellowish brown. — Batsch. 220.

Smooth, pale brown, surface

XXIV. Mushrooms. Crypt. Fungi.

Very small, somewhat waxen, thickish, sitting, rough with small prominent points, rather woolly, cup scolloped. — (Bulliard.) E. Fungi. 177.

Rotten wood, October, Common.

Obs:

Center ash-colored.

1126. Two-colored Peziza. — P. bicolor.

On a very short stem, white underneath, cup orange-scarlet. — E. Fungi. 17. excellent.

Rotten sticks, February, Common.

1127. Hairy Peziza. — P. hispida.

Sitting, cup between scolloped and vesicular, outside brown, hairy, inside bluish-white, smooth. — E. Fungi. 147. excellent.

Mosses in woods, October, Common.

1128. Rasp Peziza. — P. leporina.

Stem short, cap between flesh-colored and yellow, turned in, like a rasp underneath. — E. Fungi. 79.

Woods, October, Rare.

Silsoe.

1129. Gelatinous Peziza. — P. tremelloidea. S.D.

At first sitting, like a Tremella, then running into a thickish stem, cup between flat and concave. — Bulliard. 410. 1.

U u

Stumps,

XXIV. Mushrooms. Crypt. Fungi.

Stumps, December, Common.

1130. *Variable Peziza*. — *P. polymorpha*.

Top-shaped, first hollow, then flat or convex, wrinkled on the outside. — Bulliard.
116. — 460. 1.

Stumps, October, Common.

1131. *Lentil Peziza*. — *P. lenticularis*.

Small, thick, brittle, smooth, sitting or top-shaped, smooth underneath, between shield and cup-shaped. — Bulliard. 300.

Stumps, October, Common.

1132. *Green Peziza*. — *P. viridis*.

Stemless, cap concave, green, border blackish. — Bolton. 109. 1.

Stumps, October, Rare.

Clophill.

1133. *Black Peziza*. — *P. atra*. **S.D.**

Stemless, concave, black. — (*Hudson.*)

Park pales, Autumn, Rare.

Ampibill.

1134. *Blue Peziza*. — *P. cærulea*. **S.D.**

Stemless, cap azure, fringed. — Bolton. 108. 2.

Stumps, October, Rare.

Sheerbatch Wood.

Lead-colored

1135. Downy Peziza. — *P. hydnoides.*

Downy, ash-colored within, without yellowish
white — *P. Fungi.* 178.

Bits of Sticks in Woods, Winter, Rare.

Penholt, Clapham Park and between Woods

Obs:

Plant smaller than the figure in Sowerby
but perfectly resembling it in every
other respect; if suffered to become dry,
the disc closes and the orifice is no
longer seen —

331

XXIV. Mushrooms. Crypt. Fungi.

1135. Lead-colored Peziza. — *P. cinerea.*

Sitting, ash-colored, bent back, border cur-
led. — *E. Fungi.* 64.

Under old stumps, November, Common.

1136. Bell-shaped Peziza. — *P. campanulata.*

Lead-colored, smooth, border expanding,
seeds smooth. — *E. Fungi.* 28.

Bean-fields, Autumn, Common.

1137. Scored Peziza. — *P. striata.*

Conical, brown, outwardly woolly, inwardly
fluted, seeds downy underneath, — *E. Fun-
gi.* 29.

Branches of dead firs, November, Rare.

Warden.

1138. Crucible Peziza. — *P. laevis.*

Dirty-yellow, smooth within, border upright,
seeds smooth. — (*Sibthorpe.*) *E. Fungi.* 30.

Branches of living firs on the ground, Octo-
ber, Rare.

Warden.

1139. Crowded Peziza. — *P. spadicea.* **SP.**

Sitting, almost stemless, unequal, shining a-
bove, rough beneath, border finely scol-
loped. — *Batsch.* 217.

Fallen elms, January, Rare.

Potton Wood.

U u . 2

Cone

XXIV. Mushrooms. Crypt. Fungi.

1140. *Cone Peziza*. — *P. pineti*. S.D.

Sitting, greyish-white, bent back, border ochry-brown. — Batsch. 140.

Cones of decayed firs, October, Rare.

Warden, — *Ampibill*.

Club-top. CLAVARIA. Gen. Pl. 1682.

Mushroom smooth, club-shaped, quite simple, emitting seeds from every part of the surface.

1141. *Hollow Clavaria*. — *C. pistillaris*.

Club-shaped, very entire, hollow, tapering at the base, point roundish. — Bulliard. 463. 1.

E. Fungi. 255

253. Heaths, September, Common.

1142. *Adder's-tongue Clavaria*. — *C. ophio-glossoides*.

Club-shaped, very entire, flattened, black, head furrowed. — E. Fungi. 83.

Heaths, October, Rare.

Warden, — *Clophill*.

1143. *Coral Clavaria*. — *C. coralloides*.

Branches crowded, very much branched, bluntnish at top, yellow. — Bulliard. 222.

Heaths, woods, November, Common.

1144. *Flat-topped Clavaria*. — *C. fastigiata*.

Branches,

* 1140. Jew's Pass - P. auricula.

Stump of a dead Elm at
the Friars found by Mr. Abbot
Sep: 16. 1800. —

XXIV. Mushrooms, Crypt. Fungi.

Branches crowded, very much branched, flat at top, straw-colored. — Bolton. 112. 2.

Woods, September, Rare.

Potton Wood.

1145. Pointed Clavaria. — C. muscoides.

Branches subdivided, pointed, unequal, yellow. — Schaeffer. 173. E. Fungi. 157.

Woods, October, Common.

1146. Fingered Clavaria. — C. digitata.

Branched, branches nearly sitting, oblong, points blunt. — E. Fungi. 69.

Door-posts, Autumn, Common.

1147. Flat-horned Clavaria. — C. Hypoxylon.

Branched, compressed, horned. — E. Fungi. 55.

Stumps, Winter, Common.

1148. Scarlet Clavaria. — C. coccinea.

Almost sitting, Scarlet, shaped like a Pizza. E. Fungi. 291.

Bark in Autumn. Common

Mouse-colored, branches coral-like, solid, thick and short, points depressed. — Bulliard. 354.

Heaths, September, Rare.

Warden, — Clophill.

1149. Cauliflower Clavaria. — C. anthocephala.

Leathery, growing in a turf, rust-colored, with fan-shaped lobes. — Bulliard. 452. 1.

E. Fungi. 156. Heaths,

XXIV. Mushrooms. Crypt. Fungi.

Heaths, Autumn, Rare.

Sandy Warren.

1150. Leathery Clavaria. — C. coriacea.

Leathery, softish, branches coral-like, depressed, scored, points fringed. — Bulliard.

452. 2.

Heaths, Commons, October, Rare.

Warden.—Beeston Leys.

1151. Thorny Clavaria. — C. aculeiformis.

Small, yellow, very brittle, single or double, end pointed. — E. Fungi. 40.

Stumps, January, Rare.

Timber-yards, Bedford.

1152. Twisted Clavaria. — C. gyrans.

Stem lengthened out, hair-like, ending in a small club, rather lengthened out, cylindrical, oblong at either end. — Batsch. 164.

Woods, October, Rare.

Clapham between Woods.

1153. Orange Clavaria. — C. militaris.

Club-shaped, very entire, head scaly. — E. Fungi. 60.

Bogs, October, very Rare.

Ampthill.

Trufle.

1150. Leathery Clavaria. C. epiphylla.
Club-shaped, very entire, head blunt,
Hollow, ochrey red, Stem yellowish white
8. Fungi. 243.

Peat holes & Woods, Autumn rare
Selsoe New Wood and Ampthill Heath.

1151. Forked Clavaria. C. dichotoma
Subterraneous or immersed, horny,
coral-like black, pointed.

8. Fungi. 248. Obs:

like fibres of trees or roots.

1151. Obs. Plant mostly clustered,
~~wax-like~~, points obtusely conical,
rather than thorn-like, surface
velvety—substances between wax-like and
glutinous. Elges' Gard. St. Mary's Parish

Dec? 17th 1790-

1150

Thorn like Clavaria. C. spinosa
Subterraneous, ragged, of an
equal lengths, pointed, dark
brown — Woods. Common.

8. Fungi. 299.
Obs:

Like a ragged tapel.

XXIV. Mushrooms. Crypt. Fungi.

VIXX

Trufle. TUBER. Bulliard. 73.Mushroom growing under ground, fleshy,
firm, containing the seeds within itself.1154. *Common Trufle.* — *T. cibarium.*Round, solid, warty, without a root. — Bul-
liard. 356. E. Fungi. 309.

Woods, pastures, Rare.

Renhold Wood, — Willington Pastures.

1155. *Rooted Trufle.* — *T. radicatum.*Roundish, compressed, fibres of the root ri-
sing from the surface, growing so as to form
a root. — Bolton. 116.

Woods, September, Rare.

Puff-ball. LYCOPERDON. Gen. Pl. 1603.

Mushroom roundish, emitting powdery seeds.

1156. *Star Puff-ball.* — *L. stellatum.*Wrapper manifold, expanding, head smooth,
mouth with pointed plaits. — Ray's Synop-
sis. I. i. E. Fungi. 312.

Pastures, October, Rare.

Ford-End.

Obs:

Head never sitting, star with from 5 to 7 segments.

Hygrometer

XXIV. Mushrooms. Crypt. Fungi.

1157. *Hygrometer Puff-ball.* — L. *recolligens.*

Wrapper with many spear-shaped divisions, head sitting, between round and compressed, mouth tapering upwards. — (Woodward.) E. Fungi. 80.

Fir groves, August, Rare.

Amptbill.

1158. *Projectile Puff-ball.* — L. *carpobolus.*

Wrapper manifold, fruit globular, composed of united seeds. — (See our fig. No. 6.)

Rotten pales, October, extremely rare.

Amptbill.

First found together with the two preceding species by Mrs. ABBOT.

Obs:

Plant growing in clusters: an ovate vesicle starts up from the disc, scattering the seed with its elastic motion.

1159. *Long-stalked Puff-ball.* — L. *pedunculatum.*

Stem long, head round, smooth, cylindrical, very entire. — Batsch. 167.

Walls, October, Common.

1160. *Variable Puff-ball.* — L. *Proteus.*

Roundish, top-shaped, or nearly on a stem, flesh white with soot-colored seeds, skin thin, flaccid. — Bulliard. 447.

Pastures, August, Common.

Pitted



1793 Published by James Sowerby London.

1159. Obs:

Mouth extremely small, stem cylindrical,
generally upright, sometimes leaning.

XXIV. Mushrooms. Crypt. Fungi.

1161. Obs.
Head yellowish, pitted as with the marks
of the small-pox.

1162. Obs.
Skin of this mushroom tender, brittle,
rough and branny.

1163. Obs.
Plant growing in clusters, either scarlet,
or brown with a cast of blue.

1165. Horseshoe Puffball. *L. equinum.*
Plant brown: white. Stem solid cylindrical
head globular, rather hollowed underneath
withering
E Fungi. 292.

Hors & Horseshoos. Winter - Rare.

1165. Obs. Evertor. Clapham.
Top-shaped, sitting, or nearly on a
stem.

1161. *Pitted Puff-ball.* — *L. verrucosum.*

Roundish, reddish-brown, stem bluish-red,
seeds brown, neck of the root with very
broad plaits. — Bulliard. 24.

Heaths, August, Rare.

*Ampthill.*1162. *Ash-colored Puff-ball.* — *L. cinereum.*

Blue-grey, globular, rough and branny, seeds
like sand, large, black, intermixed with
small white zigzag fibres. — Batsch. 169.

Rotten leaves and wood, August, Rare.

*Clapham between Woods.*1163. *Purple Puff-ball.* — *L. epidendrum.*

Skin and meal purple. — E. Fungi. 52.
Rotten wood, October, Common.

1164. *Slate-colored Puff-ball.* — *L. ardosiacum.*
Roundish, between azure and lead-colored,
flesh red, seeds brown, skin flexible. —
Bulliard. 192.

Pastures, January, Rare.

*Ford-End.*1165. *Cotton Puff-ball.* — *L. gossypinum.*

Head pear-shaped, white, cottony, tapering
downwards, seeds brown. — Bulliard. 435. 1.

Horses' hoofs, Winter, Rare.

Potton Wood.

X x

Brittle

XXIV. Mushrooms. Crypt. Fungi.

1166. *Brittle Puff-ball.* — L. fragile.

Parasitical, nearly sitting, inversely-ovate, skin shining, brittle, meal black, intermixed with soft hairs. — E. Fungi. 136.

On mosses, October, Rare.

Silsoe.

1167. *Conjurer's Fern.* — L. innatum.

White, spherical, solitary, parasitical, sitting, containing a white powder and opening at the top with many ragged clefts. — (Withering.) E. Fungi. 53. *Æcidium fuscum.*

RETICULARIA. Bulliard.

Mushroom in it's first state soft, containing seeds entangled amidst numerous fibres.

1168. *Custard Reticularia.* — R. septica.

Slimy, yellowish, scattered, in various shapes. — (Sibthorpe.) Schæffer. 194.

Woods, September, Common.

1169. *Egg-shaped Reticularia.* — R. ovata.

Stemless, egg-shaped, mucilaginous, yellowish, gills cellular, vanishing, turning to dust, blackish: seeds black, adhering to threads. — (Schæffer.) Bulliard. 380. 1.

Grass, mosses, &c. October, Rare.

Warden.

Smut

1167.

obs:

The figure of the yet-unexpanded plant to and the reflected hairy wrapper after its expansion seem to claim this for a *Lycoperdon* —

1168.

obs:

Plant in its first state like ripe cream cheese.

Vermillion

*1174, Pink Sphaeria. *S. miniate.*

Roundish, vermillion-colored, scattered,
pith mealy.

On the bark of small sticks
in woods Worcester. Rare
Clapham between Woods.

Obs.
not the miniate of Bolton which is
the tremelloides n^o 1173 of this book.

1171.

Obs.

Shape of the plant various, some-
times single, generally double, al-
ways regularly marked with con-
centric scores-

1172.

Obs.

Mushrooms small, generally single, between
rust and root-colored

1173.

Obs.

Top shaped, crowded, not unfrequently
incorporating -

*1173 - Bosom Sphaeria. *S. papillosa.*

Simple, roundish, black, tapering
to a point, with an indented circle
round it. *S. Fungi.* 236.

Sticks in Woods. Oct. Rare
Penrholt.

339

XXIV. Mushrooms. Crypt. Fungi.

1170. Smut Reticularia. — *R. segetum.*

Brownish-black, growing on other plants, full
of threads within. — Bulliard. 472. 2.

Ears of corn, June, too common.

SPHÆRIA. Haller.

Fructifications spherical, full of a blackish
powder, often secreted under the bark.

1171. Ash Sphaeria. — *S. fraxinea.*

Roundish, black, outwardly smooth, nearly
pustular, within marked with concentric
circles. — E. Fungi. 160. excellent.

Old stumps, all the year, Rare.

Clapham Lane.

1172. Black Sphaeria. — *S. lycoperdonoides.*

Compound, convex, nearly solid, pith mealy,
black. — Weigel Obs: III. 2 a.

Fallen boughs, Autumn, Common.

1173. Purple Sphaeria. — *S. tremelloides.*

Roundish, sitting, single, smooth, scarlet. —
Bulliard. 284.

Currant trees, Autumn, Common.

1174. Gregarious Sphaeria. — *S. gregaria.*

Simple, collected into irregular clusters, ver-
million, crust whitish, tender. — Weigel.

II. 10.

X x 2

Oaks,

Obs.

The upper surface of the
leaf, where this singular fungus
grows, is marked by a reddish
dentate hollow appearing scalded.

Scidium - Person's Fungi
Scidium smooth on both sides, shell of seeds uncovered and
growing together - Scidium lacatum - Ragged Scidium
Scidium lacatum from a brownish base, tubes lacinated
growing 2. Fungi 318.
Scidium Beddoe.
Scidium Beddoe.

XXIV. Mushrooms. Crypt. Fungi.

Oaks, February, Common.

1175. Breast-shaped Sphaeria. — *S. mammosa*.

Single, breast-shaped, black.

Woods on hazel stumps, Spring, Common.

1176. Scarlet Sphaeria. — *S. Mori*.Simple, clustered, scarlet, very small. —
Bolton. 120. 1. E. Fungi. 255.

Rotten wood, October, Rare.

Renhold Wood.

1177. Blood-colored Sphaeria. — *S. sanguinea*.

Simple, oval, blood-colored, open at the end.

Bulliard. 487. 3. E. Fungi. 254.

Rotten sticks, Autumn, Rare.

Clapham Park Wood.

1178. Depressed Sphaeria. — *S. depressa*.

Stemless, clustered, black, shining. — Bolton. 122. 1.

Rotten sticks, Autumn, Common.

1179. Stick Sphaeria. — *S. corticalis*.

Black, seeds very numerous, shining, round. —

Bulliard. 492. 2.

Sticks and walls, October, Common.

1180. Clustered Sphaeria. — *S. aggregata*.

Parasitical, round, mouth entire. — Bolton.

122. 2.

* 1181. Concave Sphaeria. — *S. concava*. RottenHollow, black, penetrating the upper coat
of the leaf only. — Bulliard. 317.

Holly leaves. Autumn and winter, rare.

Everton White wood, Potton Wood.

1174. Obs.
The Sphaerules are shaped like asterisks.1175. Obs.
altho' growing many together, they never
incorporate.1176. Obs.
Narrowest at the base, orange colored
when young.* 1176. Brush-shaped Sphaeria. — *S. tentaculata*
Black, hairlike, body of the fungus sunk in
the wood. — Bulliard. 120.
on dead stems of Chelidonium Glaucum in my
garden.Obs.
The cylindrical mouths of this fungillus are
as it were very well exposed to the naked
eye by so many obs. fine black hairs.
Almost transparent, after the dispersal
of its seeds, cup-shaped.* 1178. Bark-raising Sphaeria. — *S. decorticans*.
Stemless, shining, black, button-shaped,
Sphaerules very minute.

Bulliard 468. 4. E. Fungi. 137.

Decayed sticks, Autumn & Spring - Common.

Obs.
When the Sphaeria grow thick, they cause
the outer rind of the bark under which they
grow, to peel off, whence the name.

1180.

Obs:

Plant crusting over the fallen branches of trees with an inner and second bark, sphaerules very minute and scarcely visible without a magnifier.

1181.

Obs.

Very different from the Lichen pulicaris growing upon the soil at the very bottom of the aged stumps of trees, green when young, brown when old, never quite black.

* 1182 Minute Sphaeria. — *S. minuta*.
Incrusting, white, with black dots.

On the Bark of trees. Winter, rare
Renhold wood.

1184.

Obs:

Not very unlike the *S. mammosa*, but is less crowded, and appears hollow when dissected.

1185.

Obs:

When that part of the mushroom which represents the upper shell of the muscle has disappeared, and it is more advanced it acquires a blackish hue.

* 1181. Two-faced Sphaeria. — *S. bifrons*.

Almost flat, black, with the same external appearance on either side of the leaf.

E. Fungi. tab. 316.

Holly leaves, Autumn and Spring. ³⁴¹ rare

XXIV. Mushrooms. Crypt. Fungi. *Serton white*
wood

Rotten sticks, December, Common.

1181. Furrowed Sphaeria. — *S. sulcata*.

Stemless, with a longitudinal groove. — Bolton.

124? E. Fungi. 315.

Stumps, Autumn, Common.

1182. Incrufted Sphaeria. — *S. stigma*. S.D. sent

Compound, bark-like, black, dotted. — Hoff-
man. Veg. Crypt. V. 2.

Under the bark of trees, October, Common.

1183. Pointed Sphaeria. — *S. acuta*.

Single, very small, pointed, black. — E.
Fungi. 119. excellent.

Nettle stumps, January, Common.

1184. Round Sphaeria. — *S. globularis*.

One-celled, scattered, spherical, from white becoming black, smooth, breast-shaped at the point. — Bulliard. 444. 2.

Stumps in woods, October, Common.

1185. Muscle Sphaeria. — *S. ostreacea*.

One-celled, scattered, from ash-colored turning brownish-black, nearly flattened, smooth, with two valves. — Bulliard. 444. 4.

Fallen boughs, October, Common.

1186. Thorn-like Sphaeria. — *S. echinata*. S.D. sent

Many-celled,

1186. Common Sphaeria. *S. communis.*
Almost cup shaped, black, fragi:
form, open at the top.

9. Fungi. 295.

342 Limestone Walls, Common.

XXIV. Mushrooms. Crypt. Fungi.

Many-celled, black, stud-shaped, flat, surface
smooth, disc covered with prickles. — Bulli-
ard. 432. 2.

Between the inner and the outer rind of trees,
Winter, Rare.

Clapham between Woods.

1187. Club-shaped Sphaeria. — *S. clavata.*

One-celled, between thyrse-shaped and crou-
ded, scattered, lengthened out, tapering at
the base, from whitish turning black,
smooth. — Bulliard. 444. 5.

Branches of trees, Autumn, Rare.

Clapham between Woods.

1188. Black Sphaeria. — *S. atra. S.D.*

Many-celled, brownish-black, with pointed
studs, seeds elliptical, breast-shaped at the
point. — Bulliard. 432. 1.

Branches of trees, October, Common.

1189. Tendril Sphaeria. — *S. cirrhata.*

One-celled, scattered, white turning black,
receptacles bearing tendrils. — E. Fungi.
138. excellent.

Branches of trees, October, Rare.

Renhold Wood.

1190. Covered Sphaeria. — *S. obducta.*

Stemless,

1186. Obs:
Sphaerules in clusters shaped like thorns
and surmounted with entirely black
dots.

1187. Obs:
Sphaerias in their young state almost
pearshaped, whitish at the point, when
older club-shaped and very thickly
set with numberless Sphaerules.

*1187 Black Sphaeria. *S. fruticulosa.*
~~Cup-shaped, black, seeds numerous, rising~~
~~lobose.~~ Clapham. Stepp. to Colera.
Bulliard t. FIG. J. 2 Canta.
Bark of trees. Winter - London p. 34.

Obs:

1188. Sphaerias simple, Sphaerules pointed
many together & united, without
brownish-black, within whitish.

1189.

Obs:
Sphaerias button-shaped, out of which
after a little white capillary sub:
stances arise, twisting up in a
spiral manner.

1190. Obs.

Plant clustered, in various shapes.
Sometimes round, or elliptical but
often kidney shaped.

1190. Under *Bartsia*.

Black, sitting, distinct, within the
exterior bark. — *S. Fungi*. 296.
Trees, October, Common.

1191. Obs:

Plant rust-colored, stems fine as a hair.

1193. Obs:

Stems increasing in breadth at the base,
a longitudinal axis passing thro' all the
capsules.

343

XXIV. Mushrooms. Crypt. Fungi.

Stemless, clustered, rather woolly. — Bulliard. 492. 1.

Sticks, Autumn, Common.

TRICHLIA. Haller.

Mushroom growing in clusters, capsules cylindrical, between top-shaped and oblong or round, producing seeds in net-work filaments.

1191. Naked Trichia. — *T. nuda*.

On a stem, head oblong, growing on a longitudinal axis. — Bulliard. 477. 1.

Rotten wood, October, Common.

1192. Purple Trichia. — *T. denudata*.

Stem very short, capsule lengthened out, oval, not perforated, cup-shaped at the base. — *E. Fungi*. 49.

Rotten wood, October, Rare.

Southill.

1193. Cat's-tail Trichia. — *T. typhaeformis*.

Stem broader at the base, capsule always cylindrical. — *E. Fungi*. 50.

Rotten Wood, September, Rare.

Silsoe, — Southill.

1194. Ash-colored Trichia. — *T. cinerea*.

Stem short, tapering, capsules greyish-white. — Bulliard. 477. 3.

Rotten

XXIV. Mushrooms. Crypt. Fungi.

In Rotten boughs, Autumn, Common.

1195. Least Trichia. — *T. minuta*.

On a stem, bell-shaped, bearing small lentil-shaped seeds.—Hoffman. Veg. Crypt. II. 2.

On *Hypnum triquetrum*, Winter, Rare.

Clapham Park Wood.

Obs:

Stem purplish, bearing a cup resembling a wine-glass in miniature, springing from living mosses.

1196. Olive-colored Trichia. — *T. olivacea*.

On a stem, stem and head woolly, olive-colored. — Bolton. 94. 2.

Rotten wood, Autumn, Rare.

Renhold Wood.

1197. Branny Trichia. — *T. furfuracea*.

On a stem, stem greenish, thread-shaped, head round, mealy. — Batsch. 178.

Rotten wood, December, Rare.

Renhold Wood.

1198. Grey Trichia. — *T. globulifera*.

On a stem, head round, ash-colored, grey. — Bulliard. 484. 3.

Clefts of wood, Autumn, Rare.

Clapham Park Wood.

Liverwort

*. Round-headed Trichia. Sphaerocephala.
on a stem, head globular, between
cinerous and black

P. Fungi. 240.

Straws and dead sticks Winter, Common.

Obs:

Stem very short, head when in perfection black, whilst under the process of emitting the fructifying powder, ash-colored —

1197. *Obs:*

Seagreen, leaning, heads brown, branny.

1198. *Obs:*

stem thickening downwards, capsule whitish, at first inclosing all the seeds, afterwards torn and fugacious.

XXIV. Mushrooms. Crypt. Fungi.

1199. Liverwort *Trichia*. — *T. lichenoides*.

Stem cylindrical, black, head round, of the same color. — (*Sibthorpe.*) Sowerby. 414.
Lichen sphærocephalus.

Park pales, November, Rare.

Ampthill Park.

1200. Top-shaped *Trichia*. — *T. turbinata*.

Stemless, top-shaped, yellow, wool yellowish. — E. Fungi. 85.

Rotten wood, August, Common.

1201. Pear-shaped *Trichia*. — *T. pyriformis*.

Yellow, varnished, capsules top-shaped, gradually ending in a cylindrical stem. — (*Sibthorpe.*) Bulliard. 417. 2.

Rotten wood, August, Common.

1202. Black *Trichia*. — *T. recutita*.

On a stem, shaped like a fig, roundish, wool black. — Bulliard. 417. 3.

Rotten wood, September, Common.

1203. Blackish *Trichia*. — *T. fusco-ater*.

Stemless, brown-black, seeds yellow. — Bulliard. 417. 5.

Limestone walls, Autumn, Rare.

Friar's Walls.

Y y

Mould.

1200. Obs.

Heads yellow, with golden-colored seeds.

1201. Obs.

Shape of the plant much like that of the preceding, but the heads rather pear-shaped, and less crowded.

1202. Obs:

After the dispersion of the woolly substance, the skin remains and bears a clefted appearance.

XXIV. Mushrooms. Crypt. Fungi.

Mould. MUCOR. Gen. Pl. 1685.

Mushroom with vesicles standing on stems,
in which numerous seeds are contained.

1204. *Cotton Mould.* — M. Mucedo.

On a stem, head round. — Bulliard. 480. 2.
Rotten planks, October, Common.

1205. *Greenish Mould.* — M. glaucus.

On a stem, head roundish, in clusters. —
Flora Danica. 777. 2.
On rotten fruit, Common.

1206. *Dewy Mould.* — M. roridus.

On a stem, fugacious, stem hair-like, head
spherical, like a dewdrop, with a black
dot at the point. — (Relhan.) Bulliard.
480. 1.
Cow and horse-dung, August, Rare.

Paradise, near Bedford.

1207. *Pitcher-shaped Mould.* — M. urceolatus.

On a stem, fugacious, stem bellying upwards,
transparent, like a dew-drop, head roundish,
elastic, black. — Dickson. I. 111. 6.

With the former, but rarer.

1208. *Black Mould.* — M. embolus.

On a black bristle-shaped stem, hair woolly,
brown.
Rotten wood, Winter, Common.

Fingered

E. Fungi. 300.

1204.

Obs:

Stem very long, head round, first white,
afterwards black.

1209.

Obs:

In some specimens it bears a distant resemblance to the spikes of *Panicum sanguinale*.

1210.

Obs:

Heads radiating, plant nearly scarlet.

1211.

Obs:

Plant seen with the help of a magnifier, is like a small shrub.

1212.

Obs:

Seeds sometimes simple, as often in bunches.

347

XXIV. Mushrooms. Crypt. Fungi.

1209. Fingered Mould. — *M. caespitosus*.

On a branched stem, fructifications fingered. — Bulliard. 504. 11.

Rotten vegetables in woods, Winter, Rare.

Thurlegh Wood.

1210. Radiating Mould. *M. crustaceus*.

On a simple stem, fructifications radiating, terminal. — Bulliard. 504. 10.

Under mosses in woods, October, Rare.

Potton Wood.

1211. Grape Mould. — *M. botrytis*.

On a stem, fructifications in bunches. — Bulliard. 504. 7.

On the caps of old Boleti. October, Rare.

Clapham Lane.

1212. Golden Mould. — *M. chrysospermus*.

Stems erect, with very numerous roundish seeds, at first white, afterwards gold-colored. — Bulliard. 504. 1.

On the caps of old Boleti, October, Common.

1213. Stick Mould. — *M. lignifractus*.

Pimple-shaped, at first white, afterwards green, stems very fine, branched, intermixed, seeds very minute, roundish, scattered. — Bulliard. 504. 6.

Y y 2

Rotten

XXIV. Mushrooms. Crypt. Fungi.

Rotten boughs, Winter, Common.

1214. *Granulous Mould.* — *M. granulosus.*

Incrustated, blackish, seed-covers sitting, adhering as well to the upper as under surface of the leaves. — Bulliard. 504. 5.

Rotten leaves, Winter, Common.

1215. *Sessile Mould.* — *M. Erysyphe.*

White, heads brown, sitting. — Bulliard. 504. 12.

Decayed leaves, September, Common.

1216. *Cheese Mould.* — *M. caseus.*

Crust rather leather-like; stems very short, heads or seeds roundish, white, yellow or scarlet. — E. Fungi. 180.

Cheese in damp Cellars, Common.

1217. *Orange Mould.* — *M. aurantius.*

Crustaceous, stems branched, creeping, seeds few, roundish, scattered, very minute, orange-colored. — Bulliard. 504. 5.

Inside of willows, Winter, Rare.

Goldington.

Plants added.

Threads in I Set. X Males. Monad. Decand.

1218. *Bloody Crane's-bill.* — *G. sanguineum.*

Fruitstalks with 1 flower, leaves with 5 divisions, subdivisions trifid, outline circular. — Sowerby. 272.

Hedge bottoms, May, P. Rare.

Potton. Mr. RUGELY.

Threads in II Sets. X Males. Diad. Decand.

1219. *White Upright Trefoil.* — *T. hybridum.*

Heads somewhat umbel-like, shells with 4 seeds in each, stem ascending. — Vaillant. XXII. 5.

Moist pastures, May, P. Rare.

Caldwell pastures.

Threads on the Female. Gynand. Diand.

1220. *Spider Ophrys.* — *O. aranifera.*

Bulbs roundish, stem leafy, lip of the nectary roundish, entire, nicked, convex. — Sowerby. 65.

Old sand and gravel pits, May, P. Rare.

Southill. Mr. P. WALKER.

Juniper

*1220. ~~*Map Ophrys*~~ ~~*O. antrophocephala.*~~
 Bulbs roundish, stem leafy, lip ~~trap-shaped~~
~~in 3 divisions, the middlemost elongated~~
~~and cleft - English Botany.~~
~~Sowerby. 27.~~
~~pastures. June. P. Rare~~
~~Kentish, on the borders of Hants~~
~~Recd. T. Martyn. P. B.~~
~~ips~~
~~helmet brownish —~~

II Houses. Threads in many Sets. Dioec. Monad.

Juniper. JUNIPERUS. Gen. Pl. 1552.

Male. — Cup of the catkin scaly. Blossom none.

Female. — Cup of 3 divisions. Petals 3.
Styles 3. Berry 4-seeded. Tuberles of
the unequal cup three.

1221. *Juniper Tree.* — J. communis.

Leaves in threes, expanding, pointed, longer
than the berry. — Blackwell. 187.

Hilly spots, May, S. Rare.

Warden.

XXIV. Mushrooms. Crypt. Fungi.

* Agarics, solid and decurrent.

1222. *Golden Agaric.* — A. aureus.

Gills orange with a leaden cast, 4 in a set,
cap deep-orange or gold-colored, stem
lighter, ring permanent. — E. Fungi. 77.

Woods, Autumn, Rare.

Clapham Park Wood.

* * Solid and fixed.

1223. *Pinky Agaric.* — A. puniceus.

Gills white, 4 in a set, cap pinky, convex,
stem white. — (Withering.)

Woods, October, Common.

Purplish

XXIV. Mushrooms. Crypt. Fungi.

1224. *Purplish Agaric.* — *A. subpurpurascens.*

Gills reddish-brown, cap brown, purplish at the edge, stem violet-colored, scurfy, bulbous at the base. — Batsch. 74.

Pastures, October, Common.

1225. *Uncurtained Agaric.* — *A. nudus.*

Gills cool-brown, numerous, 4 or 8 in a set, cap brown, gently convex, concave with age, satiny when dry, stem brown. — (Withering.) Bulliard. 439. (the lower figures.)

Woods, October, Rare.

Warden,—Clapham Park.



13. - 13. In the Title for III read IV Males.

HYPOGRAPHICAL AND OTHER ERRORS.

- Page. Line. 13. 13.
 13. - 13. For MILUIM read MILIUM.
 32. - — In the Title for III read IV Males.
 32. - 4. For 101. read 110.
 61. - 24. For 109. read 209.
 136. - 14. For 464. read 454.
 199. - 8. Add — Watery pits, June, P. Rare.
Ford End.
 274. - 3. For Weigers read Weigel.
 319. - — Before No. 1092. Hydnum barba Jovis
 insert this mark,
 †‡ Stemless.
 320. - 12. After E. Fungi, for II. read 11.

GENERIC INDEX.

A	Pag.	A	Pag.	A	Pag.
A C E R	219	A r e m i s i a	179	C a m p a n u l a	48
Achillea	186	Arum	197	Cardamine	142
Acorus	77	Arundo	25	Carduus	174
Adoxa	91	Asperula	32	Carex	200
Ægopodium	69	Asplenium	225	Carlina	177
Æthusa	65	Astragalus	160	Carum	67
Agaricus	279	Atriplex	219	Caucalis	53
Agrimonia	104	Atropa	50	Centaurea	187
Agrostemma	100	Auricularia	321	Centunculus	39
Agrostis	13	Avena	24	Cerasium	101
Aira	15	B		Chaerophyllum	66
Ajuga	125	Ballota	130	Chara	197
Alchemilla	35	Bellis	184	Cheiranthus	144
Alisma	83	Berberis	80	Chelidonium	115
Allium	74	Betonica	130	Chenopodium	54
Alopecurus	12	Betula	207	Chironia	47
Alsine	70	Bidens	177	Chlora	85
Anagallis	46	Boletus	311	Chrysanthemum	184
Anemone	118	Borago	43	Chrysosplenium	92
Anethum	67	Brassica	145	Cichorium	173
Angelica	61	Briza	19	Cicuta	65
Anthemis	186	Bromus	22	Circaeа	7
Athericum	77	Bryonia	216	Cistus	117
Anthoxanthum	7	Bryum	236	Clavaria	332
Anthyllis	155	Bunium	59	Clematis	119
Antirrhinum	136	Bupleurum	57	Clinopodium	132
Apium	69	Butomus	91	Cochlearia	141
Aquilegia	118	Bysssus	276	Colchicum	83
Arabis	145	C		Comarum	115
Arctium	174	Callitriches	1	Conferva	274
Arenaria	97	Caltha	124	Conium	60
				Convallaria	

G E N E R I C I N D E X.

Convallaria	-	76	Fritillaria	-	75	Jungermania	-	252
Convolvulus	-	47	Fumaria	-	152	Juniperus	-	350
Conyzza	-	180	G			L		
Cornus	-	35	Galanthus	-	73	Lathraea	-	135
Corylus	-	211	Galeobdolon	-	129	Lamium	-	128
Crataegus	-	108	Galeopsis	-	129	Lapsana	-	173
Crepis	-	171	Galium	-	33	Lathyrus	-	156
Cucubalus	-	95	Genista	-	153	Lemna	-	198
Cuscuta	-	36	Gentiana	-	56	Leontodon	-	169
Cynoglossum	-	41	Geranium	-	148	Leonurus	-	131
Cynosurus	-	27	Geum	-	114	Lichen	-	256
D			Glechoma	-	128	Ligustrum	-	2
Dactylis	-	20	Gnaphalium	-	179	Limosella	-	139
Daphne	-	87	H			Linum	-	71
Daucus	-	59	Hedera	-	53	Lithospermum	-	41
Delphinium	-	118	Hedysarum	-	160	Lolium	-	26
Dianthus	-	94	Helleborus	-	124	Lonicera	-	49
Digitalis	-	139	Helvelia	-	320	Lotus	-	164
Dipsacus	-	28	Heracleum	-	61	Lychnis	-	100
Draba	-	140	Hieracium	-	170	Lycoperdon	-	335
Drosera	-	71	Hippocrepis	-	159	Lycopodium	-	228
E			Hippuris	-	1	Lycopsis	-	43
Echium	-	43	Holeus	-	217	Lycopus	-	6
Epilobium	-	84	Hordeum	-	26	Lysimachia	-	45
Equisetum	-	221	Hottonia	-	45	Lythrum	-	103
Erica	-	87	Humulus	-	214	M		
Erigeron	-	180	Hyacinthus	-	74	Malva	-	151
Eriophorum	-	11	Hydnnum	-	318	Marchantia	-	255
Erodium	-	147	Hydrocharis	-	216	Marrubium	-	131
Eryngium	-	158	Hydrocotyle	-	56	Matricaria	-	185
Erysimum	-	144	Hyoscyamus	-	50	Medicago	-	164
Evonymus	-	52	Hyoseris	-	172	Melampyrum	-	135
Eupatorium	-	178	Hypericum	-	165	Melica	-	16
Euphorbia	-	105	Hypnum	-	244	Melissa	-	133
Euphorasia	-	134	Hypochaeris	-	172	Mentha	-	127
F			I	J		Menyanthes	-	44
Fagus	-	210	Jasione	-	189	Mercurialis	-	215
Festuca	-	20	Iberis	-	141	Merulius	-	278
Filago	-	188	Ilex	-	37	Milium	-	13
Fontinalis	-	230	Inula	-	183	Mnium	-	232
Fragaria	-	112	Iris	-	9	Montia	-	28
Fraxinus	-	220	Juncus	-	78	Monotropa	-	92

Mucor

G E N E R I C I N D E X.

Mucor	-	346	Poa	-	17	Scandix	-	63
Myosotis	-	40	Polygala	-	152	Schoenus	-	9
Myosurus	-	72	Polygonum	-	88	Scirpus	-	10
Myriophyllum	208		Polypodium	-	226	Scleranthus	-	93
N			Polytrichum	-	232	Scrophularia	-	138
Narcissus	-	73	Populus	-	215	Scutellaria	-	134
Nardus	-	11	Potamogeton	-	37	Sedum	-	98
Nepeta	-	126	Potentilla	-	112	Sempervivum	-	106
Nymphaea	-	117	Poterium	-	209	Senecio	-	181
O			Primula	-	44	Serapias	-	196
Oenanthe	-	64	Prunella	-	133	Serratula	-	174
Ononis	-	154	Prunus	-	107	Sherardia	-	32
Onopordum	-	176	Pteris	-	224	Silene	-	95
Ophioglossum	223		Pulmonaria	-	41	Sinapis	-	146
Ophrys	-	194	Pyrus	-	109	Sison	-	63
Orchis	-	191	Pyrola	-	92	Sisymbrium	-	142
Origanum	-	132	Q			Sium	-	62
Ornithogalum	75		Quercus	-	210	Smyrnium	-	67
Ornithopus	-	159	R			Solanum	-	51
Orobanche	-	140	Ranunculus	-	120	Solidago	-	183
Orobus	-	155	Raphanus	-	147	Sonchus	-	169
Osmunda	-	223	Reseda	-	104	Sorbus	-	109
Oxalis	-	100	Reticularia	-	338	Sparganium	-	200
P			Rhamnus	-	52	Spartium	-	153
Papaver	-	115	Rhinanthus	-	134	Spergula	-	102
Parietaria	-	218	Ribes	-	52	Sphaeria	-	339
Paris	-	90	Rosa	-	110	Sphagnum	-	228
Parnassia	-	70	Rubus	-	111	Spiraea	-	110
Pastinaca	-	67	Rumex	-	81	Splachnum	-	281
Pedicularis	-	136	S			Stachys	-	130
Peplis	-	77	Sagina	-	39	Stellaria	-	96
Peucedanum	-	66	Sagittaria	-	209	Symphytum	-	42
Peziza	-	324	Salix	-	211	T		
Phalaris	-	13	Salvia	-	6	Tamus	-	214
Phallus	-	323	Sambucus	-	70	Tanacetum	-	178
Phascum	-	229	Samolus	-	49	Teucrium	-	125
Phellandrium	65		Sanguisorba	-	31	Thalictrum	-	120
Phleum	-	12	Sanicula	-	57	Thlaspi	-	141
Picris	-	168	Saponaria	-	94	Thymus	-	132
Pimpinella	-	68	Satyrium	-	193	Tilia	-	116
Pinguicula	-	5	Saxifraga	-	93	Tragopogon	-	168
Plantago	-	30	Scabiosa	-	29	Tremella	-	270
						Trichia		

GENERIC INDEX.

Trichia	- 343	Ulmus	- 55	Veronica	- 2
Trifolium	- 161	Ulya	- 273	Viburnum	- 69
Triglochin	- 83	Urtica	- 208	Vicia	- 157
Triticum	- 27	Utricularia	- 5	Vinca	- 53
Tuber	- 335	Vaccinium	- 86	Viola	- 189
Tulipa	- 74	Valantia	- 218	Viscum	- 213
Tussilago	- 181	Valeriana	- 8		
Typha	- 199	Verbascum	- 50		
U		Verbena	- 126		
Ulex	- 154				



ENGLISH INDEX.

A	Pag.		Pag.		
ADDER's-		Bird's-nest	- 92	Canary-grass	- 13
tongue	- 223	Bladderwort	- 5	Candy-tuft	- 141
		Blinks	- 28	Carraways	- 67
Agaric	- 279	Bloodwort	- 31	Carrot	- 59
Agrimony	- 104	Bog-moss	- 228	Carline Thistle	- 177
— Dutch	178	Bog-rush	- 9	Catchfly	- 95
— Hemp	177	Boletus	- 311	Cat-mint	- 126
Alisanders	- 67	Borage	- 43	Cat's-ear	- 172
All-heal	- 129	Bottle-moss	- 231	Cat's-foot	- 188
Anemone	- 118	Brakes	- 224	Cat's-tail-grass	- 12
Angelica	- 61	Bramble	- 111	Celandine	- 115
Archangel	- 128	Brome-grass	- 22	Centaury	- 187
Arrow-grass	- 83	Broom	- 153	Chamomile	- 186
Arrow-head	- 209	Broom-rape	- 140	Chaff-weed	- 30
Ash	- 220	Brookweed	- 49	Chervil	- 66
Auricularia	- 321	Bryony	- 216	Chickweed	- 70
		Buckthorn	- 52	Chironia	- 47
B		Buckbean	- 44	Cicely	- 65
Barberry	- 89	Bugle	- 125	Cinquefoil	- 112
Barley	- 26	Bugloss	- 43	Claver	- 164
Basil	- 132	Bur-reed	- 200	Clot-burr	- 174
Bedstraw	- 33	Burnet	- 209	Club-moss	- 228
Beech	- 210	Burnet Saxifrage	68	Club-rush	- 10
Bellflower	- 48	Butterwort	- 5	Club-top	- 332
Bennet	- 114	Byssus	- 276	Cockle	- 100
Bent-grass	- 13			Cock's-head	- 160
Betony	- 130	C		Cabbage	- 145
Bindweed	- 47	Betony	- 133	Cock's-foot grass	- 20
Birch	- 207	Calamint	- 133	Colt's-foot	- 181
Bird's-foot	- 159	Campion	- 95	Columbine	- 118

ENGLISH INDEX.

Comfrey	-	42	Fox-glove	-	139	Horse-bane	-	65
Convally	-	76	Fox-tail-grass	-	12	Horse-shoe Vetch	-	159
Cornel	-	35	Fritillary	-	75	Horse-tail	-	221
Cotton-grass	-	11	Frog-bit	-	216	Hottonia	-	45
Cotton-thistle	-	176	Fumitory	-	152	Hound's-tongue	-	41
Cow-bane	-	65	Furze	-	154	House-leek	-	106
Cow-wheat	-	135	G			Hyacinth	-	74
Crane's-bill	-	148	Garlick	-	74	Hydnum	-	318
Cross-wort	-	218	Gentian	-	56	I J		
Crowfoot	-	120	Germander	-	125	Juniper	-	359
Cuckow-pint	-	197	Goat's-beard	-	168	Ivy	-	53
Cudweed	-	179	Golden-rod	-	183	K		
Currant	-	52	Golden-Saxifrage	-	92	Knawel	-	93
D			Goose-foot	-	54	Knot-grass	-	88
Daffodil	-	73	Gout-weed	-	69	L		
Daisy	-	184	Grass of Parnassus	-	79	Ladies'-finger	-	155
Dandelion	-	169	Grass-poly	-	103	Ladies'-mantle	-	35
Darnel-grass	-	26	Greenwood	-	153	Ladies'-seal	-	214
Dill	-	67	Gromil	-	41	Ladies'-smock	-	142
Dock	-	81	Ground-ivy	-	128	Larkspur	-	118
Dodder	-	36	Groundsel	-	181	Laver	-	273
Dog's-tail grass	-	27	H			Lime-tree	-	116
Drop-wort	-	64	Hair-grass	-	15	Livergreen	-	255
Duck-weed	-	198	Hair moss	-	232	Liverwort	-	256
Dyer's-weed	-	104	Hawk's-beard	-	171	Loosestrife	-	45
E			Hawk-weed	-	170	Lousewort	-	136
Earth-moss	-	229	Hawthorn	-	108	Lungwort	-	41
Earth-nut	-	59	Hazel	-	211	Lychnis	-	100
Elder	-	79	M					
Elecampane	-	183	Hellebore	-	124	Mallow	-	151
Elm	-	55	Helleborine	-	196	Majoram	-	132
Eye-bright	-	184	Hemlock	-	60	Marigold	-	184
F			Hen-bane	-	50	Maple	-	219
Feather-moss	-	244	Hen-bit	-	130	Mare's-tail	-	1
Fescue-grass	-	20	Hen's-foot	-	58	Marshlocks	-	115
Fever-few	-	185	Herb-Paris	-	90	Marsh-marigold	-	124
Field-madder	-	32	Hog-weed	-	61	Marsh-moss	-	232
Fig-wort	-	138	Holly	-	37	Mat-grass	-	11
Flag	-	9	Hone-wort	-	63	Meadow-grass	-	17
Flax	-	71	Honey-suckle	-	49	Meadow-rue	-	120
Flea-bane	-	180	Hop	-	214	Meadow-saffron	-	83
Flowering-rush	-	91	Horehound	-	131	Meadow-sweet	-	110

Mealy-tree

Mealy-tree	-	69	Peziza	-	324	Shepherd's-purse	-	141
Medick	-	164	Pimpernel	-	46	Skull-cap	-	134
Melic-grass	-	16	Pink	-	94	Snap-dragon	-	136
Mercury	-	245	Plantain	-	30	Snow-drop	-	73
Merulius	-	278	Plumb	-	107	Soft-grass	-	217
Mezereon	-	87	Polypody	-	226	Sopewort	-	94
Milfoil	-	186	Pondweed	-	37	Southern-wood	-	179
Milkwort	-	152	Poplar	-	215	Sow-thistle	-	169
Millet-grass	-	13	Poppy	-	115	Speedwell	-	2
Mint	-	127	Primerose	-	44	Sphaeria	-	339
Mistletoe	-	213	Privet	-	2	Spider-wort	-	77
Moonwort	-	223	Puff-ball	-	335	Spikenard	-	180
Morcl	-	323	Purslane	-	77	Spindle-tree	-	52
Moschatel	-	91	Q			Spleenwort	-	225
Motherwort	-	131	Quaking-grass	-	19	Spurge	-	105
Mould	-	346	R			Spurry	-	102
Mouse-ear	-	101	Radish	-	147	St. John's-wort	-	165
Mouse-tail	-	72	Rattle	-	134	Star-jelly	-	270
Mudwort	-	139	Reed	-	77	Star of Bethlehem	-	75
Mullein	-	50	Reed-grass	-	25	Startrip	-	252
Mustard	-	146	Reed-mace	-	199	Starwort	-	1
N			Restarrow	-	154	Stitchwort	-	96
Nettle	-	208	Reticularia	-	338	Stonecrop	-	98
Night-shade	-	51	Riverweed	-	274	Stonewort	-	197
— Deadly	-	50	Rose	-	110	Stork's-bill	-	147
— Enchanter's	-	7	Rush	-	78	Strawberry	-	112
Nipplewort	-	173	S			Succory	-	173
O			Sage	-	6	Sulphurwort	-	60
Oak	-	210	Saintfoin	-	160	Sun-dew	-	71
Oat-grass	-	24	Sandwort	-	97	Sunflower	-	117
Orach	-	219	Sanicle	-	57	Swine's-succory	-	172
Orchis	-	191	Satyrion	-	193	T		
Ox-tongue	-	168	Saw-wort	-	174	Tansy	-	178
P			Saxifrage	-	93	Teasel	-	28
Parsley	-	69	Scabious	-	29	Thistle	-	174
Parsnip	-	67	Scorpion-grass	-	40	Thoroughwax	-	57
Pear	-	109	Scurvy-grass	-	141	Thread-moss	-	236
Pearlwort	-	39	Sedge	-	200	Thyme	-	132
Peasling	-	155	Self-heal	-	133	Tine-tare	-	158
Pellitory	-	218	Service	-	109	Tooth-wort	-	135
Pennywort	-	56	Sheep's-Scabious	-	189	Traveller's-joy	-	119
Periwinkle	-	53	Shepherd'sneedle	-	65	Trefoil	-	161

Trichia

ENGLISH INDEX.

Trichia	340	Viper's-bugloss	43	Wheat-grass	27
Truffle	336	Violet	189	Whitlow-grass	140
Tulip	74	W		Whorts	86
Turban-top	320	Wall-flower	144	Willow	211
Turkey-pod	145	Water-cress	142	Willow-herb	84
Tway-blade	194	Water-horehound	6	Winter-green	92
V		Water-lily	117	Wood-ruff	32
Valerian	8	Water-milfoil	208	Wood-sorrel	100
Vernal-grass	7	Water-moss	230	Wormseed	144
Vervain	186	Water-parsnip	62	Woundwort	130
Vetch	157	Water-plantain	83	Y	
Vetchling	156	Weazle-snout	129	Yellow-wort	85

In the Latin Generic Index after *Splachnum*, for 281 read 231.



INDEX AGARICORUM SPECIFICUS.

Ruginosus	—	298	crassipes	—	288
albus	—	293	orbaceus	—	288
alliaceus	—	303	crustuliniformis	—	295
amethystinus	—	282	cyathiformis	—	280.
androbaceus	—	284	cylindricus	—	304
applicatus	—	310.	dealbatus	—	288
aqenosus	—	304	drysophyllum	—	305
arancosus	—	287	burneus	—	280
aurantius	—	298	clixus	—	287
aureus	—	350	ericeus	—	282
auricomus	—	299	farinaceus	—	297
buccinalis	—	299	fascicularis	—	301
bulbosus	—	294	fibula	—	285
caseus	—	281	fimiputris	—	300
campestris	—	294	floccosus	—	300
carnosus	—	305	fragrans	—	281
ceraceus	—	300	geophyllum	—	289
cinnamomeus	—	300	glaucopterus	—	289
clavus	—	294	graveolens	—	293
clypeatus	—	299	harmologum	—	305
clypeolarius	—	304	hinnuleus	—	289
collinitus	—	294	hypni	—	301
columbarius	—	298	impuber	—	306
compreffus	—	299	infundibuliformis	—	280
conchatus	—	309	integer	—	290
confuentus	—	286	irregularis	—	297
contiguus	—	284	lachrymabundus	—	306
coriaceus	—	310.	lactifluus	—	282.
cossus	—	287.			

<i>lateralis</i>	309
<i>limacina</i>	284
<i>Listeri</i>	201
<i>Lividio-rubescens</i>	283
<i>luridus</i>	295
<i>Clycoperdonoides</i>	295
<i>Macer</i>	301
<i>marginatus</i>	306
<i>melleus</i>	301
<i>Micaceus</i>	285
<i>minutulus</i>	295
<i>Mollis</i>	310
<i>Momentaneus</i>	306
<i>Muscarius</i>	290
<i>niveus</i>	298
<i>nudus</i>	351
<i>odoros</i>	286
<i>orchades</i>	296
<i>orichalceus</i>	290
<i>ostreatus</i>	309
<i>ovatus</i>	306
<i>petalodes</i>	311
<i>peronatus</i>	291
<i>pilosus</i>	296
<i>pilulariformis</i>	307
<i>pipetatus</i>	283
<i>pistillaris</i>	280
<i>plicatilis</i>	307
<i>procerus</i>	307
<i>psittacinus</i>	302
<i>pulvinatus</i>	296
<i>puniceus</i>	350

<i>querinus</i>	311
<i>radicatus</i>	297
<i>radicosus</i>	296
<i>ramealis</i>	279
<i>rinosus</i>	291
<i>rufus</i>	291
<i>rutilus.</i>	205
<i>Semiglobatus</i>	302
<i>Semiovatus</i>	308
<i>sepsilis.</i>	310.
<i>sordidus</i>	292
<i>stipitis.</i>	291
<i>striatus</i>	302
<i>stypticus</i>	308
<i>Subpurpurascens</i>	351
<i>Sulphureus.</i>	284
<i>tener</i>	302
<i>terreus</i>	292
<i>titubans</i>	308
<i>ulmaria</i>	286
<i>umbelliferus</i>	303
<i>umbilicatus</i>	298
<i>umbraculum</i>	282
<i>varius</i>	303
<i>velatus</i>	283
<i>velutipes</i>	308
<i>verrucosus</i>	292
<i>violaceus</i>	293
<i>viridis</i>	292
<i>xerampelinus</i>	287
<i>Zylophilus.</i>	303

120-

LICHENES.

<i>albus</i>	256
<i>alcicornis</i>	269
<i>aphthosus</i>	267
<i>ater</i>	261
<i>atro: albus</i>	257
<i>Baeomyces</i>	260
<i>Byssoides</i>	260
<i>Calcareas</i>	258
<i>Calicaris</i>	264
<i>candelarius</i>	262
<i>canescens</i>	259
<i>caninus</i>	267
<i>Caperatus</i>	264
<i>carpinus</i>	260
<i>ceruleonigricans</i>	259
<i>ciliaris</i>	265
<i>cocciferus</i>	269
<i>corniculatus</i>	265
<i>cornucopiodes</i>	268
<i>cornutus</i>	269
<i>concolor</i>	265
<i>cristatus.</i>	263
<i>ericetorum</i>	260
<i>fagineus</i>	257
<i>farinaceus</i>	265
<i>filiformis</i>	268
<i>fimbriatus</i>	269
<i>flavus</i>	256
<i>flavo: rubescens</i>	261
<i>fraxineus</i>	266
<i>fuscoc: ater</i>	259
<i>horitus</i>	270
<i>horizontalis</i>	267
<i>imversus</i>	258
<i>incanus</i>	256
<i>jubatus</i>	270.
<i>muralis</i>	264
<i>Muscorum</i>	258
<i>nigrescens</i>	266
<i>olivaceus</i>	263
<i>pallens</i>	262
<i>parietinus</i>	263
<i>pertusus</i>	263
<i>phytodes</i>	266
<i>planastri</i>	257
<i>pubicaris</i>	267
<i>palmariacus</i>	268
<i>pyxidatus.</i>	268
<i>querneus.</i>	259
<i>rangiiferinus</i>	269
<i>rugosus.</i>	257
<i>sanguinarius</i>	258
<i>sakatilis.</i>	264
<i>scriptus</i>	257
<i>scruposus</i>	262
<i>stellaris</i>	264
<i>subfuscus</i>	262
<i>Sulphureus.</i>	259
<i>tenellus</i>	266
<i>Tuberocalatus.</i>	268
<i>vernalis</i>	261
<i>uncialis.</i>	270.

63-

MNIA.

- cospiticium — 235
capillare — 234
cuspitatum — 234
fontanum — 232
hornum — 233
hygrometricum — 233.
palustre — 233
proliferum — 235
punctatum — 234
purpureum — 233
triguetrum — 235
undulatum. — 235.

BRYA.

- androgynum — 238
apolarpum — 236
argentatum — 243
barbatum — 237
calcareum — 243
cerviculatum — 237
cirriatum — 238
convolutum — 237
crispum — 236
decipiens — 239
extinctorum — 239
glaucum — 239
heteromallum — 240
hypnoides. — 242
lanceolatum — 240
mureale — 240

- ovatum — 240
pellucidum — 237
poniforme — 240
pulvinatum — 243
pyriforme — 241
rigidum — 241
rurale. — 241
scoparium — 241
striatum — 236
simplic — 239
subulatum — 242
trifarium — 238
truncatulum — 242
undulatum — 243
unguiculatum — 242
virrens. — 243.

HYPNA.

- abretinum — 246
adiantoides — 244
aduncum — 248
alopecurum — 249
bryoides — 244
complanatum — 244
crista-castrensis 247
cypresiforme — 248
cuspitatum — 250
dendroides — 250
filicinum — 247
fluitans — 245
lutescens — 245

Jungermannia

- ngosuroides — 251.
parietinum — 247
plumosum — 247
prolongum — 248
proliferum — 248
parum — 250
riparium — 250
rotundifolium — 245
muscifolium — 246
ratibulum — 245
sciroides — 251
scorpioides — 249
sericeum — 251
serpens — 251
squarrosum — 249
stellatum — 246
taxifolium — 244
triguetrum — 246
velutinum — 251
viticulosa. — 252.

Sowerby.

- p. 43. Borage for 217. read 36
p. 49. Campanula glomerata for Sowerby 290. read 90.
p. 153. Genista tinctoria for Sowerby 94 read 44
p. 147. Convolvulus ^{arvensis} for Sowerby 313. read 312
p. 148. Convolvulus ^{sepium} for Sowerby 312 read 313
p. 197. Chara vulgaris for Sowerby. 236 read 336

H. T. Sowerby's further
Desiderata not in this work.

Lichen	uncinatus	with 9 page 2.
	foliatus	— 3
	lacteus	— 5
	atro-virens	— 13
	rupicola	— 13
	compositus	— 13
	rufus	— 70.
	candicans	— 17
	subimbricatus	— 18
	frustulosus	— 19
	carneus	— 20
	cupularis	— 22
	crenulatus	— 17
	leucocephalus	— 20 ?
	psora	— 26.
	luridus	— 28
	diffusus	— 32
	fulous	— 33
	marginalis?	— 34

1225
44
—
1269